BASIS STRUCTURE OF MODULES

O. T. O'MEARA

1. Introduction. Elementary divisor theory over the rational integers is classical; a treatment over an arbitrary algebraic number field can be found in Steinitz [8]. In this paper we are motivated by the following interpretation of Chevalley [4] and Asano [2]: if L_1 and L_2 are finite modules in a vector space V over a Dedekind ring $\mathfrak{o}$ with quotient field F , then L_1 and L_2 have a minimal basis in common. Simple examples show that in general this is not true for more than two modules L_1 and L_2 , but in Theorem 4.2 we establish conditions under which every module in V has a minimal basis in common with a given set of modules $L_0, \dots, L_m$. The basis structure of modules under a direct sum decomposition of V is examined in Theorem 4.3.

The results outlined above are proved over any valuation ring $\mathfrak{o}$: that they can be extended to an arbitrary Dedekind ring is a consequence of Theorem 5.4 which reduces the problem to a local one.

2. Preliminaries. Throughout this paper $\mathfrak{o}$ will denote an integral domain with a unit and with quotient field F ; further restrictions will be placed on $\mathfrak{o}$ in the following paragraphs. The modules L considered will be torsion-free ($\alpha X = 0$ with $\alpha \in \mathfrak{o}$, $X \in L$, implies $\alpha = 0$ or $X = 0$), finitely generated, and of finite rank; hence [2; 4; 6] each L has a replica in a finite dimensional F -space V ; we will always assume that the modules under consideration are contained in such a vector space V ; n will denote the dimension of this vector space; the subspace of V that is spanned by L will be written $F \cdot L$ and we have $\text{rank } L = \dim F \cdot L \leq \dim V = n$.

In addition all modules will have at least one minimal basis in V —that is, a basis $\{\xi\}$ for V and (possibly zero) ideals $\mathfrak{a}_i$ in F such that $L = \mathfrak{a}_1 \xi_1 + \dots + \mathfrak{a}_n \xi_n$; such a basis is known to exist for finitely generated modules in the following case:

THEOREM 2.1. *Let F be such that every finitely generated nonzero $\mathfrak{o}$ -ideal is invertible. If L is a finitely generated $\mathfrak{o}$ -module in V and if $\xi_1, \dots, \xi_n$ is any basis for V , then there exist (finitely generated) ideals $\mathfrak{a}_i$ and a basis $\eta_\lambda = c_{1\lambda} \xi_1 + \dots + c_{\lambda-1\lambda} \xi_{\lambda-1} + \xi_\lambda$ for V such that $L = \sum \mathfrak{a}_\lambda \eta_\lambda$.*

This is proved in Satz 1 of [2] and the proof found there, though given for Dedekind rings, applies to fields of the above type; it is not necessary to assume that $\text{rank } L = n$. Note that the number of nonzero $\mathfrak{a}_i$ is equal to the rank of L . Compare the proofs in [4; 6].

Received by the editors September 23, 1954.

If we say that a result $P(n, m)$, where n and m are non-negative integers, is proved by double induction we mean

$$(2.1) \quad P(0, m) \text{ and } P(n, 0) \text{ hold for all } n \text{ and } m,$$

$$(2.2) \quad P(n, m-1) \text{ and } P(n-1, m) \Rightarrow P(n, m).$$

One notational matter: we will denote the running indices in a summation by Greek letters, fixed indices by Roman or Gothic letters.

3. Minimal bases for L . In this section we investigate which bases in V qualify as minimal bases for L . The ring $\mathfrak{o}$ is quite general but it is assumed that L is a direct sum of $\mathfrak{o}$ -ideals of F ,

$$(3.1) \quad L = a_1\xi_1 + \cdots + a_n\xi_n.$$

Let us introduce the $\mathfrak{o}$ -module

$$(3.2) \quad M = b_1\eta_1 + \cdots + b_n\eta_n,$$

the b_i being $\mathfrak{o}$ -ideals and $\{\eta\}$ being another basis for V connected with $\{\xi\}$ by the equations

$$(3.3) \quad \eta_i = \sum a_{\lambda i} \xi_{\lambda}, \quad \xi_j = \sum b_{\lambda j} \eta_{\lambda}.$$

LEMMA 3.1. $L \subseteq M$ if and only if $b_{ij}a_j \subseteq b_i$; $L = M$ if and only if $b_{ij}a_j \subseteq b_i$ and $a_{ij}b_j \subseteq a_i$ for all i and j .

PROOF. It suffices to prove the first part of the lemma. Substituting (3.3) in (3.1) we see that $L \subseteq M$ is equivalent to $\sum a_{\lambda i} (\sum b_{\lambda \mu} \eta_{\mu}) \subseteq \sum b_{\lambda} \eta_{\lambda}$; that is, $a_j b_{ij} \subseteq b_i$ for all i and j .

LEMMA 3.2. The basis $\{\eta\}$ is a minimal basis for L if and only if $a_{ij}b_{jk}a_k \subseteq a_i$ for all i, j, k .

PROOF. The necessity is an immediate consequence of Lemma 3.1: for if $a_{ij} \neq 0$, then $b_{jk}a_k \subseteq b_j \subseteq a_i a_{ij}^{-1}$. To prove the sufficiency we introduce the ideals $b_i = \sum b_{i\lambda} a_{\lambda}$ and then define M by means of equation (3.2). Then $L = M$ (again by Lemma 3.1) and so $\{\eta\}$ is a minimal basis for L .

LEMMA 3.3. If $L = M$ then $a_{ij}b_j \subseteq a_i$ and $\det \{a_{ij}\} \prod b_{\lambda} = \prod a_{\lambda}$; conversely, if these conditions are satisfied and if the b_i are invertible, then $L = M$.

PROOF. Necessity. It follows from Lemma 3.1 and the expansion $\det \{a_{ij}\} = \sum \pm a_{1\alpha} a_{2\beta} \cdots a_{n\omega}$ that $\det \{a_{ij}\} \prod b_{\lambda} \subseteq \prod a_{\lambda}$. Similarly $\prod b_{\lambda} \supseteq \det \{b_{ij}\} \prod a_{\lambda}$. The remark $\det \{a_{ij}\} = (\det \{b_{ij}\})^{-1}$ completes the proof of the necessity.

Sufficiency. Let a'_{rs} denote the cofactor of a_{rs} . Then $a'_{rs} = \sum \pm a_{1\alpha} \cdots a_{n\omega}$, the first subscript avoiding r , the second s . Hence $a'_{rs}a_{rs}^{-1} \subseteq \prod a_{\lambda} b_{\lambda}^{-1} = \det \{a_{ij}\}$ and so $b_{rs}a_{rs} \subseteq b_s$. The result follows from Lemma 3.1.

REMARK. It follows from the previous lemma that if L and M are isomorphic, then $\prod a_{\lambda}$ and $\prod b_{\lambda}$ are in the same ideal class. Compare [6].

LEMMA 3.4. Suppose that $L = M$ and let the bases $\{\xi\}$ and $\{\eta\}$ have the property that $F\xi_{m+1} + \cdots + F\xi_n = F\eta_{m+1} + \cdots + F\eta_n$ for some m , $0 \leq m \leq n-1$. Then $a_{m+1}\xi_{m+1} + \cdots + a_n\xi_n = b_{m+1}\eta_{m+1} + \cdots + b_n\eta_n$.

PROOF. The given restrictions ensure that all entries a_{ij} and b_{ij} are zero if $i \leq m$ and $j \geq m+1$. Then $\{a_{ij}\}_{i,j \geq m+1}$ has $\{b_{ij}\}_{i,j \geq m+1}$ as its inverse. But a_{ij} and b_{ij} satisfy the conditions of Lemma 3.1 for all i, j and hence for $i, j \geq m+1$. Q.E.D.

We conclude this section with some remarks on the multiplication of modules by nonzero scalars $t \in F$. (a) $t \cdot L$ is again an $\mathfrak{o}$ -module in V . (b) If $\{L_{\lambda}\}$ is a collection of $\mathfrak{o}$ -modules in V and if $\{t_{\lambda}\}$ is a corresponding collection of scalars, then the L_{λ} have a basis $\{\xi\}$ in common if and only if all the $t_{\lambda} \cdot L_{\lambda}$ have the same minimal basis in common. (c) If $\mathfrak{o}$ is a valuation ring or a Dedekind ring, there are nonzero scalars t_1, t_2 such that $t_1 \cdot L \subseteq M$ and $t_2 \cdot L \supseteq M$, provided that L and M are both of rank n (Lemma 3.1).

4. Local theory. The ring $\mathfrak{o}$ is now taken to be a valuation ring with quotient field F : $\mathfrak{o}$ is an integral domain such that for any α in F , either $\alpha \in \mathfrak{o}$ or $\alpha^{-1} \in \mathfrak{o}$. Associated with $\mathfrak{o}$ is a valuation $|\alpha|$, $\alpha \in F$, with values in an ordered multiplicative group [1; 7]. We denote the maximal prime ideal in $\mathfrak{o}$ with the letter $\mathfrak{p}$. Observe that all finitely generated $\mathfrak{o}$ -ideals are principal and hence invertible. All modules will be finitely generated and hence of the form of equation (3.1) with the a_i principal by Theorem 2.1. Hence there is a basis for L in which

$$(4.1) \quad L = (p_1)\xi_1 + \cdots + (p_n)\xi_n, \quad p_i \in F;$$

putting $\xi'_i = p_i\xi_i$ we see that we can take $L = \sum \mathfrak{o}\xi'_i$ if desired. Furthermore, if $L = \sum b_{\lambda}\eta_{\lambda}$ is another representation in a minimal basis $\{\eta\}$, then the b_{λ} must all be principal.

LEMMA 4.1. Let $L_j = \sum a_{\lambda j}\xi_{\lambda} = \sum b_{\lambda j}\eta_{\lambda}$ be modules in V , $0 \leq j \leq m$, and let $L_0 = \sum \mathfrak{o}\xi_{\lambda} = \sum \mathfrak{o}\eta_{\lambda}$. Then there is a reordering of the $\{\eta\}$ in which $a_{ij} = b_{ij}$ for all i, j .

PROOF. By induction to n . In the notation of (3.3), $\{a_{ij}\}$ must be

unimodular. Now $\sum a_{1\lambda}b_{\lambda 1}=1$; therefore there is a k for which $|a_{1k}|=1=|b_{k1}|=|a'_{1k}|$. Interchanging η_1 and η_k we have

$$(4.2) \quad |a_{11}| = 1 = |b_{11}|.$$

Then $a_{11}b_{1j} \subseteq a_{1j}$ by Lemma 3.1 and so $b_{1j} \subseteq a_{1j}$. Similarly $a_{1j} \subseteq b_{1j}$. Hence

$$(4.3) \quad a_{1j} = b_{1j} \quad \text{for all } j.$$

Now $a_{1\lambda}b_{\lambda j} \subseteq a_{1j}$ together with (4.2) and (4.3) implies that $a_{1\lambda}a_{11}^{-1}b_{\lambda j} \subseteq b_{1j}$ and therefore $\sum b_{\lambda j}\eta_\lambda = b_{1j}\eta_1 + \sum b_{\lambda j}(\eta_\lambda - a_{1\lambda}a_{11}^{-1}\eta_1)$. But $(\eta_\lambda - a_{1\lambda}a_{11}^{-1}\eta_1) \in F\xi_2 + \cdots + F\xi_n$ for $\lambda \geq 2$; hence $\sum b_{\lambda j}(\eta_\lambda - a_{1\lambda}a_{11}^{-1}\eta_1) = \sum a_{\lambda j}\xi_\lambda$, with $\lambda \geq 2$, in virtue of Lemma 3.4. Induction completes the proof.

NOTE. In the case $m=1$ the previous lemma expresses the "uniqueness of the elementary divisors."

THEOREM 4.2. *Let L_j , $0 \leq j \leq m$, be given rank n modules in V . Then every rank n module K in V has a minimal basis $\{\xi^K\}$ which is also a minimal basis for all the L_j if and only if either $tL_j \subseteq L_i$ or $tL_j \supseteq L_i$ holds for all i, j and all scalars $t \in F$.*

PROOF. *Sufficiency.* By double induction. (2.1) is true by the elementary divisor theorem which holds for valuation rings. We must prove (2.2).

By the induction and (4.1) we can write $L_0, \dots, L_{m-1}$ and L_m in a common minimal basis such that $L_0 = \sum o\xi_\lambda$, $L_j = \sum (p_{\lambda j})\xi_\lambda$, the $(p_{\lambda j})$ being principal ideals in F . By considering $t_j L_j$ if necessary we can make the further assumption that $L_j \subseteq L_0$ and $L_0 \neq \pi^{-1}L_j \supseteq L_0$ for any $\pi \in \mathfrak{p}$, $0 \leq j \leq m$. Reordering $L_1, \dots, L_m$ we obtain $L_0 \supseteq L_1 \supseteq L_2 \supseteq \dots \supseteq L_m$; hence

$$(4.4) \quad |p_{\lambda m}| \leq |p_{\lambda m-1}| \leq \dots \leq |p_{\lambda 0}| = 1, \quad (p_{\lambda m}) = \mathfrak{p} \text{ or } o.$$

Let $\gamma(\lambda)$ denote the number of p 's with absolute value 1 in this inequality.

Let K be of the form $\sum o\eta_\nu$ with $\xi_i = \sum b_{\mu i}\eta_\mu$. By rearranging $\{\xi\}$ and $\{\eta\}$ we can assume that

$$(4.5) \quad |b_{11}| = \max |b_{i\lambda}|, \quad |b_{1\lambda}| = |b_{11}| \Rightarrow \gamma(1) \geq \gamma(\lambda)'.$$

Now define $H_1 = \sum b_{\lambda 1}b_{11}^{-1}\eta_\lambda$ and $\Xi_\lambda = (\xi_\lambda - b_{1\lambda}b_{11}^{-1}\xi_1)$. Then $(p_{1j})\xi_1 + (p_{\lambda j})\xi_\lambda = (p_{1j})\xi_1 + (p_{\lambda j})\Xi_\lambda$, since $|b_{1\lambda}b_{11}^{-1}| < 1$ implies $|p_{\lambda j}b_{1\lambda}b_{11}^{-1}| \leq |p_{1j}|$, while if $|b_{1\lambda}b_{11}^{-1}| = 1$, then $\gamma(\lambda) \leq \gamma(1)$ implies $|p_{\lambda j}| \leq |p_{1j}|$. Hence $L_j = (p_{1j})\xi_1 + L'_j$ where $L'_j = \sum (p_{\lambda j})\Xi_\lambda$. Since $|b_{\lambda 1}b_{11}^{-1}| \leq 1$, we must have $K = oH_1 + K'$ where $K' = o\eta_2 + \dots + o\eta_n$. But $F\xi_1 = FH_1$

and $\sum F\Xi_\lambda = F\eta_2 + \cdots + F\eta_n$ by definition of H_1 and Ξ_λ . In addition $L'_{j+1} \subseteq L'_j$ and $L'_{j+1} \supseteq \mathfrak{p}L'_j$ by (4.4). Applying the induction again we see that L'_j , $0 \leq j \leq m$, and K' have a minimal basis $\{\omega\}$ in common. Hence $\xi_1, \omega_2, \cdots, \omega_n$ is a minimal basis for L_j , $0 \leq j \leq m$, and K . This proves (2.2) and hence establishes the induction.

Necessity. Given any pair L_i, L_j there is a minimal basis $\{\xi\}$ such that $L_i = \sum \mathfrak{o}\xi_\lambda$ and $L_j = \sum \mathfrak{b}_\lambda \xi_\lambda$ by the elementary divisor theorem. Hence if the given conditions on the L_j , $0 \leq j \leq m$, do not hold there must be a pair, L_0 and L_1 say, such that $L_0 = \sum \mathfrak{o}\xi_\lambda$ and $tL_1 = (p)\xi_1 + \mathfrak{o}\xi_2 + (p_3)\xi_3 + \cdots + (p_n)\xi_n$ with $p_\lambda \in \mathfrak{o}$ and $|p| < |\pi| < 1$, for some $\pi \in \mathfrak{p}$. By a change of notation let us refer to tL_1 as L_1 . Now define a new basis $\{\eta\}$ for V by the equations $\{\eta\}B = \{\xi\}$ where

$$(4.6) \quad \xi_1 = \eta_1, \quad \xi_2 = \pi\eta_1 + p\eta_2, \quad \xi_i = \eta_i, \quad i > 2.$$

Put $K = \sum \mathfrak{o}\eta_\lambda$. Contention: L_0, L_1 and K do not have a common minimal basis.

Suppose that the contrary is true and let their common minimal basis be

$$(4.7) \quad \{\zeta\} = \{\xi\}C = \{\eta\}BC.$$

Put $C = \{c_{ij}\}$ and $C^{-1} = \{d_{ij}\}$. Then the first row and k th column of BC has entry $(c_{1k} + \pi c_{2k})$ while the k th row and second column of $(BC)^{-1}$ has entry $(d_{k2} - \pi d_{k1})p^{-1}$. Now apply Lemma 3.2 (i) to L_0 : $c_{ik}d_{kj} \in \mathfrak{o}$ all i, j, k ; (ii) to L_1 : $c_{1k}d_{k2} \in (p)$ for all k ; (iii) to K : $(c_{1k} + \pi c_{2k})(d_{k2} - \pi d_{k1})p^{-1} \in \mathfrak{o}$. Then these three conditions yield (iv): $c_{1k}d_{k1} - c_{2k}d_{k2} \in \mathfrak{p}$. Now $(c_{1k}d_{k1})(c_{2k}d_{k2}) \in (p)$ by (i) and (ii) and so either $c_{1k}d_{k1} \in \mathfrak{p}$ or $c_{2k}d_{k2} \in \mathfrak{p}$ and hence both must be in $\mathfrak{p}$ by (iv). Hence $1 = \sum c_{1\lambda}d_{\lambda 1} \in \mathfrak{p}$ and this is a contradiction. Q.E.D.

REMARK. There is a different way of stating Theorem 4.2—by assuming that $FL_0 = V$ but dropping the conditions on the ranks of the L_j , $0 \leq j \leq m$, and on K .

Sufficiency. It is easy to see that the conditions of the theorem guarantee that all L_j must be of the same rank and therefore span V . If in addition rank $K = n$, the proof is as before. If this is not so, write $K = \mathfrak{o} \cdot \eta_1 + \mathfrak{a}_2 \eta_2 + \cdots + \mathfrak{a}_n \eta_n$; instead of (4.5) we can arrange to have

$$|b_{11}| = \max |b_{1\lambda}|, \quad |b_{1\lambda}| = |b_{11}| \Rightarrow \gamma(1) \geq \gamma(\lambda)$$

by permuting the $\{\xi\}$ only. The rest of the proof is as before.

Necessity. If all L_j , $0 \leq j \leq m$, are of rank n the previous proof is valid. If not, we can take $L_0 = \sum \mathfrak{o}\xi_\lambda$ and $L_1 = \mathfrak{o} \cdot \xi_1 + \mathfrak{o}\xi_2 + \mathfrak{a}_3\xi_3 + \cdots + \mathfrak{a}_n\xi_n$ by the elementary divisor theorem; put $K = \mathfrak{o}(\xi_1 + \pi^{-1}\xi_2)$ where

$\pi \in \mathfrak{p}$; then it is easy to see (e.g. by using Theorem 2.1 and Lemma 3.4) that L_0 , L_1 and K do not have a common minimal basis.

THEOREM 4.3. *If L_j , $0 \leq j \leq m$, are modules in V having a minimal basis in common and if each L_j has a decomposition $L_j = L'_j + L''_j$ corresponding to a fixed direct sum decomposition $V = V' + V''$, then the L'_j (resp. L''_j) have a common minimal basis in V' (resp. V'').*

First we prove the following special case.

LEMMA. *The theorem is true if $FL_0 = V$.*

PROOF OF LEMMA. By double induction. (2.1) is true. We must prove (2.2). By the induction we have

$$(4.8) \quad L'_j = \sum_1^q (p_{\lambda j}) \xi_\lambda \subseteq V', \quad L''_j = \sum_{q+1}^n (p_{\lambda j}) \xi_\lambda \subseteq V'',$$

$$0 \leq j \leq m-1,$$

with $L_0 = \sum \mathfrak{o} \xi_\lambda$, that is $p_{\lambda 0} = 1$ for all λ . Let $\{\zeta\}$ be a minimal basis common to all the L_j , $0 \leq j \leq m$; there is no loss of generality in taking

$$(4.9) \quad L_0 = \sum \mathfrak{o} \zeta_\lambda, \quad L_j = \sum (\bar{p}_{\lambda j}) \zeta_\lambda, \quad L_m = \mathfrak{o} \zeta_1 + \sum_2^n (\bar{p}_{\lambda m}) \zeta_\lambda,$$

$$\bar{p}_{\lambda m} \in \mathfrak{o}.$$

If $\zeta_i = \sum c_{\lambda i} \xi_\lambda$, then $\{c_{ij}\}$ must be unimodular and we can permute $\{\xi\}$ as in Lemma 4.1 until $|c_{11}| = 1 = |d_{11}|$, interchange V' and V'' if necessary, and then permute $\zeta_2, \dots, \zeta_n$ until

$$(4.10) \quad \sum (p_{\lambda j}) \xi_\lambda = L_j = \sum (p_{\lambda j}) \zeta_\lambda, \quad 0 \leq j \leq m-1,$$

$$(4.11) \quad |c_{11}| = 1, \quad c_{\lambda \mu} p_{\mu j} \in (p_{\lambda j}), \quad 0 \leq j \leq m-1.$$

Note that with a corresponding change of notation, (4.8) and (4.9) are preserved.

Now put $Z_1 = \zeta_1$, $Z_\lambda = (\zeta_\lambda - c_{1\lambda} c_{11}^{-1} \zeta_1)$, $\Xi_1 = \sum_1^q c_{\lambda 1} c_{11}^{-1} \xi_\lambda$, $\Xi_\lambda = \xi_\lambda$ and then write ζ and ξ instead of Z and Ξ . In virtue of (4.11) and these substitutions, (4.8), (4.9) and (4.10) are still valid and

$$(4.12) \quad \zeta_1 = c_{11} \xi_1 + \sum_{q+1}^n c_{\lambda 1} \xi_\lambda, \quad \zeta_i = \sum_2^n c_{\lambda i} \xi_\lambda, \quad i > 1.$$

It follows from the direct sum decomposition of L_m and (4.9) that $\sum_{q+1}^n c_{\lambda 1} \xi_\lambda \in L_m$ and $c_{11} \xi_1 \in L_m$. Hence we see from (4.9) and (4.12) that $L_m = \sum_2^n (\bar{p}_{\lambda m}) \zeta_\lambda + \mathfrak{o} \xi_1$. Now $\sum_2^n (\bar{p}_{\lambda m}) \zeta_\lambda$ and $\sum_2^n (p_{\lambda j}) \xi_\lambda$, $0 \leq j \leq m-1$, have $\zeta_2, \dots, \zeta_n$ as a common minimal basis in virtue of Lemma 3.4 and (4.12). Contention: there is a direct sum decomposition of $\sum_2^n (\bar{p}_{\lambda m}) \zeta_\lambda$ corresponding to $\sum_2^q F \xi_\lambda \oplus \sum_{q+1}^n F \xi_\lambda$: for $L_m = L'_m + L''_m$;

and $L'_m = 0\xi_1 + \sum_2^q a_\lambda \eta_\lambda$ with $\eta_\lambda = \alpha_{1\lambda}\xi_1 + \cdots + \xi_\lambda$ in virtue of Theorem 2.1; but $a_\lambda \alpha_{1\lambda} \subseteq 0$ since $L_m \subseteq L_0$; hence we can assume that $\alpha_{1\lambda} = 0$, $2 \leq \lambda \leq q$; then by Lemma 3.4, $\sum_2^n (\bar{p}_{\lambda m})\zeta_\lambda = \sum_2^q a_\lambda \eta_\lambda \oplus L''_m$. This proves the contention. Induction completes the proof. Q.E.D.

PROOF OF THEOREM. By double induction. By applying the elementary divisor theorem to L'_0 (resp. L''_0) and L'_1 (resp. L''_1) we can write

$$(4.13) \quad \begin{aligned} L_0 &= 0\xi_1 + \cdots + 0\xi_r + \cdots + 0\xi_t, \\ L_1 &= \quad \quad \quad p_r\xi_r + \cdots + p_t\xi_t + 0\xi_{t+1} + \cdots + 0\xi_s \end{aligned}$$

with $s \leq n$ and $\{\xi\}$ such that either $\xi_\lambda \in V'$ or $\xi_\lambda \in V''$; there is no loss of generality in assuming that $p_\lambda \subseteq 0$, $r \leq \lambda \leq t$. On the other hand an easy computation shows that we can assume that equations (4.13) are true if we replace $\{\xi\}$ by $\{\zeta\}$ where $\{\zeta\}$ is a minimal basis common to all L_j , $0 \leq j \leq m$. Define K invariantly as the smallest 0 -module containing L_0 and L_1 . Then

$$\sum_1^s 0\xi_\lambda = K = \sum_1^s 0\zeta_\lambda.$$

CASE I. $F \cdot K = V$. Then $K, L_0, L_1, \cdots, L_m$ satisfy the conditions of the lemma; the theorem therefore holds for these modules; hence it holds for L_j , $0 \leq j \leq m$.

CASE II. $F \cdot K \neq V$. By the induction $K, L_2, L_3, \cdots, L_m$ have a common basis $\{\eta\}$ such that $K = \sum_1^s 0\eta_\lambda$ and either $\eta_\lambda \in V'$ or $\eta_\lambda \in V''$ for $1 \leq \lambda \leq n$. By means of Lemma 3.4 we see that L_0, L_1 and $\sum_1^s a_{\lambda j} \eta_\lambda$, $2 \leq j \leq m$, must have $\zeta_1, \cdots, \zeta_s$ as a common minimal basis and hence have a basis $\omega_1, \cdots, \omega_s$ with the required property. Then $\omega_1, \cdots, \omega_s, \eta_{s+1}, \cdots, \eta_n$ is the basis required. This completes the proof of the theorem.

LEMMA 4.4. Let L_j be rank n modules in V and let X be any vector in L_0 . Then there is a nonzero α (independent of X) such that if $\{\xi\}$ is any basis for all L_j with $L_0 = \sum 0\xi_\lambda$, then so is $\xi_1, \cdots, \xi_{i-1}, \xi_i + \alpha'X, \xi_{i+1}, \cdots, \xi_n$ where $|\alpha'| \leq |\alpha|$.

PROOF. Write $L_j = \sum a_{\lambda j} \xi_\lambda$ and $X = \sum \alpha_\lambda \xi_\lambda$ where $\alpha_\lambda \in 0$; choose $\alpha \in \mathfrak{p}$ such that $\alpha \in a_{\lambda j} a_{\mu j}^{-1}$ for all λ, μ, j . Then

$$L_j = a_{1j}\xi_1 + \cdots + a_{ij}(\xi_i + \alpha'X) + \cdots + a_{nj}\xi_n.$$

5. **Dedekind rings.** In this section we assume that classical ideal theory holds in $F/0$, that is to say the ideals with bounded denominator form a group under multiplication. We denote by $\mathfrak{v}_\mathfrak{p}$ the valuation ring corresponding to the prime ideal $\mathfrak{p}$; the same letter $\mathfrak{p}$ will be used

for the maximal prime ideal in $\mathfrak{o}_{\mathfrak{p}}$. The $\mathfrak{o}_{\mathfrak{p}}$ -ideals in F are all powers of $\mathfrak{p}$ and the value group is discrete and real. If $\mathfrak{a} = \mathfrak{p}^r \cdots$ is a factorization of the $\mathfrak{o}$ -ideal $\mathfrak{a}$ we define $\mathfrak{p}(\mathfrak{a})$ as the $\mathfrak{o}_{\mathfrak{p}}$ -ideal $\mathfrak{p}^r$. All modules will be finitely generated and hence of the form (3.1) with the α_i of bounded denominator and invertible. Note that if $\xi_1, \dots, \xi_n$ is any basis for V , then by Theorem 2.1 there is a minimal basis $\eta_1, \dots, \eta_n$ for L such that

$$(5.1) \quad \eta_i = c_{1i}\xi_1 + \cdots + c_{i-1i}\xi_{i-1} + \xi_i, \quad c_{ij} \in F.$$

If $L = \sum \alpha_{\lambda} \xi_{\lambda}$ put $\mathfrak{p}(L) = \sum \mathfrak{p}(\alpha_{\lambda}) \xi_{\lambda}$: that this is well-defined is an immediate consequence of Lemma 3.1. By taking $\xi'_{\lambda} = \alpha_{\lambda} \xi_{\lambda}$ instead of ξ_{λ} it is always possible to choose a basis for L such that $\mathfrak{p}(L) = \sum \mathfrak{o}_{\mathfrak{p}} \xi'_{\lambda}$ at a given finite set of primes $\mathfrak{p}$ provided that $\text{rank } L = n$. By means of Lemma 3.1 and the unique factorization into primes it is easy to prove the following result.

LEMMA 5.1. *$L = M$ if and only if $\mathfrak{p}(L) = \mathfrak{p}(M)$ at all $\mathfrak{p}$. In any case, if $F \cdot L = F \cdot M$, then $\mathfrak{p}(L) = \mathfrak{p}(M)$ holds at all but a finite number of primes $\mathfrak{p}$.*

LEMMA 5.2. *Let L_j , $0 \leq j \leq m$, be rank n modules in V and let W be an r dimensional subspace of V . If $\mathfrak{p}(L_j)$, $0 \leq j \leq m$, have a common minimal basis $\{\eta^{\mathfrak{p}}\}$ such that $\eta_1^{\mathfrak{p}}, \dots, \eta_r^{\mathfrak{p}} \in W$ at each $\mathfrak{p}$ in a finite set of primes S , then there is a basis $\{\xi\}$ for L_0 which is also a basis for all the $\mathfrak{p}(L_j)$, $\mathfrak{p} \in S$, and such that $\xi_1, \dots, \xi_r \in W$.*

PROOF. By induction on n . All bases (whether defined or derived) that appear in the proof will be such that their first r elements span W . Let us write $L_0 = \sum \alpha_{\lambda 0} \xi_{\lambda}$ and $\eta_i^{\mathfrak{p}} = \sum \alpha_{\lambda i}^{\mathfrak{p}} \xi_{\lambda}$; we can assume that $\mathfrak{p}(\alpha_{\lambda 0}) = \mathfrak{o}_{\mathfrak{p}}$ and $\mathfrak{p}(L_0) = \sum \mathfrak{o}_{\mathfrak{p}} \eta_{\lambda}^{\mathfrak{p}}$ at all $\mathfrak{p} \in S$. Apply the approximation theorem to the entries of the matrices $\{a_{ij}^{\mathfrak{p}}\}$ in such a way that a_{ij} approximates to $a_{ij}^{\mathfrak{p}}$ with $a_{ij} = 0$ whenever $a_{ij}^{\mathfrak{p}} = 0$ is true for all $\mathfrak{p} \in S$; then in virtue of Lemma 4.4 a sufficiently high approximation yields a basis $\eta_j = \sum \alpha_{\lambda j} \xi_{\lambda}$ such that

$$(5.2) \quad \mathfrak{p}(L_0) = \sum \mathfrak{o}_{\mathfrak{p}} \xi_{\lambda}, \quad \mathfrak{p}(L_j) = \sum \mathfrak{p}_{\lambda j} \eta_{\lambda}, \quad \text{all } \mathfrak{p} \in S.$$

Now write each $L_j = \sum \alpha_{\lambda j} \xi'_{\lambda j}$ with $\xi'_{\lambda j} \in F\eta_1 + \cdots + F\eta_{\lambda}$, by (5.1). Then all the modules $\sum_{i=1}^{n-1} \alpha_{\lambda j} \xi'_{\lambda j}$ have $\eta_1, \dots, \eta_{n-1}$ as a common minimal basis at each $\mathfrak{p} \in S$, in virtue of Lemma 3.4; by the induction there is a basis (and we continue to call it $\{\xi\}$) for L_0 such that $\eta_1, \dots, \eta_n$ is a basis for the $\mathfrak{p}(L_j)$, $\mathfrak{p} \in S$, where $\eta_i = \xi_i$ for all $i \leq n-1$; we can assume that

$$(5.3) \quad \sum \mathfrak{o}_{\mathfrak{p}} \xi_{\lambda} = \mathfrak{p}(L_0) = \sum \mathfrak{o}_{\mathfrak{p}} \eta_{\lambda}, \quad \mathfrak{p} \in S.$$

Suppose that $\eta_n = \alpha_1 \xi_1 + \cdots + \alpha_n \xi_n$; then $|\alpha_i| \leq 1$ and $|\alpha_n| = 1$ at all $\mathfrak{p} \in S$; by suitably scaling η_n we can make the further assumptions

$$(5.4) \quad \alpha_i \in \mathfrak{o}, \quad \alpha_i \in \mathfrak{a}_{i0} \mathfrak{a}_{n0}^{-1} \quad \text{for all } i.$$

Let $\alpha, \alpha' \in \mathfrak{o}$ be such that $|\alpha|, |\alpha'|$ are arbitrarily small at $\mathfrak{p} \in S$, but either $|\alpha| = 1$ or $|\alpha'| = 1$ outside S ; then $\text{g.c.d.}(\alpha, \alpha', \alpha_n) = \mathfrak{o}$ and there must exist $\beta, \beta', \gamma \in \mathfrak{o}$ such that $\alpha\beta + \alpha'\beta' + \gamma\alpha_n = 1$; clearly γ must be a unit at all $\mathfrak{p} \in S$. Define

$$(5.5) \quad \zeta_i = \xi_i, \quad i < n, \quad \zeta_n = \alpha\beta\xi_n + \alpha'\beta'\xi_n + \gamma\eta_n.$$

Then $\zeta_n = \sum_{\lambda=1}^{n-1} \gamma_{\lambda} \xi_{\lambda} + \xi_n$ and so $\{\zeta\}$ is a basis for L_0 . Since $|\alpha|, |\alpha'|$ can be made arbitrarily small at $\mathfrak{p} \in S$, we have $\{\zeta\}$ as a basis for $\mathfrak{p}(L_j)$, $\mathfrak{p} \in S$, by Lemma 4.4. Q.E.D.

LEMMA 5.3. *If the assumptions of the previous lemma are satisfied at ALL primes $\mathfrak{p}$, then there is a common basis $\{\zeta\}$ for all L_j , $0 \leq j \leq m$, such that $\zeta_1, \cdots, \zeta_r \in W$.*

PROOF. Let S be a finite set of primes such that $\mathfrak{p}(L_0) = \mathfrak{p}(L_j)$ if $\mathfrak{p} \notin S$. Let $\{\zeta\}$ be a basis having the properties of Lemma 5.2; then $\mathfrak{p}(L_0) = \sum \mathfrak{p}_{\lambda 0} \zeta_{\lambda} = \mathfrak{p}(L_j)$ if $\mathfrak{p} \notin S$, and $\mathfrak{p}(L_j) = \sum \mathfrak{p}_{\lambda j} \zeta_{\lambda}$ if $\mathfrak{p} \in S$. Define

$$\mathfrak{a}_{\lambda j} = \prod_{\mathfrak{p} \in S} \mathfrak{p}_{\lambda j} \prod_{\mathfrak{p} \notin S} \mathfrak{p}_{\lambda 0}.$$

Then $L_j = \sum \mathfrak{a}_{\lambda j} \zeta_{\lambda}$ by Lemma 5.1.

THEOREM 5.4. *The modules L_j have a common minimal basis if and only if their local components $\mathfrak{p}(L_j)$ have a common minimal basis at all primes $\mathfrak{p}$, $0 \leq j \leq m$.*

PROOF. By double induction. By the elementary divisor theorem

$$L_0 = \mathfrak{a}_{10}\omega_1 + \cdots + \mathfrak{a}_{r0}\omega_r + \cdots + \mathfrak{a}_{q0}\omega_q,$$

$$L_1 = \mathfrak{a}_{r1}\omega_r + \cdots + \mathfrak{a}_{q1}\omega_q + \cdots + \mathfrak{a}_{s1}\omega_s.$$

We can assume that $\mathfrak{a}_{\lambda 1} \subseteq \mathfrak{a}_{\lambda 0}$, $r \leq \lambda \leq q$. Define K invariantly as the $\mathfrak{o}$ -module generated by L_0 and L_1 ; then $K = \sum (\mathfrak{a}_{\lambda 0} + \mathfrak{a}_{\lambda 1}) \omega_{\lambda}$.

CASE I. Rank $L_j = n$, $0 \leq j \leq m$. This follows from Lemma 5.3.

CASE II. Rank $L_0 < n$, rank $L_j = n$, $1 \leq j \leq m$. By Lemma 5.3 we can express L_j , $1 \leq j \leq m$, in a common $\{\xi\}$ with $\xi_1, \cdots, \xi_q$ spanning $F \cdot L_0$: $L_j = \sum \mathfrak{a}_{\lambda j} \xi_{\lambda}$, $1 \leq j \leq m$. Then L_0 and $\sum \mathfrak{a}_{\lambda j} \xi_{\lambda}$ have a common basis at all $\mathfrak{p}$ in virtue of Lemma 3.4 and so by the induction they must have a common basis $\zeta_1, \cdots, \zeta_q$ in the large. Then $\zeta_1, \cdots, \zeta_q, \zeta_{s+1}, \cdots, \zeta_n$ is the basis required.

CASE III. Rank $K = s < n$. Then if $\{\eta^p\}$ is a basis for $\mathfrak{p}(L_j)$, $0 \leq j \leq m$, it must also be a basis for $\mathfrak{p}(K)$; therefore by the induction $K, L_2, \dots, L_m$ have a common basis $\{\xi\}$; let us suppose that $\xi_1, \dots, \xi_s$ spans $F \cdot K$; then L_0, L_1 and $\sum_1^s \alpha_{\lambda j} \xi_\lambda$ have a common minimal basis at all $\mathfrak{p}$ in virtue of Lemma 3.4; hence they have a basis $\zeta_1, \dots, \zeta_s$ in the large. Then $\zeta_1, \dots, \zeta_s, \xi_{s+1}, \dots, \xi_n$ is the basis required.

CASE IV. Rank $L_0 < n$, rank $L_1 < n$ and rank $K = s = n$. Then by (5.1) we can express each L_j , $2 \leq j \leq m$, in a basis whose successive elements are in $F\omega_n, F\omega_n + F\omega_{n-1}$, etc.; by Lemma 3.4 and the inductive hypothesis we see that we can assume that $L_j = \sum_1^{r-1} \alpha_{\lambda j} \omega'_{\lambda j} + \sum_r^n \alpha_{\lambda j} \omega_\lambda$, for $0 \leq j \leq m$.

Contention: we can arrange to have $\omega'_{\lambda j} \in FL_0$ when $2 \leq j \leq m$. For each $\alpha \in \alpha_{\lambda j}$ we have $\alpha \omega'_{\lambda j} = \alpha \omega_\lambda + \alpha \rho_{\lambda+1 j} \omega_{\lambda+1} + \dots + \dots \in L_j$; at each $\mathfrak{p}$, $\mathfrak{p}(L_j)$ has a basis $\{\eta^p\}$ with $\eta_1^p, \dots, \eta_q^p \in FL_0$, $\eta_r^p, \dots, \eta_n^p \in FL_1$; and $\eta_\mu^p = \omega_\mu$ for $\mu \geq r$, by Lemma 3.4, hence

$$\alpha \omega'_{\lambda j} = \beta_1 \eta_1^p + \dots + \beta_q \eta_q^p + \alpha \rho_{q+1 j} \omega_{q+1} + \dots + \alpha \rho_n j \omega_n;$$

hence $\alpha \rho_{\mu j} \omega_\mu \in \mathfrak{p}(L_j)$ holds for all $\mathfrak{p}$, $\mu \geq q+1$; and so $\alpha_{\lambda j} \rho_{\mu j} \omega_\mu \subseteq L_j$. This proves the contention.

By the induction and Lemma 3.4 we can express $\sum_1^{r-1} \alpha_{\lambda j} \omega'_{\lambda j} + \sum_r^n \alpha_{\lambda j} \omega_\lambda$, $0 \leq j \leq m$, in a common minimal basis $\zeta_1, \dots, \zeta_q$. Then $\zeta_1, \dots, \zeta_q, \omega_{q+1}, \dots, \omega_n$ is the basis required. We have exhausted all possibilities and the theorem is now proved.

REFERENCES

1. E. Artin, *Algebraic numbers and algebraic functions*, Princeton University, New York University, 1950.
2. K. Asano, *Über Moduln und Elementarteilertheorie im Körper, in dem Arithmetik definiert ist*, Jap. J. Math. vol. 20 (1950) pp. 55–71.
3. N. Bourbaki, *Éléments de mathématique*, Livre II, Actualités Scientifiques et Industrielles, no. 1179, Paris, 1952.
4. C. Chevalley, *L'arithmétique dans les algèbres de matrices*, Actualités Scientifiques et Industrielles, no. 323, Paris, 1936.
5. I. Kaplansky, *Elementary divisors and modules*, Trans. Amer. Math. Soc. vol. 66 (1949) pp. 464–491.
6. ———, *Modules over Dedekind rings and valuation rings*, Trans. Amer. Math. Soc. vol. 72 (1952) pp. 327–340.
7. W. Krull, *Allgemeine Bewertungstheorie*, J. Reine Angew. Math. vol. 167 (1932) pp. 160–196.
8. E. Steinitz, *Rechteckige Systeme und Moduln in algebraischen Zahlkörpern*, Math. Ann. vol. 71 (1912) pp. 328–354.

UNIVERSITY OF OTAGO, NEW ZEALAND