EVALUATION OF A TRIGONOMETRIC INTEGRAL

R. MOHANTY

1. Introduction. It is assumed throughout the present paper that $f(t) \in L(-\infty, \infty)$. Then the Fourier integral of $f(t)$ at $t=x$ is

$$(1) \quad \frac{1}{\pi} \int_0^\infty du \int_{-\infty}^\infty f(t) \cos u(x-t) dt.$$

This may be written in a form analogous to that of the Fourier series as

$$(2) \quad \int_0^\infty \{a(u) \cos xu + b(u) \sin xu\} du,$$

where

$$(3) \quad a(u) = \frac{1}{\pi} \int_{-\infty}^\infty f(t) \cos ut dt, \quad b(u) = \frac{1}{\pi} \int_{-\infty}^\infty f(t) \sin ut dt.$$

The allied integral of the Fourier integral is then

$$(4) \quad \begin{aligned} \int_0^\infty \{b(u) \cos xu - a(u) \sin xu\} du \\ = \frac{1}{\pi} \int_0^\infty du \int_{-\infty}^\infty f(t) \sin u(t-x) dt. \end{aligned}$$

Thus the allied integral of an odd function $f(t)$ belonging to $L(0, \infty)$ is

$$(5) \quad \frac{2}{\pi} \int_0^\infty \cos xudu \int_0^\infty f(t) \sin ut dt.$$

We write

$$(6) \quad \psi(t) = f(x+t) - f(x-t).$$

DEFINITION. We say that the integral $\int_0^\infty g(u)du$ is summable $(C, 1)$ to sum I , if

$$\lim_{\lambda \rightarrow \infty} \int_0^\lambda \left(1 - \frac{u}{\lambda}\right) g(u) du = I.$$

In this paper we are concerned with the integrals of the type

Received by the editors December 19, 1955.

$$(7) \quad \int_0^\infty u^\alpha \{b(u) \cos xu - a(u) \sin xu\} du,$$

where $0 < \alpha < 1$, and first prove the following

THEOREM. *Integral (7) is summable $(C, 1)$ to*

$$\frac{\Gamma(1 + \alpha)}{\pi} \cos \frac{1}{2} \alpha \pi \int_{-\infty}^\infty \frac{\psi(t)}{t^{1+\alpha}} dt,$$

whenever this integral exists, and

$$(8) \quad \psi(t) = o(t^\alpha) \text{ as } t \rightarrow 0.$$

2. **Proof of the theorem.** We are to consider the behaviour of

$$\begin{aligned} I &= \frac{1}{\pi} \int_0^\lambda \left(1 - \frac{u}{\lambda}\right) u^\alpha du \int_{-\infty}^\infty f(t) \sin u(t - x) dt \\ &= \frac{1}{\pi} \int_0^\lambda \left(1 - \frac{u}{\lambda}\right) u^\alpha du \int_0^\infty \psi(t) \sin ut dt \end{aligned}$$

as $\lambda \rightarrow \infty$. Now

$$\int_0^\lambda \left(1 - \frac{u}{\lambda}\right) u^\alpha du \int_0^\infty \psi(t) \sin ut dt = \int_0^\infty \psi(t) K(\lambda, t) dt,$$

where

$$\begin{aligned} K(\lambda, t) &= \int_0^\lambda \left(1 - \frac{u}{\lambda}\right) u^\alpha \sin ut du \\ &= \lambda^{1+\alpha} \int_0^1 (\omega^\alpha - \omega^{\alpha+1}) \sin \omega \lambda t d\omega \\ &= \lambda^{1+\alpha} \left[(\omega^\alpha - \omega^{\alpha+1}) \frac{\cos \omega \lambda t}{-\lambda t} \right]_0^1 \\ &\quad + \frac{\lambda^\alpha}{t} \int_0^1 [\alpha \omega^{\alpha-1} - (\alpha + 1) \omega^\alpha] \cos \omega \lambda t d\omega \\ &= \frac{\alpha \lambda^\alpha}{t} \int_0^1 \frac{\cos \omega \lambda t}{\omega^{1-\alpha}} d\omega - \frac{(\alpha + 1) \lambda^\alpha}{t} \int_0^1 \omega^\alpha \cos \omega \lambda t d\omega \\ &= \frac{\alpha}{t^{1+\alpha}} \int_0^{\lambda t} \frac{\cos \theta}{\theta^{1-\alpha}} d\theta + O\left(\frac{1}{\lambda^{1-\alpha} t^2}\right). \end{aligned}$$

Hence

$$K(\lambda, t) - \frac{\alpha}{t^{1+\alpha}} \int_0^\infty \frac{\cos \theta}{\theta^{1-\alpha}} d\theta = \frac{-\alpha}{t^{1+\alpha}} \int_{\lambda t}^\infty \frac{\cos \theta}{\theta^{1-\alpha}} d\theta + O\left(\frac{1}{\lambda^{1-\alpha} t^2}\right).$$

By the second mean value theorem, we have

$$\left| \int_{\lambda t}^\infty \frac{\cos \theta}{\theta^{1-\alpha}} d\theta \right| \leq \frac{2}{\lambda^{1-\alpha} t^{1-\alpha}},$$

and consequently

$$(9) \quad \left| K(\lambda, t) - \frac{\Gamma(\alpha + 1)}{t^{1+\alpha}} \cos \frac{1}{2} \alpha \pi \right| < \frac{A}{\lambda^{1-\alpha} t^2}.$$

Plainly

$$(10) \quad |K(\lambda, t)| < A \lambda^{1+\alpha}.$$

It follows at once that, if $\delta > 0$,

$$\lim_{\lambda \rightarrow \infty} \frac{1}{\pi} \int_\delta^\infty \psi(t) K(\lambda, t) dt = \frac{\Gamma(1 + \alpha)}{\pi} \cos \frac{1}{2} \alpha \pi \int_\delta^\infty \frac{\psi(t)}{t^{1+\alpha}} dt.$$

Also by (9), we have

$$\begin{aligned} \frac{1}{\pi} \int_{\lambda^{-1}}^\delta \psi(t) K(\lambda, t) dt \\ = \frac{\Gamma(1 + \alpha)}{\pi} \cos \frac{1}{2} \alpha \pi \int_{\lambda^{-1}}^\delta \frac{\psi(t)}{t^{1+\alpha}} dt + O \left\{ \int_{\lambda^{-1}}^\delta \frac{|\psi(t)|}{t^2 \lambda^{1-\alpha}} dt \right\} \end{aligned}$$

and, if condition (8) holds, the last term is

$$o \left(\frac{1}{\lambda^{1-\alpha}} \int_{\lambda^{-1}}^\delta t^{\alpha-2} dt \right) = o(1)$$

by choosing first δ then λ .

Finally, by (8) and (10), we have

$$\begin{aligned} \left| \frac{1}{\pi} \int_0^{\lambda^{-1}} \psi(t) K(\lambda, t) dt \right| &< A \lambda^{1+\alpha} \int_0^{\lambda^{-1}} |\psi(t)| dt \\ &= o(1). \end{aligned}$$

This proves the theorem.

3. Evaluation of a definite integral. Let us consider the odd function $e^{-t} t^{\beta-1}$ ($\beta > 1 + \alpha$) defined in $(0, \infty)$, then the integral (7) at $x=0$ reduces for the present function to

$$\begin{aligned} \frac{2}{\pi} \int_0^\infty u^\alpha du \int_0^\infty e^{-t} t^{\beta-1} \sin ut dt \\ = \frac{2}{\pi} \Gamma(\beta) \int_0^\infty \frac{u^\alpha \sin \{\beta \tan^{-1} u\} du}{(1+u^2)^{\beta/2}}, \quad \text{where } \beta > 1 + \alpha. \end{aligned}$$

Obviously this integral is a convergent integral, and since this special function satisfies the conditions of the theorem, we have by the theorem

$$(11) \quad \int_0^\infty \frac{u^\alpha \sin (\beta \tan^{-1} u)}{(u^2 + 1)^{\beta/2}} du = \frac{\Gamma(\alpha + 1) \Gamma(\beta - \alpha - 1)}{\Gamma(\beta)} \cos \frac{1}{2} \alpha \pi,$$

where

$$\beta > \alpha + 1.$$

RAVENSHAW COLLEGE, CUTTACK, INDIA