SOME CONSEQUENCES OF THE APPROXIMATION THEOREM OF BING

O. G. HARROLD, JR.¹

1. Introduction. In the following paragraphs we consider several unsolved problems in the topology of 3-space. Among the results is included a new formulation of properties characterizing tame curves in three-space without any explicit mention of polyhedral sets.

Three-space is denoted by R^3 , the null-set by $\square$, and the set of points in A but not in B by $A \setminus B$. The combinatorial boundary of chain E is ∂E and the set of points on such a boundary by $|\partial E|$. The closure of a set A is denoted by $\text{Cl } A$, or $\overline{A}$.

THEOREM (Bing). *Given a compact 2-manifold M of R^3 with or without boundaries, a compact subset D of R^3 and a positive number ϵ . There is a 2-manifold N and a homeomorphism f of M on N such that no point of M is moved more than ϵ and N is locally polyhedral at each point of $N \setminus D$. If $D \neq \square$, f may also be taken so that no point x is moved more than the minimum of ϵ and the distance from x to D . In particular, f is the identity map at all points of $D \cap M$ and $D \cap M = D \cap N$ [1].*

2. The "orthogonal" disk problem. Some simple closed curves in R^3 have the property that, if J is the curve, there is a disk D whose boundary links J and such that J and D meet in a single point. This property is preserved by homeomorphisms acting on R^3 . The class of curves having this property is known to include curves $\{J\}$ such that under some homeomorphism of R^3 on itself some sub-arc of J is mapped on a rectifiable arc [5]. By the use of his approximation theorem, R. H. Bing [2] has constructed an example of a curve that admits no orthogonal disk.

3. A new formulation of properties $\mathcal{P}$ and $\mathcal{Q}$. A 1-manifold J is said to have property $\mathcal{P}$ if at each point x of J and for positive ϵ there is a set $K(x, \epsilon)$ of diameter less than ϵ that is a topological 2-sphere whose interior contains x , that meets J at a set of points whose cardinal does not exceed the order of x in J , and such that K is locally polyhedral modulo J .

To see that the locally polyhedral character of K may be omitted

Presented to the Society, November 15, 1955; received by the editors March 10, 1956.

¹ This paper was written at the Point Set Topology Institute (University of Wisconsin, 1955) which was supported by a National Science Foundation Grant.

suppose K has the other requirements and $\epsilon' < \epsilon - \text{diameter } K$. Using $M = K$, $D = J$ and ϵ' in the approximation theorem, then $N = K_1(x, \epsilon)$ has the original requirements.

A 1-manifold J is said to have property $\mathcal{Q}$ at x if there is a disk D_1 such that $J \cap D_1$ is the closure of a neighborhood of x in J and D_1 is locally polyhedral at the points of $D_1 \setminus J$. To see that the locally polyhedral character of D_1 may be omitted, suppose D_2 has the requirements of D_1 apart from locally polyhedral character. If $M = D_2$ and $D = J$ and ϵ arbitrary in the approximation theorem we find a disk $N = D_1$ that is locally polyhedral at points of $N \setminus J$ and has the other requirements.

4. 1-cells contained in the interior of 2-cells. It has been known since 1921 when Antoine's thesis appeared that an arc in R^3 may be a subset of no topological 2-cell in R^3 . Such an arc certainly does not lie on the boundary of a 2-cell and still less on the boundary of a 3-cell. (That a 2-cell in R^3 may be a subset of no topological 2-sphere and hence lie on the boundary of no 3-cell is implied by a result announced by Kapuano [7]). We limit ourselves here chiefly to the special case where the 1-cell lies interior (relative) to a 2-cell.

A fairly simple proof is now available that arcs exist lying on no 2-cell. Example 1.4 of Fox-Artin [3] is an example of a wild arc that is the union of two tame arcs, having only an end point in common. If J is the set of points on this arc and J_1 is a sub-arc of J whose interior contains the "wild point" of the imbedded arc, then J_1 is also wild. Now if J were a part of the boundary of some disk D in R^3 , then J_1 has properties $\mathcal{P}$ and $\mathcal{Q}$ (in the new form) and, by Theorem 6 of [6], would be tame. But J_1 is wild.

THEOREM. *Let J be a closed 1-cell in R^3 , then J lies in the interior of a closed 2-cell if and only if J lies on the boundary of some bounded, open, connected ulc^1 subset of R^3 .*

If: Let A be the bounded, open, connected subset of R^3 that is ulc^1 and has J on its boundary. By the results of Wilder, [8, Theorem 8.3, p. 311], every nondegenerate component of the boundary M of A is a classical 2-manifold. If M_1 is the component of M determined by J , then, by triangulation of M_1 , an open 2-cell E in M_1 may be found such that J is in $\text{Int } E$ (rel M_1).

Only if: If J lies in the interior of a disk E , there is a sub-disk E_1 of E such that J is in $|\partial E_1|$. Using the approximation theorem (with $D = J$, $M = E_1$) there is an "almost" polyhedral disk N whose boundary contains J that is locally polyhedral at points of $N \setminus J$. One may "inflate" N to obtain a closed 3-cell F that is locally polyhedral at

points $F \setminus J$ and $|\partial F|$ contains J . Then J is contained in the closure of the interior of F and $\text{Int } F$ is the required bounded, open, connected, ulc^1 subset.

5. A special case of the union of two locally tame sets. It is known that locally tame sets are tame. However, the union of two tame sets having a single common point need not be tame. In this connection the following theorem is of interest.

THEOREM. *Let J be a simple closed curve that is polyhedral and let J be the boundary of any disk F in R^3 . Then J is the boundary of a polyhedral disk.*

(Thus J is tame and unknotted).

PROOF. If J is the boundary of any disk D , then J has the (new) property $\mathcal{Q}$ at each point. Since J is polyhedral, it has property $\mathcal{P}$ at each point. By Theorem VII of [6] it is locally tamely imbedded, hence tamely imbedded. By Graeub [4, p. 39], this implies there is a polyhedral 2-cell of which J is the boundary.

REFERENCES

1. R. H. Bing, *Approximating surfaces with polyhedral ones*, Bull. Amer. Math. Soc. Abstract 61-6-824. (A sketch of the proof may be found in the Proceedings of the Point Set Topology Institute, University of Wisconsin, 1955).
2. ———, *A simple closed curve which pierces no disk*, Bull. Amer. Math. Soc. Abstract 62-1-156.
3. R. Fox and E. Artin, *Some wild cells and spheres in 3-dimensional space*, Ann. of Math. vol. 49 (1948) pp. 979–990.
4. W. Graeub, *Die semi-linearen Abbildungen*, Sitzungsberichte der Heidelberger Akad., 1950, Part 4.
5. O. G. Harrold, Jr., *A theorem on disks*, Proc. Amer. Math. Soc. vol. 7 (1956) pp. 153–154.
6. O. G. Harrold, Jr., H. C. Griffith and E. E. Posey, *A characterization of tame curves in 3-space*, Trans. Amer. Math. Soc. vol. 79 (1955) pp. 12–35.
7. I. Kapuano, *Sur les surfaces homéomorphes à un disque dans un R^3* , C. R. Acad. Sci. Paris vol. 256 (1953) pp. 1229–1230.
8. R. L. Wilder, *Topology of manifolds*, Amer. Math. Soc. Colloquium Publications, vol. 32, 1949.

THE UNIVERSITY OF TENNESSEE