AN INTERPOLATION THEOREM FOR SUBLINEAR OPERATORS ON H_p SPACES

GUIDO WEISS

The purpose of this paper is to extend an interpolation theorem for sublinear operators on L_p spaces, obtained by Calderón and Zygmund, to similar operators acting on H_p spaces, $p > 0$ [1]. We begin by making a few definitions.

Let T be a mapping defined on a class of functions that are analytic in the interior of the unit circle $|w| < 1$, taking values that are measurable functions on some measure space (M, μ) . We say that T is *sublinear* if

- (i) TF is uniquely defined if $F = F_1 + F_2$ and TF_1 and TF_2 are defined;
- (ii) For any constant k , $T(kF)$ is defined if TF is defined;
- (iii) $|T(F_1 + F_2)| \leq |TF_1| + |TF_2|$, $|T(kF)| = |k| \cdot |T(F)|$ almost everywhere.

We shall also say that T is of *type* (a, b) , $a, b > 0$, if the domain of T includes H_a and there exists a constant M , independent of F in H_a , such that $\|TF\|_b \leq M \cdot \|F\|_a$ (the norm on the left is the L_b norm and the norm on the right is the H_a norm).

For the results in the theory of H_p spaces we shall use, we refer the reader to [2].

THEOREM. *Let T be a sublinear operator of types $(1/\alpha_1, 1/\beta_1)$ and $(1/\alpha_2, 1/\beta_2)$ with constants M_1 and M_2 respectively, where $1/\alpha_i > 0$, $1/\beta_i \geq 1$ ($i = 1, 2$). Let $\alpha = (1 - t_0)\alpha_1 + t_0\alpha_2$ and $\beta = (1 - t_0)\beta_1 + t_0\beta_2$, where $0 \leq t_0 \leq 1$. Then there exists an absolute constant, K , such that*

$$\|TF\|_{1/\beta} \leq KM_1^{(1-t_0)} M_2^{t_0} \|F\|_{1/\alpha}$$

for all F in $H_{1/\alpha}$ and, thus, T is of type $(1/\alpha, 1/\beta)$.

We remark that the above theorem in the special case where T is linear is known [3].

Let us assume that $\alpha_1 \leq \alpha \leq \alpha_2$. Let n be an integer such that $n/\alpha_1, n/\alpha_2 > 1$. In the proof we shall suppose that $n = 2$, the proof being easily extensible to the general case.

Let $f = \sum_k c_k \epsilon_k \chi_k$, where $c_k > 0$, $|\epsilon_k| = 1$ and the χ_k 's are a finite number of characteristic functions of mutually disjoint measurable subsets, E_k , of $[0, 2\pi]$. Let

Received by the editors March 24, 1956.

$$(1) \quad F(w) = 1/2\pi \int_0^{2\pi} \frac{e^{it} + w}{e^{it} - w} f(t) dt.$$

It is well known that F is in H_p for all $p > 0$ and, thus, F^2 is in H_p for all $p > 0$. Hence, $T(F^2)$ is well defined. We are going to make use of the fact that

$$(2) \quad \|T(F^2)\|_{1/\beta} = \sup \left\{ \int_M |T(F^2)| g d\mu; g \text{ simple, } g \geq 0, \|g\|_{1/1-\beta} = 1 \right\}.$$

We make the further assumption that $\|f\|_{2/\alpha} = 1$.

Set $g = \sum_i c_i' \chi_i'$ where $c_i' > 0$ and χ_i' is the characteristic function of the measurable set E_i' of M , and suppose that $\|g\|_{1/1-\beta} = 1$. We shall now estimate

$$(3) \quad \int_M |T(F^2)| g d\mu.$$

This estimate will then give us a bound for $\|T(F^2)\|_{1/\beta}$.

Let $z = x + iy$ be any complex number and define

$$F_z(w) = 1/2\pi \int_0^{2\pi} \frac{e^{it} + w}{e^{it} - w} |f(t)|^{\alpha(z)/\alpha} e^{i \arg f(t)} dt$$

where $\alpha(z) = (1-z)\alpha_1 + z\alpha_2$ and $\beta(z) = (1-z)\beta_1 + z\beta_2$. We now consider

$$(4) \quad \Phi(z) = \int_M |T(F_z^2)| g^{1-\beta(z)/1-\beta} d\mu.$$

Clearly, when $z = t_0$, $\Phi(z)$ reduces to (3).

Our plan is to show that $\log \Phi(z)$ is a continuous subharmonic function that is bounded in every vertical strip in the plane. An application of the three line theorem for such functions will then give us

$$(5) \quad \log \Phi(t_0 + iy) \leq (1 - t_0) \log \max_y \Phi(iy) + t_0 \log \max_y \Phi(1 + iy).$$

But, making use of Hölder's inequality and M. Riesz's inequality [2], which implies that if h is in $L_p(0, 2\pi)$, $p > 1$, and

$$H(w) = 1/2\pi \int_0^{2\pi} \frac{e^{it} + w}{e^{it} - w} h(t) dt$$

then there exists a constant, A_p , independent of h , such that $\|H\|_p \leq A_p \cdot \|h\|_p$, we have, on the line $x = 0$

$$\begin{aligned}\Phi(iy) &\leq \|TF_{iy}^2\|_{1/\beta_1} \cdot \|g^{1-\beta(0)/1-\beta}\|_{1/1-\beta_1} \leq M_1 \|F_{iy}^2\|_{1/\alpha_1} \cdot 1 = M_1 \|F_{iy}\|_{2/\alpha_1}^2 \\ &\leq M_1 \cdot A_{2/\alpha_1}^2 \| |f(t)|^{\alpha(iy)/\alpha} \|_{2/\alpha_1}^2 = M_1 A_{2/\alpha_1}^2\end{aligned}$$

since $\|f\|_{2/\alpha}=1$ and $\|g\|_{1/1-\beta}=1$ and $\left\{ \|f\|^{\alpha(iy)/\alpha} \right\}_{2/\alpha_1} = \left\{ |f|^\alpha \right\}_{2/\alpha_1} = \|f\|_{2/\alpha}^{2/\alpha}$ and, similarly, $\left\{ \|(g)^{1-\beta(0)/1-\beta}\| \right\}_{1/1-\beta_1} = \|g\|_{1/1-\beta}^{1-\beta}$.

Likewise, on the line $x=1$ we have $\Phi(1+iy) \leq M_2 A_{2/\alpha_2}^2$.

Thus, letting $y=0$ in (5), since g was an arbitrary simple function satisfying $\|g\|_{1/1-\beta}=1$, we obtain, upon comparing with (2)

$$(6) \quad \|T(F^2)\|_{1/\beta} \leq KM_1^{(1-t_0)} M_2^{t_0}$$

where $K = \{A_{2/\alpha_1}^{(1-t_0)} A_{2/\alpha_2}^{t_0}\}^2$.

The last inequality is the conclusion of the theorem for the special case of functions in $H_{1/\alpha}$ that are squares of functions of the type (1) with $\|f\|_{2/\alpha}=1$. We will show later on, however, that we can deduce the theorem from (6).

The fact that $\log \Phi(z)$ is bounded in every vertical strip in the plane is immediate. For expanding (4) and using (iii) we obtain

$$\Phi(z) \leq \sum_i c_i'^{1-\beta(z)/1-\beta} \int_{E_i} \sum_{n,m} (c_n c_m)^{\alpha(z)/\alpha} |T(\Delta_n \cdot \Delta_m)| d\mu,$$

where

$$\Delta_s(w) = 1/2\pi \int_0^{2\pi} \frac{e^{it} + w}{e^{it} - w} \chi_s(t) dt.$$

We now proceed to show that $\log \Phi(z)$ is continuous. We have

$$\Phi(z) = \sum_i \int_{E_i'} c_i'^{1-\beta(z)/1-\beta} |T(F_s^2)| d\mu = \sum_i \int_{E_i'} |T c_i'^{1-\beta(z)/1-\beta} (F_s^2)| d\mu.$$

We shall investigate each of these summands. Let

$$\psi_{i,s} = \psi_s = c_i'^{1-\beta(z)/1-\beta} (F_s^2)$$

and $\Psi_i(z) = \Psi(z) = \int_{E_i'} |T\psi_s| d\mu$.

It follows immediately from (i), (ii), and (iii) that if TF_1 and TF_2 are defined so is $T(F_1 - F_2)$ and we have

$$(7) \quad \left| |TF_1| - |TF_2| \right| \leq |T(F_1 - F_2)|.$$

It clearly suffices to show that $\Psi(z)$ is continuous. Using (7) and Hölder's inequality, we obtain

$$\begin{aligned}
|\Psi(z) - \Psi(z + \Delta z)| &= \left| \int_{E_l} |T\psi_{z+\Delta z}| d\mu - \int_{E_l} |T\psi_z| d\mu \right| \\
&\leq \int_{E_l} |T(\psi_{z+\Delta z} - \psi_z)| d\mu \\
&\leq \|T(\psi_{z+\Delta z} - \psi_z)\|_{1/\beta_1} \cdot \{\mu(E_l')\}^{1/1-\beta_1} \\
&\leq M_1 \cdot \|\psi_{z+\Delta z} - \psi_z\|_{1/\alpha_1} \cdot \{\mu(E_l')\}^{1/1-\beta_1}.
\end{aligned}$$

Thus, if we can show that $\|\psi_{z+\Delta z} - \psi_z\|_{1/\alpha_1} \rightarrow 0$ as $\Delta z \rightarrow 0$ we have the desired continuity. We proceed to do this:

$$|\psi_{z+\Delta z} - \psi_z| = |\{c_l'^{1-\beta(z+\Delta z)/1-\beta}\} F_{z+\Delta z}^2 - \{c_l'^{1-\beta(z)/1-\beta}\} F_z^2| \leq A + B,$$

where

$$A = |\{c_l'^{1-\beta(z+\Delta z)/1-\beta}\} (F_{z+\Delta z}^2 - F_z^2)|$$

and

$$B = |(c_l'^{1-\beta(z+\Delta z)/1-\beta} - c_l'^{1-\beta(z)/1-\beta}) F_z^2|.$$

If $1/\alpha_1 \leq 1$ we thus have

$$(8) \quad \|\psi_{z+\Delta z} - \psi_z\|_{1/\alpha_1}^{1/\alpha_1} \leq \|A\|_{1/\alpha_1}^{1/\alpha_1} + \|B\|_{1/\alpha_1}^{1/\alpha_1}.$$

(If $1/\alpha_1 > 1$ we do not take the $1/\alpha_1$ powers of the norms and use Minkowski's inequality instead of (8) and the remaining part of the proof is essentially the same.)

We have, using Schwarz's inequality:

$$\begin{aligned}
\|A\|_{1/\alpha_1} &= |c_l'^{1-\beta(z+\Delta z)/1-\beta}| \cdot \|F_{z+\Delta z}^2 - F_z^2\|_{1/\alpha_1} \\
&\leq |c_l'^{1-\beta(z+\Delta z)/1-\beta}| \cdot \|F_{z+\Delta z} - F_z\|_{2/\alpha_1} \cdot \|F_{z+\Delta z} + F_z\|_{2/\alpha_1}.
\end{aligned}$$

Using M. Riesz's inequality we have that the last terms are dominated by

$$\begin{aligned}
A_{2/\alpha_1}^2 |c_l'^{1-\beta(z+\Delta z)/1-\beta}| \cdot \| |f(t)|^{\alpha(z+\Delta z)/\alpha} - |f(t)|^{\alpha(z)/\alpha} \|_{2/\alpha_1} \\
\cdot \| |f(t)|^{\alpha(z+\Delta z)/\alpha} + |f(t)|^{\alpha(z)/\alpha} \|_{2/\alpha_1}.
\end{aligned}$$

But $|f(t)|^{\alpha(z+\Delta z)/\alpha} - |f(t)|^{\alpha(z)/\alpha}$ is 0 outside $\cup E_k$ and tends to 0 uniformly in t there as $\Delta z \rightarrow 0$. Therefore,

$$\| |f(t)|^{\alpha(z+\Delta z)/\alpha} - |f(t)|^{\alpha(z)/\alpha} \|_{2/\alpha_1} \rightarrow 0 \text{ as } \Delta z \rightarrow 0.$$

On the other hand, both $|c_l^{1-\beta(z+\Delta z)/1-\beta}|$ and $\| |f(t)|^{\alpha(z+\Delta z)/\alpha} + |f(t)|^{\alpha(z)/\alpha} \|_{2/\alpha_1}$ are bounded for Δz small. This shows that $\|A\|_{1/\alpha_1} \rightarrow 0$ as $\Delta z \rightarrow 0$. We also have

$$\|B\|_{1/\alpha_1} = |c_l^{1-\beta(z+\Delta z)/1-\beta} - c_l^{1-\beta(z)/1-\beta}| \cdot \|F_z^2\|_{1/\alpha_1} \rightarrow 0$$

as $\Delta z \rightarrow 0$.

Hence, we have shown that $\|\psi_{z+\Delta z} - \psi_z\|_{1/\alpha_1} \rightarrow 0$ as $\Delta z \rightarrow 0$ and, thus, $\Psi(z)$ is continuous.

We shall now show that $\log \Phi(z)$ is subharmonic. It is sufficient to show that if $h(z)$ is any harmonic function then $\Phi(z)e^{h(z)}$ is subharmonic. To achieve this end it is sufficient to show that $\Psi_l(z)e^{h(z)} = \Psi(z)e^{h(z)}$ is subharmonic, for then

$$\Phi(z)e^{h(z)} = \sum_l \Psi_l(z)e^{h(z)}$$

being a sum of subharmonic functions is subharmonic. Thus we fix $h(z)$, an arbitrary harmonic function, and denote by $H(z)$ an analytic function whose real part is $h(z)$. Since the problem is local, we may consider h and H in a given circle. We write

$$\psi_z^* = \psi_z e^{H(z)} \text{ and } \Psi^*(z) = \Psi(z)e^{h(z)} = \int_{E_1} |T\psi_z^*| d\mu.$$

Fix z and let $\rho > 0$, and denote by $z_1, z_2, \dots, z_p$ a collection of points equally spaced over the circumference of the circle with center z and radius ρ . We have

$$(9) \quad \psi_z^*(w) = \lim_{q \rightarrow \infty} 1/q \sum_{j=1}^q \psi_{z_j}^*(w)$$

since $\psi_{z_j}^*(w)$ is analytic for each w in the unit circle. This is easily seen since

$$\psi_z(w) = c_l^{1-\beta(z)/1-\beta} \left\{ \sum_k c_k^{\alpha(z)/\alpha} \epsilon_k \Delta_k(w) \right\}^2$$

is clearly analytic for each w . It will be necessary for our purpose to analyze further the convergence in (9). More precisely, we will show

$$(10) \quad \|\psi_z^*(w) - 1/q \sum_{j=1}^q \psi_{z_j}^*(w)\|_{1/\alpha_1} \rightarrow 0 \text{ as } q \rightarrow \infty.$$

Suppose that $1/\alpha_1 \leq 1$ (if $1/\alpha_1 > 1$ we do not take the $1/\alpha_1$ powers and, as before, use Minkowski's inequality, the rest of the proof being the same). We have

$$\begin{aligned}
& \left\| \psi_z^*(w) - 1/q \sum_{j=1}^q \psi_{z_j}^*(w) \right\|_{1/\alpha_1}^{1/\alpha_1} \\
&= \left\| e^{H(z)} c_l^{1-\beta(z)/1-\beta} F_z^2 - 1/q \sum_{j=1}^q e^{H(z_j)} c_l^{1-\beta(z_j)/1-\beta} F_{z_j}^2 \right\|_{1/\alpha_1}^{1/\alpha_1} \\
&= \left\| e^{H(z)} c_l^{1-\beta(z)/1-\beta} \sum_{n,m} (c_n c_m)^{\alpha(z)/\alpha} \epsilon_n \epsilon_m \Delta_n(w) \Delta_m(w) \right. \\
&\quad \left. - 1/q \sum_{j=1}^q e^{H(z_j)} c_l^{1-\beta(z_j)/1-\beta} \sum_{n,m} (c_n c_m)^{\alpha(z_j)/\alpha} \epsilon_n \epsilon_m \Delta_n(w) \Delta_m(w) \right\|_{1/\alpha_1}^{1/\alpha_1} \\
&\leq \sum_{n,m} \left\{ \left| e^{H(z)} c_l^{1-\beta(z)/1-\beta} (c_n c_m)^{\alpha(z)/\alpha} \epsilon_n \epsilon_m \right. \right. \\
&\quad \left. \left. - 1/q \sum_{j=1}^q e^{H(z_j)} c_l^{1-\beta(z_j)/1-\beta} (c_n c_m)^{\alpha(z_j)/\alpha} \epsilon_n \epsilon_m \right| \cdot \left\| \Delta_n(w) \Delta_m(w) \right\|_{1/\alpha_1} \right\}^{1/\alpha_1}.
\end{aligned}$$

But each of the terms

$$e^{H(z)} c_l^{1-\beta(z)/1-\beta} (c_n c_m)^{\alpha(z)/\alpha} - 1/q \sum_{j=1}^q e^{H(z_j)} c_l^{1-\beta(z_j)/1-\beta} (c_n c_m)^{\alpha(z_j)/\alpha}$$

converge to 0 as $q \rightarrow \infty$. This proves (10).

We thus have

$$\begin{aligned}
\int_{E_l} \left| T \left(\psi_z^* - 1/q \sum_1^q \psi_{z_j}^* \right) \right| d\mu &\leq \left\| T \left(\psi_z^* - 1/q \sum_1^q \psi_{z_j}^* \right) \right\|_{1/\beta_1} \{ \mu(E_l') \}^{1/1-\beta_1} \\
&\leq M_1 \left\| \psi_z^* - 1/q \sum_1^q \psi_{z_j}^* \right\|_{1/\alpha_1} \cdot \{ \mu(E_l') \}^{1/1-\beta_1} \rightarrow 0 \text{ as } q \rightarrow \infty.
\end{aligned}$$

A fortiori, we have

$$e_q = \int_{E_l} \left| |T\psi_z^*| - \left| T \left(1/q \sum_1^q \psi_{z_j}^* \right) \right| \right| d\mu \rightarrow 0, \text{ as } q \rightarrow \infty.$$

Thus, since

$$\int_{E_l} |T\psi_q^*| d\mu \leq e_q + 1/q \sum_1^q \int_{E_l} |T\psi_{z_j}^*| d\mu$$

we obtain

$$\Psi^*(z) \leq \lim_{q \rightarrow \infty} 1/q \sum_1^q \Psi^*(z_j) = 1/2\pi \int_0^{2\pi} \Psi^*(z + \rho e^{it}) dt.$$

We have thus proved that $\log \Phi(z)$ is subharmonic. Thus our next task is to show that (6) implies the theorem.

Suppose that F is defined as in equation (1) where f is a simple function, but we no longer assume that $\|f\|_{2/\alpha}=1$. We then have $\|f\|_{2/\alpha} \leq \|F\|_{2/\alpha}$. Thus

$$\begin{aligned} \|T(F^2/\|F\|_{2/\alpha}^2)\|_{1/\beta} &= 1/\|F\|_{2/\alpha}^2 \cdot \|T(F^2)\|_{1/\beta} \leq 1/\|f\|_{2/\alpha}^2 \cdot \|T(F^2)\|_{1/\beta} \\ &= \|T(F^2/\|f\|_{2/\alpha}^2)\|_{1/\beta} \leq KM_1^{(1-t_0)} M_2^{t_0} \end{aligned}$$

since $F/\|f\|_{2/\alpha}$ is a function of type (1) for which this inequality was proved. But

$$\|F\|_{2/\alpha}^2 = \|F^2\|_{1/\alpha}.$$

We thus have, using the last part of (iii)

$$(11) \quad \|T(F^2)\|_{1/\beta} \leq KM_1^{(1-t_0)} M_2^{t_0} \|F^2\|_{1/\alpha}$$

for any function of type (1) where f is simple.

Now let G be a function in $H_{1/\alpha}$ that is the square of a function F in $H_{2/\alpha}$. Let f be a function in $L^{2/\alpha}(0, 2\pi)$ such that $F(w) = 1/2\pi \int_0^{2\pi} ((e^{it} + w/e^{it} - w))f(t)dt$ (i.e. f is the real part of $F(e^{it})$). Let $\{f_n\}$ be a sequence of simple functions such that $\|f_n - f\|_{2/\alpha} \rightarrow 0$ as $n \rightarrow \infty$ and set $F_n(w) = 1/2\pi \int_0^{2\pi} ((e^{it} + w)/(e^{it} - w))f_n(t)dt$. Then $F^2 - F_n^2$ is in $H_{1/\alpha} \subset H_{1/\alpha_2}$ (since $1/\alpha \geq 1/\alpha_2$). Thus, in particular, F^2 is a sum of a function F_n in H_{1/α_2} and a function $F^2 - F_n^2$ in H_{1/α_2} , showing, by (i), that $T(F^2)$ is defined. We also have

$$(12) \quad |TF^2| - |TF_n^2| \leq |T(F^2 - F_n^2)|.$$

But $\|T(F^2 - F_n^2)\|_{1/\beta_2} \leq M_2 \|F^2 - F_n^2\|_{1/\alpha_2} \leq M_2 \|F - F_n\|_{2/\alpha_2} \cdot \|F + F_n\|_{2/\alpha_2} \leq M_2 A_{2/\alpha_2}^2 \|f - f_n\|_{2/\alpha_2} \cdot \|f + f_n\|_{2/\alpha_2} \leq CM_2 A_{2/\alpha_2}^2 \|f - f_n\|_{2/\alpha} \cdot \|f + f_n\|_{2/\alpha}$ where C does not depend on the functions f and f_n but is such that $\|g\|_{2/\alpha_2} \leq C^{1/2} \|g\|_{2/\alpha}$ for any g in $L_{2/\alpha}(0, 2\pi)$ (C exists because the measure space is finite). The last factor is bounded since $\|f_n - f\|_{2/\alpha} \rightarrow 0$ and, by this very last fact, the entire right-hand side of the inequality goes to 0 as $n \rightarrow \infty$. Thus there exists a subsequence $|T(F^2 - F_{n_k}^2)| \rightarrow 0$ almost everywhere and, consequently, $|T(F_{n_k}^2)| \rightarrow |T(F^2)|$ almost everywhere, by (12). By Fatou's lemma, therefore, we have

$$\begin{aligned} \int |T(F^2)|^{1/\beta} d\mu &\leq \liminf \int |T(F_{n_k}^2)|^{1/\beta} d\mu \\ &\leq \liminf \{KM_1^{(1-t_0)} M_2^{t_0} \|F_{n_k}^2\|_{1/\alpha}\}^{1/\beta} \\ &= \{KM_1^{(1-t_0)} M_2^{t_0} \|F^2\|_{1/\alpha}\}^{1/\beta}. \end{aligned}$$

We have shown

$$(13) \quad \|T(F^2)\|_{1/\beta} \leq KM_1^{(1-t_0)} M_2^{t_0} \|F^2\|_{1/\alpha}$$

for any function F in $H_{2/\alpha}$.

But any function G in $H_{1/\alpha}$ is a sum of two functions, G_1 and G_2 , with no zeros, both in $H_{1/\alpha}$, such that $\|G_1\|_{1/\alpha} = \|G\|_{1/\alpha}$ and $\|G_2\|_{1/\alpha} \leq 2\|G\|_{1/\alpha}$. This is easily seen by the Blaschke product decomposition, $G = BG_1$ where B is the Blaschke product which satisfies $|B(z)| < 1$ if $|z| < 1$, and G_1 has no zeros. Thus, $G = (B-1)G_1 + G_1$ is the desired decomposition with $G_2 = (B-1)G_1$. The norm inequalities follow from the fact that $\|G_1\|_{1/\alpha} = \|G\|_{1/\alpha}$ and $|B-1| \leq 2$ in the unit circle. Since G_1 and G_2 have no zeros they can be written as the squares of their square roots, and, clearly, these square roots are in $H_{2/\alpha}$. Thus, using (iii)

$$\begin{aligned} \|TG\|_{1/\beta} &= \|T(G_1 + G_2)\|_{1/\beta} \leq \| |TG_1| + |TG_2| \|_{1/\beta} \leq \|TG_1\|_{1/\beta} + \|TG_2\|_{1/\beta} \\ &\leq KM_1^{(1-t_0)} M_2^{t_0} \{ \|G_1\|_{1/\alpha} + \|G_2\|_{1/\alpha} \} \\ &\leq KM_1^{(1-t_0)} M_2^{t_0} \{ \|G\|_{1/\alpha} + 2\|G\|_{1/\alpha} \} = 3KM_1^{(1-t_0)} M_2^{t_0} \|G\|_{1/\alpha}. \end{aligned}$$

This proves the theorem.

REFERENCES

1. A. P. Calderon and A. Zygmund, *A note on the interpolation of sublinear operations*, Amer. J. Math. vol. 78 (1956).
2. A. Zygmund, *Trigonometrical series*, Warsaw, 1935. (New York, Stechert).
3. A. P. Calderon and A. Zygmund, *On the theorem of Hausdorff-Young, contributions to Fourier analysis*, Annals of Mathematics Studies, no. 25, Princeton University Press, 1950, pp. 166-168.

DE PAUL UNIVERSITY