ON THE SOLUTION OF $f(f(z)) = e^z - 1$ AND ITS DOMAIN OF REGULARITY

ROBERT OSSERMAN

The problem of examining the complex solutions $f(z)$ of the equation

$$(1) \quad f(f(z)) = e^z - 1$$

was suggested to the author by S. Chowla who, together with Kempner, Rivlin, and Thron, proved that $f(z)$ cannot be an entire function. This result is contained as a special case in two recent papers [1; 3] using the theory of entire functions. The purpose of the present note is to show how the use of elementary geometric methods leads quite easily to a slightly stronger theorem which gives some information on a maximum domain of regularity for $f(z)$. This method may also be applied to the equation

$$(2) \quad f(f(z)) = e^z$$

which was treated in detail by Kneser [2] who proved the existence of a solution analytic on the whole real axis.

THEOREM. *Let $z = x + iy$ and let Ω denote an infinite strip $|y| < b$ for some constant $b > \pi$. Let $f(z)$ be a function defined in some domain D such that $\Omega \subset D$ and $f(\Omega) \subset D$. If $f(z)$ satisfies (1) throughout D then it cannot be analytic in Ω .*

PROOF. Denote by R the image region under $w = f(z)$ of the strip $S: |y| < \pi$. Then if $\zeta = f(w) = f(f(z)) = e^z - 1$, the image under $f(w)$ of R must be the region T consisting of the ζ -plane slit along the negative real axis from -1 to $-\infty$. Since the composed map $\zeta = f(f(z))$ is a one-one map of S onto T , the maps f of S onto R and of R onto T must also be one-one. Furthermore, since the given correspondences between S and R and between S and T are conformal, so is the correspondence between R and T .

We note next that $f(0) = 0$. Namely, if $f(0) = a$, then $f(a) = f(f(0)) = 0$, so that $a = f(0) = f(f(a)) = e^a - 1$. But evaluating the derivative of equation (1) at the points 0 and a respectively, we find $f'(a)f'(0) = 1$ and $f'(0)f'(a) = e^a = a + 1$, whence $a = 0$.

Presented to the Society, August 24, 1956 under the title *On the solution of a functional equation*; received by the editors March 15, 1956 and, in revised form, June 27, 1956.

We may thus write the power expansion for $f(z)$ at the origin in the form $f(z) = a_1 z + a_2 z^2 + \dots$, and inserting this in equation (1) shows that $a_1 = 1$ and all the a_n are real. In other words, $f'(0) = 1$ and $f(z)$ is real on the real axis. But the image of the negative x -axis cannot be the whole negative axis in the w -plane, since the composed map $f(f(z))$ must take the negative x -axis onto the interval $(-1, 0)$. Hence $w = f(z)$ must map the negative x -axis onto an interval $(-c, 0)$ such that $f: (-c, 0) \rightarrow (-1, 0)$. This implies that $c > 1$.

We can now show explicitly that the function $f(z)$ must have a singularity at the point $z_0 = \log(c-1) + i\pi$ of the region Ω . To see this, denote by R' a copy of the region R placed in the ζ -plane. Then we can find a sequence of points ζ_n in the upper half ζ -plane such that $\zeta_n \in R' \cap T$ for all n , and $\zeta_n \rightarrow -c$. Then the corresponding points z_n of S such that $\zeta_n = f(f(z_n))$ will approach z_0 . On the other hand, their images $w_n = f(z_n)$ must approach $-\infty$. To see this we need only note that $w_n = f^{-1}(\zeta_n)$, so that if we consider the points $z'_n = w_n$ of the z -plane and the points $w'_n = \zeta_n$ of the w -plane, then $z'_n = f^{-1}(w'_n)$, and $w'_n \rightarrow -c$ implies $z'_n \rightarrow -\infty$.

BIBLIOGRAPHY

1. Baker, *The iteration of entire transcendental functions and the solution of the functional equation $f(f(z)) = F(z)$* , Math. Ann. vol. 129 (1955) pp. 174-180.
2. Kneser, *Reelle analytische Lösungen der Gleichung $\phi(\phi(x)) = e^x$* , J. Reine Angew. Math. vol. 187 (1950) pp. 56-67.
3. Thron, *Entire solutions of the functional equation $f(f(z)) = g(z)$* , Canadian Journal of Mathematics vol. 8 (1956) pp. 47-48.

STANFORD UNIVERSITY