A THEOREM OF DICKSON ON NONVANISHING CUBIC FORMS IN A FINITE FIELD

L. CARLITZ

Dickson stated that a ternary cubic form with coefficients in $GF(q)$, $q = p^n$, $p > 2$, that vanishes only for $x = y = z = 0$, is the norm of a linear form with coefficients in $GF(q^3)$. The proof [2, p. 161, particularly line 15] does not seem adequate. In this note we sketch another proof.

Let $f = f(x, y, z)$ be a homogeneous cubic polynomial with coefficients in $GF(q)$ that vanishes in $GF(q)$ only for $x = y = z = 0$. Put

$$(1) \quad f = f_0 z^3 + f_1 z^2 + f_2 z + f_3,$$

where $f_j = f_j(x, y)$ is homogeneous of degree j . Assume first that $f_0 \neq 0$, so that we may take $f_0 = 1$; also we assume to begin with that $p \neq 3$. Then by means of a linear transformation we may suppose that $f_1 = 0$. Hence consider

$$(2) \quad f = z^3 + f_2 z + f_3.$$

Now for fixed $a, b \in GF(q)$, a, b not both 0, it is clear that $f(a, b, z)$ is irreducible. Consequently its discriminant is a nonzero square of $GF(q)$; that is

$$(3) \quad D = D(a, b) = -4f_2^3(a, b) - 27f_3^2(a, b)$$

is a square for all $a, b \in GF(q)$, $D = 0$ only for $a = b = 0$. But Dickson [1, Theorem 3] has proved that for $q \geq 13$, this implies

$$D(x, y) = C^2(x, y),$$

where $C(x, y)$ is a homogeneous cubic polynomial. Thus (3) becomes

$$(4) \quad C^2(x, y) = -4f_2^3(x, y) - 27f_3^2(x, y).$$

Clearly it is necessary that $f_3(x, y)$ be irreducible, for otherwise there exist two number a, b of the field, not both 0, such that $f_3(a, b) = 0$, and therefore $f(a, b, 0) = 0$. Thus either $f_2 = 0$ or $(f_2, f_3) = (C, f_3) = 1$. Leaving the case $f_2 = 0$ for the moment, we rewrite (4) as

$$C^2 + 27f_3^2 = -4f_2^3$$

and put $\alpha^2 = -27$, where $\alpha \in GF(q^2)$. Then clearly

Received by the editors June 5, 1956 and, in revised form, February 18, 1957.

$$(5) \quad C + \alpha f_3 = 2\beta u^3, \quad C - \alpha f_3 = -2\beta^{-1}v^3,$$

where $\beta \in GF(q^2)$ and u, v are linear forms with coefficients in $GF(q^2)$. Since (5) implies $C = \beta u^3 - \beta^{-1}v^3$, $\alpha f_3 = \beta u^3 + \beta^{-1}v^3$, $f_2 = uv$, (2) becomes

$$f = z^3 + uvz + \alpha^{-1}\beta u^3 + \alpha^{-1}\beta^{-1}v^3$$

or

$$(6) \quad f = z^3 - 3UVz + U^3 + V^3,$$

where U, V are linear forms in some $GF(q^k)$. But (6) evidently implies $f(-U-V)=0$, from which it follows that the linear form $U+V$ has coefficients in $GF(q^3)$. Indeed if $-U-V=\lambda x+\mu y$, then

$$f(x, y, \lambda x + \mu y) = 0$$

implies $f(x, 0, \lambda x)=0$, so that $\lambda \in GF(q^3)$; similarly for μ . Therefore f is the norm of a linear form in $GF(q^3)$.

In the excluded case $f_2=0$, we have

$$f = z^3 + f_3(x, y).$$

Since f_3 is irreducible, it is only necessary to consider the case

$$(7) \quad f = z^3 + ax^3 + by^3 \quad (ab \neq 0),$$

for the occurrence of a term x^2y or xy^2 can be treated as in the previous proof. Next we need only consider the case $q \equiv 1 \pmod{3}$, $ab=1$, a not a cube of $GF(q)$. But by a theorem of Hurwitz [3] which may be extended without difficulty to finite fields, the equation

$$ax^3 + by^3 + z^3 = 0,$$

where a, b satisfy the stated conditions, has at least one solution with $xyz \neq 0$.

Returning to (1), suppose that $f_0=0$. If a term in x^3 or y^3 occurs, the previous proof applies. We need then consider only the case $f_3 = ax^2y + bxy^2$, and this obviously leads to a contradiction.

When $p=3$ the above discussion must be modified somewhat. Returning to (1), we may assume $f_0=1$ but cannot require that $f_1=0$. If indeed $f_1=0$, then (4) reduces to

$$C^2(x, y) = -4f_2^3(x, y),$$

so that C is the cube of a linear form; this contradicts the requirement that C vanishes only for $x=y=0$.

Employing the discriminant of (1) we get

$$(8) \quad -f_1^3 f_3 - f_2^3 + f_1^2 f_2^2 = C^2.$$

If $f_2=0$, (8) becomes $-f_1^3 f_3 = C^2$, which contradicts the irreducibility of f_3 . We need then consider only the case $f_1 f_2 \neq 0$. By means of a linear transformation we may suppose that $f_1 = x$, $f_2 = ay^2$; then (8) reduces to

$$-a^3 y^6 + a^2 x^2 y^4 - x^3 f_3 = C^2.$$

We may take $a = -1$, $C = y^3 - x^2 y + cx^3$, which yields

$$f_3 = cy^3 - xy^2 + cx^2 y + c^2 x^3.$$

In (1) take $f_1 = x$, $f_2 = -y^2$, $z = \alpha x + \beta y$. Substituting in $f(x, y, z) = 0$, we obtain the following conditions on α, β :

$$\alpha^3 + \alpha^2 = c^2, \quad \beta^3 - \beta = c, \quad \alpha\beta = c, \quad \beta^2 - \alpha = 1.$$

It is easily verified that this system is equivalent to

$$(9) \quad \beta^3 - \beta = c, \quad \alpha\beta = c.$$

Clearly (9) implies $\alpha, \beta \in GF(q^3)$. Hence $f(x, y, z)$ has the linear factor $z - \alpha x - \beta y$, and it follows at once that

$$f(x, y, z) = (z - \alpha x - \beta y)(z - \alpha^q x - \beta^q y)(z - \alpha^{q^2} x - \beta^{q^2} y).$$

In other words, $f(x, y, z)$ is the norm of a linear form with coefficients in $GF(q^3)$; moreover $1, \alpha, \beta$ are linearly independent with respect to $GF(q)$.

We may state the following

THEOREM. *Let $f(x, y, z)$ be a homogeneous cubic polynomial with coefficients in $GF(q)$ that vanishes in $GF(q)$ only for $x=y=z=0$. Then if $p > 2$, $q \geq 13$, $f(x, y, z)$ is the norm of a linear form $\alpha x + \beta y + \gamma z$, where α, β, γ are in $GF(q^3)$ and are linearly independent with respect to $GF(q)$.*

The converse of the theorem is evidently true. Possibly the theorem holds for $p=2$; the above proof obviously does not apply in that case.

REFERENCES

1. L. E. Dickson, *Definite forms in a finite field*, Trans. Amer. Math. Soc. vol. 10 (1909) pp. 109-122.
2. ———, *On triple algebras and ternary cubic forms*, Bull. Amer. Math. Soc. vol. 14 (1907-1908) pp. 160-169.
3. A. Hurwitz, *Über die Kongruenz $ax^e + by^e + cz^e \equiv 0 \pmod{p}$* , J. Reine Angew. Math. vol. 136 (1909) pp. 272-292 (= Mathematische Werke, vol. 2, Basel, 1933, pp. 430-445).