ON COHOMOLOGY OF LIE ALGEBRAS

GEORGE F. LEGER, JR.

Let G be a Lie algebra, L an ideal of G and M a G -module. The n -dimensional cochains for L in M are n -linear alternating functions on L^n with values in M . These cochains form a vector space $C^n(L, M)$ over the ground field of L ; we identify $C^0(L, M)$ with M . $C^0(L, M)$ is already a G -module and for $n > 0$, f in $C^n(L, M)$ and γ in G put

$$\begin{aligned} (\gamma \cdot f)(\sigma_1, \dots, \sigma_n) \\ = \gamma \cdot f(\sigma_1, \dots, \sigma_n) - \sum_{i=1}^n f(\sigma_1, \dots, \sigma_{i-1}, [\gamma, \sigma_i], \sigma_{i+1}, \dots, \sigma_n) \end{aligned}$$

for all $\sigma_1, \dots, \sigma_n$ in L .

This definition makes $C^n(L, M)$ into a G -module and further $\gamma \cdot (\delta f) = \delta(\gamma \cdot f)$ where δ is the coboundary operator: $C^n(L, M) \rightarrow C^{n+1}(L, M)$ so that the cohomology groups $H^n(L, M)$ take on the structure of a G -module. Professor Hochschild suggested to the author that he define an operation of G on the standard interpretations of the low dimensional cohomology groups which would agree with the operation of G on $H^n(L, M)$ as given above.

This problem has an analogue in the cohomology of groups. There, if L is a normal subgroup of a group G and f is any function of L^{n+1} to M , the operation

$$(\gamma \cdot f)(\sigma_0, \dots, \sigma_n) = \gamma \cdot f(\gamma^{-1}\sigma_0\gamma, \dots, \gamma^{-1}\sigma_n\gamma)$$

for γ in G and $\sigma_0, \dots, \sigma_n$ in L , again induces the structure of a G -module on the cohomology groups $H^n(L, M)$. In an appendix we outline briefly a way of making the standard interpretations of the low dimensional cohomology groups into G -modules in a way consistent with the operation of G on the cohomology groups.

The author is pleased to acknowledge his indebtedness to Professor Hochschild for his suggestions and encouragement.

1. Let G be a Lie algebra over a ground field of *characteristic* 0, L an ideal of G such that G/L is semi-simple and let M be a G -module. We denote by ρ_n the restriction homomorphism of $H^n(G, M)$ into $H^n(L, M)$. For any G -module U , let $U^G = \{u \mid \gamma \cdot u = 0 \text{ for all } \gamma \text{ in } G\}$. A special case of a theorem of Hochschild-Serre [1, p. 603, Theorem 13]¹ is

Presented to the Society, April 21, 1956 and December 29, 1956; received by the editors May 15, 1956 and, in revised form, March 2, 1957.

¹ Numbers in square brackets refer to the bibliography at the end of the paper.

1.1. THEOREM. *If G/L is semi-simple then ρ_n is an isomorphism of $H^n(G, M)$ onto $H^n(L, M)^G$ for $n=0, 1, 2$.*

Hochschild-Serre's proof of their Theorem 13 depends on the theory of spectral sequences, and it seems desirable to sketch a direct proof of 1.1 here. Accordingly let $G=K+L$ be a Levi decomposition of G .

(1.2) $H^n(L, M)^G = H^n(L, M)^K$ for all $n \geq 0$.

(1.3) Every element of $H^n(L, M)^K$ has a representative cocycle in $C^n(L, M)^K$.

(1.4)² If $df \in C^n(L, M)^K$ for f in $C^{n-1}(L, M)$ then $\exists g$ in $C^{n-1}(L, M)^K$ $\ni df = dg$.

(1.5) Let f be in $Z^n(G, M)$ where $n=1$ or 2 then there exists f^* in $Z^n(G, M) \ni f^*$ is cohomologous to f and

(a) $f^*(k) = 0$ for all k in K (if $n=1$),

(b) $f^*(\gamma_1, \gamma_2) = 0$ if γ_1 or γ_2 is in K (if $n=2$).

PROOF. (a) If $f \in Z^1(G, M)$ then $f|_K \in Z^1(K, M)$ whence $\exists m$ in $C^0(K, M) = M \ni f(k) = k \cdot m$ for all k in K . Put

$$f^*(\gamma) = (f - dm)(\gamma) \text{ for all } \gamma \text{ in } G.$$

(b) If $f \in Z^2(G, M)$, define a Lie algebra extension (E, π) of M by G as follows: $E = (G, M)$ (direct product) as a vector space with $[(\gamma_1, m_1), (\gamma_2, m_2)] = ([\gamma_1, \gamma_2], \gamma_1 \cdot m_2 - \gamma_2 \cdot m_1 + f(\gamma_1, \gamma_2))$ for all γ_1, γ_2 in G, m_1, m_2 in M ; π the projection of E onto G .

Now $G=K+L$ with K semi-simple. $\pi^{-1}(L)$ is an ideal of E containing M and $E/\pi^{-1}(L) \cong K$. Thus we can find a linear map $\sigma: G \rightarrow E \ni \pi\sigma(\gamma) = \gamma$ for all γ in G and $\sigma|_K$ is a Lie algebra isomorphism of K into E . Consider $\pi^{-1}(L)$ and M as $\sigma(K)$ -modules in the regular representation. Then since $\sigma(K)$ is semi-simple, there exists a submodule H of $\pi^{-1}(L) \ni \pi^{-1}(L) = H + M$ and $H \cap M = (0)$. $\pi|_H$ is a vector space isomorphism of H onto L so we see that we can choose the linear map σ so that

(1) $\sigma(k_1, k_2) = [\sigma(k_1), \sigma(k_2)]$ for all k_1, k_2 in K .

(2) $E = \sigma(K) + \sigma(L) + M$ direct sum as $\sigma(K)$ -modules. Now note that

$$[\sigma(k), \sigma(x)] - \sigma[k, x] \in \sigma(L) \rightarrow M = (0) \text{ for all } k \text{ in } K, x \text{ in } L.$$

The cocycle $f^* \ni f^*(\gamma_1, \gamma_2) = [\sigma(\gamma_1), \sigma(\gamma_2)] - \sigma[\gamma_1, \gamma_2]$ is cohomologous to f and has the desired properties.

PROOF OF 1.1. If $n=0$ this is trivial. If $n=1$ or 2 and $h \in H^n(G, M)$, choose a representative cocycle f of h as in (1.5). It is easy to see that $k \cdot f|_L = 0$ for all k in K where $f|_L$ is the restriction of f to L so that

² (1.2), (1.3) and (1.4) are proved directly in [1, p. 603].

$\rho_n(H^n(G, M)) \subseteq H^n(L, M)^G$. Using (1.4) one sees quickly that ρ_n is an isomorphism.

To see that ρ_n is onto, take $\bar{h}$ in $H^n(L, M)^G$ ($n=1, 2$) and choose a representative f of $\bar{h}$ in $Z^n(L, M)^K$ by (1.2) and (1.3). Extend f to $\hat{f}$ in $Z^n(G, M)$ by

$$\begin{aligned} n = 1 & \quad \hat{f}(k+x) = f(x) \quad \text{for all } x \text{ in } L, k \text{ in } K; \\ n = 2 & \quad \begin{cases} \hat{f}(k, \gamma) = 0 & \text{for all } \gamma \text{ in } G, k \text{ in } K, \\ \hat{f}(x_1, x_2) = f(x_1, x_2) & \text{for all } x_1, x_2 \text{ in } L \end{cases} \end{aligned}$$

then $\rho_n(\hat{f} + dC^{n-1}(G, M)) = \bar{h}$.

1.6. DEFINITION. Let G be a Lie algebra and V an ideal of G . If there exists a subalgebra U of G such that $G = U + V$ and $U \cap V = (0)$ we say that G splits over V .

As a corollary of 1.2 we have

1.7. Let G be a Lie algebra over a ground field of characteristic 0 and let $L \geq M$ be ideals of G such that G/L is semi-simple, M is abelian and L splits over M . Then G splits over M .

It would be interesting to know if the hypothesis that M be abelian can be removed from 1.7.

2. Module extensions. Let P and Q be G -modules, L any ideal of G . (We drop the requirement that G/L be semi-simple and that the ground field have characteristic 0.) If $P \subseteq E$ and if the sequence $0 \rightarrow P \rightarrow {}^iE \rightarrow {}^oQ \rightarrow 0$ of L -modules is exact we say that the pair (E, ϕ) is an L -extension of P by Q . Given two L -extensions (E_1, ϕ_1) and (E_2, ϕ_2) of P by Q , they are called L -equivalent if there is an L -isomorphism τ of E_1 onto E_2 such that $\tau(p) = p$ for all p in P and such that $\phi_2\tau = \phi_1$. We denote by $\{(E, \phi)\}$ the class of extensions L -equivalent to (E, ϕ) and recall that these classes form a vector space, over the ground field of L , which we denote by $\text{Ext}_L[P, Q]$.

Let M be the vector space of linear maps of Q into P made into a G (and thus also an L)-module by setting

$$(\gamma \cdot f)(q) = \gamma \cdot f(q) - f(\gamma \cdot q) \text{ for all } \gamma \text{ in } G, f \text{ in } M$$

and q in Q . It is known that $\text{Ext}_L[P, Q] \cong H^1(L, M)$ as a vector space [2, p. 562] and we now define an operation of G on $\text{Ext}_L[P, Q]$ which agrees with the operation of G on $H^1(L, M)$ in the sense that this vector space isomorphism becomes a G -isomorphism. Take γ in G , and (E, ϕ) an L -module extension of P by Q . As a vector space, put

$$E_\gamma^* = \{(e_1, e_2) \mid e_1, e_2 \text{ in } E \text{ and } \gamma \cdot \phi(e_1) = \phi(e_2)\}.$$

Make E_γ^* into an L -module such that

$$x \cdot (e_1, e_2) = (x \cdot e_1, x \cdot e_2 + [\gamma, x] \cdot e_1) \quad \text{for } x \text{ in } L \text{ and}$$

(e_1, e_2) in E_γ^* . Define a map ϕ_γ^* of E_γ^* onto Q by $\phi_\gamma^*(e_1, e_2) = \phi(e_1)$. The kernel of ϕ_γ^* is (P, P) and the map $w_\gamma: (P, P) \rightarrow P$ such that

$$w_\gamma(p_1, p_2) = \gamma \cdot p_1 - p_2$$

is an L -homomorphism of (P, P) onto P . The kernel N_γ of w_γ consists of all pairs (p_1, p_2) for which $p_2 = \gamma \cdot p_1$. N_γ is an L -submodule of E_γ^* so that we may use w_γ to induce an L -identification of $(P, P)/N_\gamma$ with P . Finally we put

$$\gamma \cdot (E, \phi) = (E_\gamma^*/N_\gamma, \phi_\gamma^*)$$

[ϕ_γ^* is used again to denote the L -homomorphism of E_γ^*/N_γ induced by the original ϕ_γ^* .]

Suppose now that (E_1, ϕ_1) is an extension of P by Q which is L -equivalent to (E, ϕ) and let ψ be an L -isomorphism of E onto E_1 . The map $\hat{\psi}: E_\gamma^* \rightarrow E_{1\gamma}^*$ such that

$$\hat{\psi}(e_1, e_2) = (\psi(e_1), \psi(e_2))$$

is an L -homomorphism which takes N_γ onto $N_{1\gamma}$ and induces an L -isomorphism of E_γ^*/N_γ onto $E_{1\gamma}^*/N_{1\gamma}$. Thus we may put

$$\gamma\{(E, \phi)\} = \{\gamma \cdot (E, \phi)\}.$$

2.1. PROPOSITION. *If h is the element of $H^1(L, M)$ determined by (E, ϕ) then the element of $H^1(L, M)$ determined by $\gamma \cdot (E, \phi)$ is $\gamma \cdot h$ [$M = \text{linear maps of } Q \text{ into } P$].*

PROOF. Let ψ be a linear map of Q into E such that $\phi\psi(q) = q$ for all q in Q . Then we know that the f in $C^1(L, M)$ such that

$$f(x)\{q\} = x \cdot \psi(q) - \psi(x \cdot q) \quad \text{for all } x \text{ in } L$$

is an element of $Z^1(L, M)$ which represents h . Define a linear map ψ^* of Q into E_γ^*/N_γ by

$$\psi^*(q) = (\psi(q), \psi(\gamma \cdot q)) + N_\gamma.$$

Then $\phi_\gamma^*\psi^*(q) = q$ for all q in Q and

$$x \cdot \psi^*(q) - \psi^*(x \cdot q) = (\gamma \cdot f)(x)\{q\}$$

for all x in L .

2.2. COROLLARY. *The operation of G on $\text{Ext}_L[P, Q]$ induces the structure of a G -module on $\text{Ext}_L[P, Q]$ and the vector space isomorphism*

of $\text{Ext}_L [P, Q]$ onto $H^1(L, M)$ becomes a G -isomorphism.

Now define a map $\rho: \text{Ext}_G [P, Q] \rightarrow \text{Ext}_L [P, Q]$ as follows. If (E, ϕ) is a G -extension of P by Q , define an L -extension $(\bar{E}, \bar{\phi})$ of P by Q by restricting the operators on E to elements of L . As an L -module, $\bar{E} \equiv E$ and as an L -homomorphism $\bar{\phi} \equiv \phi$. It is evident that if h is the element of $H^1(G, M)$ determined by $\{(E, \phi)\}$ then $\rho_1(h)$ is the element of $H^1(L, M)$ determined by $\rho(\{(E, \phi)\})$ where ρ_1 is the restriction homomorphism. Thus we have:

2.3. PROPOSITION. $\rho(\text{Ext}_G [P, Q]) \subseteq \text{Ext}_L [P, Q]^G$.

Now we have the diagram:

$$\begin{array}{ccc} \text{Ext}_G [P, Q] & \xrightarrow{\rho} & \text{Ext}_L [P, Q]^G \\ \uparrow & & \uparrow \\ H(G, M) & \xrightarrow{\rho_1} & H^1(L, M)^G \end{array}$$

where ρ and ρ_1 are homomorphisms into. Further this diagram is commutative.

It is easy to find examples showing that ρ_1 (thus also ρ) need not be an isomorphism in general. Moreover, the following example (due to Hochschild) shows that ρ_1 need not even be onto in general: G is 3 dimensional, with basis x, y, z such that $[x, y] = z$ and z is central. Put $L = (z)$, M a trivial 1 dimensional G -module. Then every 1-cocycle for G in M must map L into (0) ; but the map f , where $f(az) = a$, represents a nonzero element of $H^1(L, M)^G$.

In fact it is an easy consequence of 1.1 and Hochschild-Serre's Theorem 6 [1, p. 596] that

2.4. PROPOSITION. *If the ground field of G has characteristic 0 then $\rho_1: H^1(G, M) \rightarrow H^1(L, M)^G$ is an isomorphism for all G -modules M if and only if G/L is semi-simple.*

3. **Lie algebra extensions.** Let L be an ideal of G and M a G -module which we consider as an abelian Lie algebra. An extension of M by L is a pair (U, π) such that M is an ideal of U and such that the sequence $0 \rightarrow M \rightarrow U \rightarrow U/M \rightarrow 0$ is exact. Two such extensions (U_1, π_1) and (U_2, π_2) are called equivalent if there is an isomorphism $\sigma: U_1 \rightarrow U_2$ such that $\sigma(m) = m$ for all m in M and $\pi_1 = \pi_2 \sigma$. Using a procedure analogous to the Baer product, one can define directly the structure of a vector space on the classes of extensions $\{(U, \pi)\}$. We denote this vector space by $\text{Ext}(M, L)$. Further it is known, [2, p. 566], that this vector space is isomorphic to $H^2(L, M)$.

We define an operation of G on $\text{Ext}(L, M)$ as follows: take γ in

G and (U, π) an extension of M by L . Put

$$U_\gamma = \{ \langle u_1, u_2 \rangle \mid u_1, u_2 \text{ in } U, [\gamma, \pi(u_1)] = \pi(u_2) \}$$

as a vector space and make U_γ into a Lie algebra by setting

$$[\langle u_1, u_2 \rangle, \langle u'_1, u'_2 \rangle] = \langle [u_1, u'_1], [u_2, u'_2] \rangle + \langle [u_1, u'_2], [u_2, u'_1] \rangle$$

for all $\langle u_1, u_2 \rangle$ and $\langle u'_1, u'_2 \rangle$ in U_γ . Define a homomorphism π_γ of U_γ onto L by $\pi_\gamma \langle u_1, u_2 \rangle = \pi(u_1)$. The kernel of π_γ is $\langle M, M \rangle$. This situation makes $\langle M, M \rangle$ into an L -module with L operating on $\langle M, M \rangle$ via U_γ . The map $\omega_\gamma: \langle M, M \rangle \rightarrow M$ such that

$$\omega_\gamma \langle m_1, m_2 \rangle = \gamma \cdot m_1 - m_2$$

is an L -homomorphism of $\langle M, M \rangle$ onto M . The kernel M_γ of ω_γ consists of pairs $\langle m, \gamma \cdot m \rangle$ and M_γ is an ideal of U_γ . Thus π_γ induces a homomorphism (still denoted by π_γ) of U_γ/M_γ onto L with kernel $\langle M, M \rangle/M_\gamma$ which we identify with M using ω_γ . Now define

$$\gamma \cdot (U, \pi) = (U_\gamma/M_\gamma, \pi_\gamma).$$

If (U_1, π_1) is equivalent to (U, π) under the isomorphism ψ , the map ψ^* such that $\psi^* - \langle u_1, u_2 \rangle = \langle \psi(u_1), \psi(u_2) \rangle$ induces an equivalence isomorphism of $(U_\gamma/M_\gamma, \pi_\gamma)$ with $(U_{1\gamma}/M_{1\gamma}, \pi_{1\gamma})$ so that we may put

$$\gamma \cdot \{ (U, \pi) \} = \{ \gamma \cdot (U, \pi) \}.$$

3.1. PROPOSITION. *Let h be the element of $H^2(L, M)$ determined by (U, π) . Then the element of $H^2(L, M)$ determined by $\gamma \cdot \{ (U, \pi) \}$ is $\gamma \cdot h$.*

PROOF. Let α be a linear map of L into U such that $\pi\alpha(x) = x$ for all x in L . The element f of $C^2(L, M)$ such that

$$f(x_1, x_2) = [\alpha(x_1), \alpha(x_2)] - \alpha([x_1, x_2])$$

is in $Z^2(L, M)$ and represents h . Define a linear map $\hat{\alpha}: L \rightarrow U_\gamma/M_\gamma$ by

$$\hat{\alpha}(x) = \langle \alpha(x), \alpha([\gamma, x]) \rangle + M_\gamma.$$

Then

$$[\hat{\alpha}(x_1), \hat{\alpha}(x_2)] - \hat{\alpha}[x_1, x_2] = (\gamma \cdot f)(x_1, x_2) \quad \text{for all } x_1, x_2 \text{ in } L.$$

3.2. COROLLARY. *The operation of G on $\text{Ext}(M, L)$ induces the structure of a G -module on $\text{Ext}(M, L)$ and the vector space isomorphism of $\text{Ext}(M, L)$ onto $H^2(L, M)$ becomes a G -isomorphism.*

Next consider $\text{Ext}(M, G)$. If $\{ (U, \pi) \}$ is in $\text{Ext}(M, G)$ put

$$\rho \{ (U, \pi) \} = \{ (\pi^{-1}(L), \bar{\pi}) \}$$

in $\text{Ext}(M, L)$ where $\bar{\pi}$ is the restriction of π to $\pi^{-1}(L)$. ρ is a homomorphism of $\text{Ext}(M, G)$ into $\text{Ext}[M, L]$ and it is evident that the cohomology class in $H^2(L, M)$ belonging to $\rho\{(U, \pi)\}$ is $\rho_2(h)$ where g is the cohomology class in $H^2(G, M)$ belonging to $\{(U, \pi)\}$ and ρ_2 is the restriction homomorphism.

3.3. PROPOSITION. $\rho \text{ Ext } [M, G] \subseteq \text{Ext } [M, L]^G$.

PROOF. $\bar{U} = \pi^{-1}(L)$ is an ideal in U . Let $\bar{\pi} = \pi|_{\bar{U}}$ then $\gamma \cdot (\bar{U}, \bar{\pi})$ is equivalent to the split extension (L, M) of M by L . In fact, the map

$$\langle \bar{u}_1, \bar{u}_2 \rangle + M_\gamma \rightarrow (\bar{\pi}(\bar{u}_1), \bar{u}_2 - [\gamma^1, \bar{u}_1])$$

$$(\bar{u}_i \text{ in } \bar{U}, \gamma' \text{ in } \bar{U} \text{ such that } \pi(\gamma') = \gamma)$$

of $\bar{U}_\gamma/M_\gamma$ onto (L, M) is an equivalence isomorphism.

Again we have a commutative diagram:

$$\begin{array}{ccc} \text{Ext } (M, G) & \xrightarrow{\rho} & \text{Ext } (M, L)^G \\ \uparrow & & \uparrow \\ H^2(\bar{G}, M) & \xrightarrow{\rho_1} & H^2(\bar{L}, M)^G \end{array}$$

where ρ and ρ_2 are homomorphisms into. If G/L is semi-simple and if the ground field has characteristic 0 we know that ρ_2 is an isomorphism onto by 1.1. Thus

3.5. PROPOSITION. *If G/L is semi-simple and if the ground field of G has characteristic 0, ρ is an isomorphism of $\text{Ext } (M, G)$ onto $\text{Ext } (M, L)^G$.*

4. Lie algebra kernels. As before, let G be a Lie algebra, L an ideal of G and M a G -module. The derivations of L form a Lie algebra (denoted by $D(L)$) and the inner derivations form an ideal of $D(L)$ denoted by $I(L)$. An L -kernel with nucleus M is a pair (K, ϕ) consisting of a Lie algebra K and a homomorphism ϕ of L into $D(K)/I(K)$, M is required to be the center of K and ϕ must induce the given L -module structure on M .

Hochschild [3, p. 699 et seq.], has defined a rule of composition of L -kernels and a partitioning of the set of L -kernels with nucleus M into similarity classes such that the resulting set of similarity classes of kernels, $\text{Kern } [L, M]$, becomes a vector space over the same ground field as L . It is not difficult to show that each element of $\text{Kern } [L, M]$ determines a unique element of $H^3(L, M)$ and that this correspondence is in fact a vector space isomorphism of $\text{Kern } [L, M]$ into $H^3(L, M)$. This isomorphism is not onto in general and again we refer to [3] and [4] for the details.

Here we shall define directly an operation of G on $\text{Kern } [L, M]$ which is "correct" in the sense that if the similarity class $\{[K, \phi]\}$ determines h in $H^3(L, M)$ then $\gamma \cdot \{(K, \phi)\}$ determines $\gamma \cdot h$. An immediate consequence of this (at least if the ground field has characteristic 0) is that the effaceable elements of $H^3(L, M)$ constitute a G -submodule of $H^3(L, M)$. See [4, Theorem 5, p. 777].

In order to do this, let (K, ϕ) be an L -kernel with nucleus M where L is an ideal of G and take γ in G . Form the Lie algebra K_γ of pairs $\langle k_1, k_2 \rangle$ with k_1, k_2 in K and put

$$\langle \langle k_1, k_2 \rangle, \langle k'_1, k'_2 \rangle \rangle = \langle [k_1, k'_1], [k_1, k'_2] + [k_2, k'_1] \rangle.$$

K_γ has center $\langle M, M \rangle$ and we define a homomorphism $t_\gamma: \langle M, M \rangle \rightarrow M$ such that $t_\gamma \langle m_1, m_2 \rangle = \gamma \cdot m_1 + m_2$. The kernel T_γ of t_γ consists of pairs $\langle m, -\gamma \cdot m \rangle$ with m in M .

Now form K_γ/T_γ . The center of K_γ/T_γ is $\langle M, M \rangle/T_\gamma$; for if $[\langle a_1, a_2 \rangle, \langle k_1, k_2 \rangle]$ is in T_γ for all $\langle k_1, k_2 \rangle$ in K_γ then $[a_1, k_1]$ is in M for all k_1 in K and $[a_1, k_2] + [a_2, k_1] = -\gamma \cdot [a_1, k_1]$. In particular, for pairs of the form $\langle m, k \rangle$ with m in M , we get

$$[a_1, k] + [a_2, m] = -\gamma \cdot [a_1, m] = 0$$

whence $[a_1, k] = 0$ for all k in K so that a_1 is in M . Thus since $[a_1, k_2] + [a_2, k_1] = -\gamma \cdot [a_1, k_1]$ for all $\langle k_1, k_2 \rangle$, we see that $[a_2, k_1] = 0$ for all k_1 in K i.e. a_2 is in M .

We define a homomorphism ϕ_γ of L into $D(K_\gamma/T_\gamma)/I(K_\gamma/T_\gamma)$ as follows: let ρ be any linear map of L into $D(K)$ which is compatible with ϕ (i.e. the class mod $I(K)$ of $\rho(x)$ is $\phi(x)$ for all x in L .) Set

$$\rho^*(x)\{\langle k_1, k_2 \rangle + T_\gamma\} = \langle \rho(x)\{k_1\}, \rho(x)\{k_2\} - \rho([\gamma, x])\{k_1\} \rangle + T_\gamma.$$

It is easy to check that ρ^* is a linear map of L into $D(K_\gamma/T_\gamma)$ and that the class mod $I(K_\gamma/T_\gamma)$ of $\rho^*(x)$ does not depend on the choice of ρ compatible with ϕ . Now we note that if

$$[\rho(x_1), \rho(x_2)] - \rho([x_1, x_2]) = D_{\tau(x_1, x_2)} \quad \text{in } I(K) \text{ for } x_1, x_2 \text{ in } L,$$

then

$$[\rho^*(x_1), \rho^*(x_2)] - \rho^*([x_1, x_2]) = D_{\tau^*(x_1, x_2)} \quad \text{in } I(K_\gamma/T_\gamma)$$

where $\tau^*(x_1, x_2) = \langle \tau(x_1, x_2), -\tau([\gamma, x_1], x_2) - \tau(x_1, [\gamma, x_2]) \rangle + T_\gamma$. Now put $\phi_\gamma(x) = \phi^*(x) + I(K_\gamma/T_\gamma)$. One sees immediately that ϕ_γ induces the given L -module structure on the center $\langle M, M \rangle/T_\gamma$ and that t_γ induces an L -isomorphism of $\langle M, M \rangle/T_\gamma$ onto M so that the pair $(K_\gamma/T_\gamma, \phi_\gamma)$ is an L -kernel with nucleus M which we denote by $\gamma \cdot (K, \phi)$.

The function $\tau: L \times L \rightarrow K$ is called a hindrance of (K, ϕ) and its (formal) coboundary $\delta\tau$ is an element of $Z^3(L, M)$ called a primary deviation. The cohomology class of $\delta\tau$ (called the obstruction of (K, ϕ)) is independent of the choice of ρ compatible with ϕ and of the choice of τ such that $[\rho(x_1), \rho(x_2)] - \rho([x_1, x_2]) = D_{\tau(x_1, x_2)}$.

In this terminology, $\tau^*(x_1, x_2)$ is a hindrance of $\gamma \cdot (K, \phi)$ and a straightforward computation shows that

$$\delta\tau^* = \gamma \cdot \delta\tau.$$

Since two L -kernels have the same obstruction if and only if they are similar, we have

4.1. PROPOSITION. *The map $\{(K, \phi)\} \rightarrow \{\gamma \cdot (K, \phi)\}$ (where braces indicate similarity class) induces the structure of a G -module on $\text{Kern}[L, M]$ and further, if h is the obstruction of $\{(K, \phi)\}$, then $\gamma \cdot h$ is the obstruction of $\{\gamma \cdot (K, \phi)\}$.*

As before we define a map ρ of $\text{Kern}[G, M]$ into $\text{Kern}[L, M]$ by setting $\rho\{(\hat{K}, \hat{\phi})\} = \{(\hat{K}, \hat{\phi}|_L)\}$ where $\hat{\phi}|_L$ is the restriction of $\hat{\phi}$ to L . It is evident that if h is the obstruction of $\{(\hat{K}, \hat{\phi})\}$ then $\rho_3(h)$ is the obstruction of $\{(\hat{K}, \hat{\phi}|_L)\}$ where ρ_3 is the restriction map of $H^3(G, M)$ into $H^3(L, M)$. Again we have a commutative diagram:

$$\begin{array}{ccc} H^3(G, M) & \rightarrow & H^3(L, M)^G \\ \downarrow & & \downarrow \\ \text{Kern}[G, M] & \rightarrow & \text{Kern}[L, M]^G \end{array}$$

APPENDIX

For details concerning the following discussion the reader is referred to [5].

A. Group enlargements.

DEFINITION. Let G be an arbitrary group (written multiplicatively) and P, Q groups on which G operates from the left. In this section, P is assumed commutative but Q need not be. A G -enlargement of P by Q is a pair (E, ϕ) such that

- (i) G operates on E from the left,
- (ii) P is a subgroup of E and a direct summand of E (as a group),
- (iii) ϕ is a G -homomorphism of E onto Q with kernel P .

Two G -enlargements $(E_1, \phi_1), (E_2, \phi_2)$ are called *equivalent* if there exists a G -isomorphism $\tau: E_1 \rightarrow E_2$ such that $\tau(p) = p$ for all p in P and $\phi_2\tau = \phi_1$.

Eilenberg gives a definition of multiplication of equivalence classes

of G -enlargements so that these classes form an abelian group denoted by $\text{Enl}(G, Q, P)$. If θ is an element of $\text{Hom}(Q, P)$ and if one puts $(\sigma \cdot \theta)\{q\} = \sigma \cdot \theta(\sigma^{-1}q)$ for all σ in G and q in Q then $\text{Hom}(Q, P)$ becomes a G -module and Eilenberg then showed that $\text{ENL}(G, Q, P) \cong H^1(G, \text{Hom}(Q, P))$.

Now suppose that G is a normal subgroup of a group Γ and that Γ also operates on P and Q . Let (E, ϕ) denote a G -enlargement of P by Q . Take γ in Γ and let E_γ be the subgroup of $E \times E$ consisting of pairs (e_1, e_2) such that $\phi(e_1) = \gamma \cdot \phi(e_2)$. Make E_γ into a G -module such that $\sigma \cdot (e_1, e_2) = (\sigma \cdot e_1, \sigma^\gamma e_2)$ where $\sigma^\gamma = \gamma^{-1} \sigma \gamma$ for all σ in G . Let ϕ_γ^* denote the G -homomorphism $(e_1, e_2) \rightarrow \phi(e_1)$ of E_γ onto Q . The kernel of ϕ_γ^* is (P, P) and the map $\alpha_\gamma: (P, P) \rightarrow P$ such that $\alpha_\gamma(p_1, p_2) = \gamma \cdot p_2$ is a G -homomorphism of (P, P) whose kernel $\ker(\alpha_\gamma)$ consists of pairs $(p, 1)$ where 1 is the identity of P . ϕ_γ^* induces a G -homomorphism ϕ_γ of $E_\gamma/\ker(\alpha_\gamma)$ onto Q so that the pair $(E_\gamma/\ker(\alpha_\gamma), \phi_\gamma)$ is a G -enlargement of P by Q which we denote by $\gamma \cdot (E, \phi)$. It is not difficult to show that if (E, ϕ) is equivalent to (E', ϕ') then $\gamma \cdot (E, \phi)$ is equivalent to $\gamma \cdot (E', \phi')$ and that this operation induces the structure of a Γ -module on $\text{Enl}(G, Q, P)$. Further one can check that this structure is compatible with the Γ -module structure on $H^1(G, \text{Hom}(Q, P))$ so that the natural isomorphism of $\text{Enl}(G, Q, P)$ onto $H^1(G, \text{Hom}(Q, P))$ becomes a Γ -isomorphism.

B. Group extensions. If M is a subgroup of U and if the sequence $0 \rightarrow M \rightarrow U \rightarrow \pi G \rightarrow 0$ is exact then we say that the pair (U, π) is an extension of G by M . If M is abelian, this situation makes M into a G -module. Two such extensions (U_1, π_1) and (U_2, π_2) of G by M are called equivalent if there exists an isomorphism $\tau: U_1 \rightarrow U_2$ such that $\tau(m) = m$ for all m in M and such that $\pi_1 = \pi_2 \tau$. The equivalence classes of such extensions form an abelian group $A(G, M)$ which is isomorphic with $H^2(G, M)$ see [5].

Now suppose that G is a normal subgroup of a group Γ and that M is also a Γ -module. Let (U, π) be an extension of G by M and take γ in Γ . Form $U_\gamma = \{\langle u_1, u_2 \rangle \mid u_i \text{ in } U, \pi(u_1) = \pi(u_2)^\gamma\}$ with $\langle u_1, u_2 \rangle \langle u'_1, u'_2 \rangle = \langle u_1 u'_1, u_2 u'_2 \rangle$. Set $\pi_\gamma \langle u_1, u_2 \rangle = \pi(u_1)$ and note that the kernel of π_γ is $\langle M, M \rangle$. Define a homomorphism $\omega_\gamma: \langle M, M \rangle \rightarrow M$ by $\omega_\gamma \langle m_1, m_2 \rangle = \gamma \cdot m_2$ for all $\langle m_1, m_2 \rangle$ in M . ω_γ is a G -homomorphism of $\langle M, M \rangle$ onto M and the kernel of $\omega_\gamma \equiv \langle M, 0 \rangle$ is a normal subgroup of $U_\gamma \cdot \pi_\gamma$ induces a homomorphism (still denoted by π_γ) of $U_\gamma/\langle M, 0 \rangle$ onto M and we put

$$\gamma \cdot (U, \pi) = (U_\gamma/\langle M, 0 \rangle, \pi_\gamma).$$

One can check that if (U, π) is equivalent to (U_1, π_1) then $\gamma \cdot (U, \pi)$

is equivalent to $\gamma \cdot (U_1, \pi_1)$ and that this operation induces the structure of a Γ -module on $A(G, M)$. Further, if h is the element of $H^2(G, M)$ belonging to (U, π) then $\gamma \cdot h$ belongs to $\gamma \cdot (U, \pi)$ so that the isomorphism of $A(G, M)$ onto $H^2(G, M)$ becomes a Γ -isomorphism.

C. Group kernels. A G -kernel is a pair (K, θ) where θ is a homomorphism: $G \rightarrow A(K)/I(K)$ (the outer automorphisms of K). If M is the center of K , then θ makes M into a G -module called the nucleus of (K, θ) . Eilenberg-MacLane [5, p. 12] have given a rule of composition of G -kernels with fixed nucleus M and a partitioning of the set of G -kernels with nucleus M into similarity classes such that the resulting set of similarity classes $\mathcal{K}(G, M)$ becomes an abelian group. They then show that $\mathcal{K}(G, M)$ is isomorphic to $H^3(G, M)$.

Now suppose that G is a normal subgroup of a group Γ and that M is also a Γ -module. Let (K, θ) be a G -kernel with nucleus M and take γ in Γ . As a group, put $K_\gamma = K$ and put $\theta_\gamma(\sigma) = \theta(\gamma^{-1}\sigma\gamma)$ for all σ in G . The center M_γ of K_γ is still M (as a group) but now the operation of G on M_γ is given by

$$\sigma * m = \gamma^{-1}\sigma\gamma \cdot m \quad \text{for all } \sigma \text{ in } G, m \text{ in } M.$$

The map $m \rightarrow \gamma \cdot m$ is a G -isomorphism of M_γ onto M which we use to identify M_γ with M as a G -module. Then the pair $(K_\gamma, \theta_\gamma)$ is a G -kernel with nucleus M which we denote by $\gamma \cdot (K, \theta)$. One can check that this operation induces the structure of a Γ -module on $\mathcal{K}(G, M)$ and sees that if h is the element of $H^3(G, M)$ belonging to (K, θ) then $\gamma \cdot h$ belongs to $\gamma \cdot (K, \theta)$ so that the isomorphism of $\mathcal{K}(G, M)$ with $H^3(G, M)$ becomes a Γ -isomorphism.

BIBLIOGRAPHY

1. G. Hochschild and J.-P. Serre, *Cohomology of Lie algebras*, Ann. of Math. vol. 57 (1953) pp. 591-603.
2. G. Hochschild, *Cohomology of restricted Lie algebras*, Amer. J. Math. vol. 66 (1954) pp. 555-580.
3. ———, *Lie Algebra kernels and cohomology*, Amer. J. Math. vol. 66 (1954) pp. 698-716.
4. ———, *Cohomology classes of finite type and finite dimensional kernels for Lie algebras*, Amer. J. Math. vol. 66 (1954) pp. 763-778.
5. Samuel Eilenberg, *Topological methods in abstract algebra*, Bull. Amer. Math. Soc. vol. 55 (1949) pp. 3-37.

SYRACUSE UNIVERSITY AND
UNIVERSITY OF PITTSBURGH