ON THE GIBBS' PHENOMENON IN A CERTAIN EIGENFUNCTION SERIES

LUNA I. MISHOE

I. Introduction. In a paper by B. Friedman and Luna Mishoe [1] it is shown that the series,

$$(1) \quad \sum_{-\infty}^{\infty} a_n u_n(x);$$

where

$$a_n = \left(\int_0^1 u_n(\xi) B^* v_n(\xi) d\xi \right)^{-1} \int_0^1 f(\xi) B^* v_n(\xi) d\xi; \quad u_n(x)$$

being eigenfunctions or nonzero solutions of the system

$$(2) \quad u'' + q(x)u + \lambda[p(x)u - u'] = 0, \quad u(0) = u(1) = 0$$

and $v_n(x)$ being eigenfunctions of the system adjoint to (2), where $B^* = d/dx + p(x)$; converges to;

$$(3) \quad 1/2[f(x+) + f(x-)] + c \exp \int_0^x p dt;$$

provided $f(x)$ is of bounded variation, and $p(x)$ possesses a continuous second derivative, and $q(x)$ is continuous on $(0, 1)$. In the expression (3) above

$$(4) \quad c = -1/2 \left[f(0+) + f(1-) \exp - \int_0^1 p dt \right].$$

In this paper the following theorem is proved:

THEOREM. *The partial sum $\sigma_n(x) \equiv \sum_{k=-n}^n a_k u_k(x)$ exhibits Gibbs' phenomenon for the function $g(x) \equiv f(x) + c \exp \int_0^x p dt$ whenever the Fourier partial sum $S_n(x) \equiv \sum_{k=-n}^n \exp(2k\pi i x) \int_0^1 f(\xi) \exp(-2k\pi i \xi) d\xi$ exhibits Gibbs' phenomenon for $f(x)$.*

II. A relationship between $\sigma_n(x)$ and $S_n(x)$. In [1] it is shown that:

$$(5) \quad u_n(x) = \lambda_n^{-1} \left[\exp \left(\lambda_n x - \int_0^x p dt \right) - \exp \left(\int_0^x p dt \right) \right] + O \left(\frac{1}{\lambda_n^2} \right),$$

$$B^* v_n(x) = \exp \left(-\lambda_n x + \int_0^x p dt \right) + O \left(\frac{e^{-\lambda_n x}}{\lambda_n} \right)$$

Received by the editors April 29, 1957.

where $O(1)$ means a bounded function as $|\lambda| \rightarrow \infty$.

In [2] it is shown that:

$$(6) \quad \lambda_n = 2n\pi i + \int_0^1 p dt + O\left(\frac{1}{n}\right).$$

From (5) and (6) it follows that

$$(7) \quad \int_0^1 u_n(\xi) B^* v_n(\xi) d\xi = \int_0^1 \frac{1}{\lambda} d\xi + \int_0^1 O\left(\frac{1}{n}\right) d\xi \\ - \int_0^1 \exp\left(-\lambda_n \xi + \int_0^\xi p dt\right) d\xi = \frac{1 + O(1/\lambda_n)}{\lambda_n}$$

and consequently

$$(8) \quad \left[\int_0^1 u_n(\xi) B^* v_n(\xi) d\xi \right]^{-1} = \frac{\lambda_n}{1 + O(1/n)} = \lambda_n \left[1 + O\left(\frac{1}{n}\right) \right]$$

for large n .

Now in (5) if we use the fact that $\exp(-\int_0^x p dt) = 1 + O(1)$ we have

$$(9) \quad u_n(x) = \lambda_n^{-1} \left[e^{\lambda_n x} + O(1) + O\left(\frac{1}{n}\right) \right]$$

and

$$(10) \quad \int_0^1 f(\xi) B^* v_n(\xi) d\xi = \int_0^1 f(\xi) e^{-\lambda_n \xi} d\xi + O(1).$$

From (8), (9), and (10) we obtain

$$(11) \quad a_n u_n(x) \equiv u_n(x) \left[\int_0^1 u_n(\xi) B^* v_n(\xi) d\xi \right]^{-1} \int_0^1 f(\xi) B^* v_n(\xi) d\xi \\ = e^{\lambda_n x} \int_0^1 f(\xi) e^{-\lambda_n \xi} d\xi + O(1).$$

Using the value of λ_n as given in (6) we have

$$(12) \quad e^{\lambda_n x} = e^{2n\pi i x} \cdot \exp\left(2x \int_0^1 p dt + O\left(\frac{x}{n}\right)\right) = e^{2n\pi i x} + O(1)$$

and consequently

$$(13) \quad a_n u_n(x) = e^{2n\pi i x} \int_0^1 f(\xi) e^{-2n\pi i \xi} d\xi + E_n(x),$$

where

$$E_n(x) = O(1) + O(1/n).$$

And we have,

$$(14) \quad \sigma_n(x) \equiv \sum_{k=-n}^n a_k u_k(x) = \sum_{k=-n}^n e^{2k\pi ix} \int_0^1 f(\xi) e^{-2k\pi i\xi} d\xi + \sum_{k=-n}^n E_k(x)$$

or $\sigma_n(x) = S_n(x) + T_n(x)$ where $S_n(x)$ is the Fourier partial sum of $f(x)$ and $T_n(x) \equiv \sum_{k=-n}^n E_k(x)$. This is our desired relationship.

III. **Uniform convergence of the series** $\sum_{-\infty}^{\infty} E_n(x)$ **in** $(0, 1)$. Since $[1] \sum_{-\infty}^{\infty} a_n u_n(x) = 1[f(x+) + f(x-)]/2 + c \exp \int_0^x p dt$; it follows that

$$(15) \quad \sum_{-\infty}^{\infty} \left[e^{2n\pi ix} \int_0^1 f(\xi) e^{-2n\pi i\xi} d\xi + E_n(x) \right] \\ = 1/2[f(x+) + f(x-)] + c \exp \left(\int_0^x p dt \right)$$

and since

$$\sum e^{2n\pi ix} \int_0^1 f(\xi) e^{-2n\pi i\xi} d\xi = 1/2[f(x+) + f(x-)] \quad \text{for } 0 < x < 1$$

we have

$$(16) \quad \sum_{-\infty}^{\infty} \left[e^{2n\pi ix} \int_0^1 f(\xi) e^{-2n\pi i\xi} d\xi + E_n(x) \right] \\ = c \exp \left(\int_0^x p dt \right) + \sum_{-\infty}^{\infty} e^{2n\pi ix} \int_0^1 f(\xi) e^{-2n\pi i\xi} d\xi$$

and it follows that

$$(17) \quad \sum_{-\infty}^{\infty} E_n(x) = c \exp \left(\int_0^x p dt \right).$$

Now since $p(x)$ possess a continuous second derivative on $(0, 1)$ it follows that $c \exp \left(\int_0^x p dt \right)$ is both continuous and of bounded variation on $(0, 1)$ and therefore may be expanded into a Fourier series which converges uniformly to $c \exp \left(\int_0^x p dt \right)$ for $0 < x < 1$. From this we conclude that

$$(18) \quad \sum_{-\infty}^{\infty} E_n(x) = c \exp \left(\int_0^x p dt \right) \quad \text{uniformly for } 0 < x < 1.$$

IV. Gibbs' Phenomenon. Let $f(x)$ possess a finite discontinuity at $x=a_1$ and let α_n and β_n be the abscissa and ordinate respectively of the first maxima (or minima) of $y=S_n(x)$ to the left or right of $x=a_1$, then as

$$(19) \quad n \rightarrow \infty, \quad \alpha_n \rightarrow a_1.$$

But on the left of a_1

$$\beta_n \rightarrow f(a_1 - 0) + \left[\frac{f(a_1 + 0) - f(a_1 - 0)}{\pi} \int_{\pi}^{\infty} \frac{\sin x}{x} dx \right]$$

and on the right of $x=a_1$

$$\beta_n \rightarrow f(a_1 + 0) - \left[\frac{f(a_1 + 0) - f(a_1 - 0)}{\pi} \int_{\pi}^{\infty} \frac{\sin x}{x} dx \right].$$

This is Gibbs' Phenomenon for the partial sum $S_n(x)$ of the Fourier Series of $f(x)$.

Since $T_n(x) = c \exp(\int_0^x p dt)$ uniformly in $(0, 1)$ as $n \rightarrow \infty$, this partial sum does not exhibit Gibbs' Phenomenon.

Note that corresponding to the maxima (or minima) (α_n, β_n) on the curve, $y=S_n(x)$ there is a point (α_n, β_n^1) on the curve $y^1=\sigma_n(x)$ such that

$$\beta_n^1 = \beta_n + c \exp\left(\int_0^{\alpha_n} p dt\right).$$

Now as $n \rightarrow \infty$, $\alpha_n \rightarrow a_1$

$$\begin{aligned} \beta_n^1 \rightarrow f(a_1 - 0) + \left[\frac{f(a_1 + 0) - f(a_1 - 0)}{\pi} \int_{\pi}^{\infty} \frac{\sin x}{x} dx \right] \\ + c \exp\left(\int_0^{a_1} p dt\right) \end{aligned}$$

on the left of $x=a_1$ and

$$\begin{aligned} \beta_n^1 \rightarrow f(a_1 + 0) - \left[\frac{f(a_1 + 0) - f(a_1 - 0)}{\pi} \int_{\pi}^{\infty} \frac{\sin x}{x} dx \right] \\ + c \exp\left(\int_0^{a_1} p dt\right) \end{aligned}$$

on the right of $x=a_1$. Or as $n \rightarrow \infty$

$$(20) \quad \alpha_n \rightarrow a_1;$$

and since $f(a_1+0) - f(a_1-0) = g(a_1+0) - g(a_1-0)$ where

$$g(x) = f(x) + c \exp\left(\int_0^x p dt\right)$$

we have

$$(21) \quad \beta_n^1 \rightarrow g(a_1-0) + \left[\frac{g(a_1+0) - g(a_1-0)}{\pi} \int_{\pi}^{\infty} \frac{\sin x}{x} dx \right]$$

on the left of $x=a_1$ and

$$(22) \quad \beta_n^1 \rightarrow g(a+0) - \left[\frac{g(a_1+0) - g(a_1-0)}{\pi} \int_{\pi}^{\infty} \frac{\sin x}{x} dx \right]$$

on the right of $x=a_1$, where $g(x) = f(x) + c \exp\left(\int_0^x p dt\right)$. The results in (20), (21), and (22) shows clearly that $y^1 = \sigma_n(x)$ exhibits Gibbs' Phenomenon for

$$g(x) = f(x) + c \exp\left(\int_0^x p dt\right)$$

wherever $y = S_n(x)$ exhibits this Phenomenon for $f(x)$.

REFERENCES

1. Bernard Friedman and L. I. Mishoe, *Eigenfunction expansions associated with a non-self-adjoint differential equation*, Pacific J. Math. vol. 6, no. 2 (1956).
2. L. I. Mishoe and G. C. Ford, *On the uniform convergence of a certain eigenfunction series*, Pacific J. Math. vol. 6, no. 2 (1956).

MORGAN STATE COLLEGE