ON RIESZ SUMMABILITY OF FOURIER SERIES

FU CHENG HSIANG

1. Let $\phi(t)$ be an even function, integrable in Lebesgue sense and periodic with period 2π . Let

$$\phi(t) \sim \frac{1}{2} a_0 + \sum_{n=1}^{\infty} a_n \cos nt.$$

Let $\lambda(x)$ be a continuous function, monotone increasing in $(0, \infty)$ and tends to infinity as $x \rightarrow \infty$. The series $\sum a_n$ is said to be summable $(R, \lambda(\omega), r)$ if $c_r(\omega)/(\lambda(\omega))^r \rightarrow s$ as $\omega \rightarrow \infty$, where

$$c_r(\omega) = \frac{1}{2} (\lambda(\omega) - \lambda(0))^r a_0 + \sum_{n \leq \omega} (\lambda(\omega) - \lambda(n))^r a_n.$$

Throughout this note, we assume $s=0$, and on writing $\lambda(x) = e^{\mu(x)}$, $\mu(x)$ satisfies the following conditions (A):

(i) $\mu(x)$ is differentiable and monotone increasing for $x > A$ and $\mu(x) > 1$ as $x \rightarrow \infty$,

(ii) $\mu'(x)$ is monotone decreasing for $x > A$ and $\mu'(x) \rightarrow 0$ as $x \rightarrow \infty$,

(iii) $(\mu'(x))^2 \geq -2\mu''(x) \geq 0$, $(\mu'(x))^3 \geq -(3\mu'(x)\mu''(x) + \mu'''(x))$ for $x > A$,

$$(iv) \quad U(x) = \int_x^\infty \frac{du}{u\mu(u)} \text{ exists for } x > A$$

and

$$(a) \quad x(\mu'(x))^\Delta U\left(\frac{1}{\mu'(x)}\right) = O(1)$$

for some $0 < \Delta < 2$,

(b) $x^\Delta U(x)$ being monotone increasing for some $0 < \Delta < 2 - \Delta$, $\mu(x)$ may be taken in the following forms:

(1) $(\log x)^{1+\Delta}$ ($\Delta > 0$, $x > A > 1$), (2) x^α ($0 < \alpha < 1/2$, $x > A > 0$),

(3) $x^\alpha (\log x)^{-\Delta}$ ($\Delta > 0$, $0 < \alpha < 1/2$, $x > A > 1$).

Hardy and Littlewood have derived a convergence test for the oscillating series $\sum a_n$ [2; 3; 4] which contains two conditions: one is

Received by the editors April 8, 1957 and, in revised form, June 21, 1957.

concerning the behavior of $\phi(t)$ near the point $t=0$, e.g., the continuity condition. The other is concerning the magnitude of a_n , e.g., the coefficients condition. Morgan [8] developed Hardy-Littlewood's result so as to permit varying types of conditions on $\phi(t)$ and a_n . Wang [5; 10; 11] proved Hardy-Littlewood's test in one-sided Tauberian condition for a_n . Sunouchi [9] joined the results of Morgan and Wang and established another type of convergence test for the series $\sum a_n$. Jurkat [6] developed Sunouchi's result as same as Hardy-Littlewood's to further steps. Very recently, Kanno [7] has extended Wang's theorem to the fractional integral $\phi_\alpha(t)$ ($\alpha \geq 1$) of $\phi(t)$. In the present note, we give the above investigations an exhaustive solution. Our result includes, however, to some degrees, the works of the previous authors. It is known that [10], for $r > 0$, $\omega > 0$,

$$c_r(\omega) = \frac{2}{\pi} \int_0^\pi \phi(t) E_r(\omega, t) dt + o((\lambda(\omega))^r),$$

where

$$E_r(\omega, t) = \frac{r}{t} \int_0^\omega (\lambda(\omega) - \lambda(x))^{r-1} \lambda'(x) \sin x t dx.$$

In addition, we define:

$$\lambda(x) = \begin{cases} e^{\mu(x)} & (x > A), \\ \lambda(A) + \lambda'(A)(x - A) & (0 \leq x \leq A). \end{cases}$$

2. The following lemmas are required.

LEMMA 1. For $\omega > A$,

$$E_2(\omega, t) = O(e^{2\mu(\omega)} \omega)$$

uniformly in t ,

$$E_2(\omega, t) = O(e^{\mu(\omega)} t^{-2}) + O(e^{2\mu(\omega)} \mu'(\omega) t^{-2}),$$

$$\frac{\partial}{\partial t} E_2(\omega, t) = O(e^{2\mu(\omega)} \omega t^{-1}),$$

and

$$\frac{\partial}{\partial t} E_2(\omega, t) = O(e^{\mu(\omega)} t^{-3}) + O(e^{2\mu(\omega)} (\mu'(\omega))^2 \omega t^{-3}).$$

PROOF. By the second mean value theorem,¹

¹ B is a numerical constant not necessarily the same at each occurrence.

$$\begin{aligned}
E_2(\omega, t) &= \int_0^\omega (\lambda(\omega) - \lambda(x))^2 \cos xtdx \\
&= (\lambda(\omega) - B)^2 \int_0^{\omega'} \cos xtdx \\
&= O(e^{2\mu(\omega)}\omega). \\
\frac{\partial}{\partial t} E_2(\omega, t) &= - \int_0^\omega (\lambda(\omega) - \lambda(x))^2 x \sin xtdx \\
&= - (\lambda(\omega) - B)^2 \omega \int_{\omega'}^{\omega''} \sin xtdx \\
&= O(e^{2\mu(\omega)}\omega t^{-1})
\end{aligned}$$

for $\omega > A$, by the second mean value theorem, where $0 < \omega' < \omega'' < \omega$.

Now,

$$\begin{aligned}
E_2(\omega, t) &= \frac{2}{t} \left(\int_0^A + \int_A^\omega \right) (\lambda(\omega) - \lambda(x)) \lambda'(x) \sin xtdx \\
&= L_1 + L_2. \\
L_1 &= \frac{2}{t} (\lambda(\omega) - B) \lambda'(A) \int_0^A \sin xtdx = O(e^{\mu(\omega)} t^{-2}), \\
L_2 &= \frac{2}{t} (\lambda(\omega) - B) \lambda'(\omega) \int_{\omega'}^{\omega''} \sin xtdx,
\end{aligned}$$

since, by (iii) of the conditions (A), $\lambda'(x)$ is monotone increasing in (A, ω) . It follows that

$$L_2 = O(e^{2\mu(\omega)} \mu'(\omega) t^{-2}).$$

Thus,

$$E_2(\omega, t) = O(e^{\mu(\omega)} t^{-2}) + O(e^{2\mu(\omega)} \mu'(\omega) t^{-2})$$

for $\omega > A$. Further,

$$\begin{aligned}
\frac{\partial L_1}{\partial t} &= - \frac{2\lambda'(A)}{t^2} \int_0^A (\lambda(\omega) - \lambda(x)) \sin xtdx \\
&\quad + \frac{2\lambda'(A)}{t} \int_0^A (\lambda(\omega) - \lambda(x)) x \cos xtdx \\
&= O(e^{\mu(\omega)} t^{-3}).
\end{aligned}$$

By integration by parts once, we find

$$\begin{aligned}
L_2 &= \frac{2}{t^2} (\lambda(\omega) - \lambda(A))\lambda'(A) \cos At \\
&\quad + \frac{2}{t^2} \int_A^\omega \frac{d}{dx} ((\lambda(\omega) - \lambda(x))\lambda'(x) \cos x) dx \\
&= \frac{2}{t^2} (\lambda(\omega) - \lambda(A))\lambda'(A) \cos At \\
&\quad - \frac{2}{t^2} \int_A^\omega ((\lambda'(x))^2 - (\lambda(\omega) - \lambda(x))\lambda''(x)) \cos x dx.
\end{aligned}$$

Denote $H(\omega, x) = (\lambda'(x))^2 - (\lambda(\omega) - \lambda(x))\lambda''(x)$. Then

$$\begin{aligned}
\frac{\partial L_2}{\partial t} &= \frac{4}{t^3} (\lambda(\omega) - \lambda(A))\lambda'(A) \cos At + \frac{2A}{t^2} (\lambda(\omega) - \lambda(A)) \sin At \\
&\quad + \frac{4}{t^3} \int_A^\omega H(\omega, x) \cos x dx + \frac{2}{t^2} \int_A^\omega H(\omega, x) x \sin x dx \\
&= O(e^{\mu(\omega)} t^{-3}) + (I_1 + I_2 + I_3 + I_4). \\
I_1 &= \frac{4}{t^3} \int_A^\omega (\lambda'(x))^2 \cos x dx = \frac{4}{t^3} (\lambda'(\omega))^2 \int_{\omega'}^\omega \cos x dx \\
&= O((\lambda'(\omega))^2 \omega t^{-3}). \\
I_2 &= -\frac{4}{t^3} \int_A^\omega (\lambda(\omega) - \lambda(x))\lambda''(x) \cos x dx \\
&= -\frac{4}{t^3} (\lambda(\omega) - B)\lambda''(\omega) \int_{\omega'}^{\omega''} \cos x dx,
\end{aligned}$$

since, by (iii) of the conditions (A) , $\lambda''(x)$ is monotone increasing in (A, ω) . It follows that

$$\begin{aligned}
I_2 &= O(t^{-3}(\lambda(\omega) - B)e^{\mu(\omega)}((\mu'(\omega))^2 + \mu''(\omega)\omega)) \\
&= O(e^{2\mu(\omega)}(\mu'(\omega))^2 \omega t^{-3}) + O(e^{2\mu(\omega)} |\mu''(\omega)| \omega t^{-3}).
\end{aligned}$$

In view of $(\mu'(x))^2 \geq -2\mu''(x) \geq 0$ for $x > A$, the second term can be absorbed in the first, so that

$$\begin{aligned}
I_2 &= O(e^{2\mu(\omega)}(\mu'(\omega))^2 \omega t^{-3}). \\
I_2 &= \frac{2}{t^2} \int_A^\omega (\lambda'(x))^2 x \sin x dx \\
&= \frac{2}{t^2} (\lambda'(\omega))^2 \omega \int_{\omega'}^\omega \sin x dx = O((\lambda'(\omega))^2 \omega t^{-3}).
\end{aligned}$$

$$\begin{aligned}
I_4 &= -\frac{2}{t^2} \int_A^\omega (\lambda(\omega) - \lambda(x)) \lambda''(x) x \sin xt dx \\
&= -\frac{2}{t^2} (\lambda(\omega) - B) \lambda''(\omega) \omega \int_{\omega'}^{\omega''} \sin xt dx \\
&= -\frac{2}{t^2} (\lambda(\omega) - B) e^{\mu(\omega)} ((\mu'(\omega))^2 + \mu''(\omega)) \omega \int_{\omega'}^{\omega''} \sin xt dx \\
&= O(e^{2\mu(\omega)} (\mu'(\omega))^2 \omega t^{-3}) + O(e^{2\mu(\omega)} |\mu''(\omega)| \omega t^{-3}) \\
&= O(e^{2\mu(\omega)} (\mu'(\omega))^2 \omega t^{-3}).
\end{aligned}$$

Thus, $(\partial/\partial t)E_2(\omega, t) = O(e^{\mu(\omega)} t^{-3}) + O(e^{2\mu(\omega)} (\mu'(\omega))^2 \omega t^{-3})$ for $\omega > A$. This proves the lemma.

LEMMA 2. If $\sum a_n$ is summable $(R, \lambda(\omega), r)$ and

$$a_n > -K \frac{\lambda(n) - \lambda(n-1)}{\lambda(n)},$$

then $\sum a_n$ converges.

This lemma is known. [1].

3. Now, we are in a position to establish the following

THEOREM 1. If

$$\phi_1(t) = \int_0^t \phi(u) du = o\left(t^\Delta U\left(\frac{1}{t}\right)\right) \quad (t \rightarrow +0),$$

then the series $\sum a_n$ is summable $(R, \lambda(\omega), 2)$ to zero.

PROOF. We have

$$\begin{aligned}
\frac{\pi}{2} c_2(\omega) &= \int_0^\pi \phi(t) E_2(\omega, t) dt + o(e^{2\mu(\omega)}) \\
&= \left(\int_0^{1/A-\epsilon} + \int_{1/A-\epsilon}^\pi \right) \phi(t) E_2(\omega, t) dt + o(e^{2\mu(\omega)}).
\end{aligned}$$

By the second relation of Lemma 1,

$$\begin{aligned}
&\int_{1/A-\epsilon}^\pi \phi(t) E_2(\omega, t) dt \\
&= O\left(e^{\mu(\omega)} \int_{1/A-\epsilon}^\pi \frac{|\phi(t)|}{t^2} dt\right) + O\left(e^{2\mu(\omega)} \mu'(\omega) \int_{1/A-\epsilon}^\pi \frac{|\phi(t)|}{t^2} dt\right) \\
&= O(e^{\mu(\omega)}) + O(e^{2\mu(\omega)} \mu'(\omega)) = o(e^{2\mu(\omega)}),
\end{aligned}$$

since $\mu'(\omega) \rightarrow 0$ as $\omega \rightarrow \infty$. By integration by parts,

$$\int_0^{1/A-\epsilon} \phi(t) E_2(\omega, t) dt = (\phi_1(t) E_2(\omega, t))_0^{1/A-\epsilon} - \int_0^{1/A-\epsilon} \phi_1(t) \frac{\partial}{\partial t} E_2(\omega, t) dt.$$

The integrated term is easily seen to be $o(e^{2\mu(\omega)})$ by the first and the second relations of Lemma 1. Write

$$\begin{aligned} \int_0^{1/A-\epsilon} \phi_1(t) \frac{\partial}{\partial t} E_2(\omega, t) dt &= \left(\int_0^{\mu'(\omega)} + \int_{\mu'(\omega)}^{1/A-\epsilon} \right) \phi_1(t) \frac{\partial}{\partial t} E_2(\omega, t) dt \\ &= J_1 + J_2. \\ J_1 &= o \left(e^{2\mu(\omega)} \omega \int_0^{\mu'(\omega)} t^{\Delta-1} U \left(\frac{1}{t} \right) dt \right) \\ &= o \left(e^{2\mu(\omega)} \omega (\mu'(\omega))^{\Delta} U \left(\frac{1}{\mu'(\omega)} \right) \right) \\ &= o(e^{2\mu(\omega)}) \end{aligned}$$

by (iv) of the conditions (A).

$$\begin{aligned} J_2 &= \int_{\mu'(\omega)}^{1/A-\epsilon} \phi_1(t) \frac{\partial}{\partial t} E_2(\omega, t) dt \\ &= o \left(e^{\mu(\omega)} \int_{\mu'(\omega)}^{1/A-\epsilon} t^{\Delta-3} U(t) dt \right) + o \left(e^{2\mu(\omega)} (\mu'(\omega))^2 \omega \int_{\mu'(\omega)}^{1/A-\epsilon} t^{\Delta-3} U(t) dt \right) \\ &= o \left(e^{\mu(\omega)} (\mu'(\omega))^{-\Delta} U \left(\frac{1}{\mu'(\omega)} \right) \int_{\mu'(\omega)}^{\tau} t^{\Delta+\Delta-3} dt \right) \\ &\quad + o \left(e^{2\mu(\omega)} (\mu'(\omega))^{2-\Delta} \omega U \left(\frac{1}{\mu'(\omega)} \right) \int_{\mu'(\omega)}^{\tau} t^{\Delta+\Delta-3} dt \right) \\ &= o \left(e^{\mu(\omega)} (\mu'(\omega))^{\Delta-2} U \left(\frac{1}{\mu'(\omega)} \right) \right) + o \left(e^{2\mu(\omega)} (\mu'(\omega))^{\Delta} \omega U \left(\frac{1}{\mu'(\omega)} \right) \right) \\ &= o \left(e^{2\mu(\omega)} (\mu(\omega) (\mu'(\omega))^2)^{-1} (\mu'(\omega))^{\Delta} U \left(\frac{1}{\mu'(\omega)} \right) \right) + o(e^{2\mu(\omega)}) \\ &= o(e^{2\mu(\omega)}) \end{aligned}$$

since $e^{\mu(x)} (\mu'(x))^2$ is positive and monotone increasing for $x > A$ by (iii) of the conditions (A). This proves the theorem.

By means of Theorem 1 and Lemma 2, we obtain immediately

THEOREM 2. If

$$(i) \quad \phi_1(t) = o \left(t^{\Delta} U \left(\frac{1}{t} \right) \right) \quad (t \rightarrow +0),$$

$$(ii) \quad a_n > -K \frac{\lambda(n) - \lambda(n-1)}{\lambda(n)}$$

for some $K > 0$, then $\sum a_n$ converges to zero.

4. Last, we illustrate by some special cases derived from our result:

(a) Let $0 < \alpha < 1/2$. If

$$(i) \quad \phi_1(t) = o\left(t^{1/(1-\alpha)}\left(\log \frac{1}{t}\right)^\Delta\right) \quad (t \rightarrow +0),$$

$$(ii) \quad a_n > -Kn^{\alpha-1}(\log n)^{-\Delta}$$

for some $K > 0$, then $\sum a_n$ converges to zero.

(b) Let $0 < \alpha < 1/2$. If

$$(i) \quad \phi_1(t) = o(t^{1/(1-\alpha)}) \quad (t \rightarrow +0),$$

$$(ii) \quad a_n > -Kn^{\alpha-1}$$

then $\sum a_n$ converges to zero.

(c) If

$$(i) \quad \phi_1(t) = o\left(t\left(\log \frac{1}{t}\right)^{-\Delta}\right) \quad (t \rightarrow +0),$$

$$(ii) \quad a_n > Kn^{-1}(\log n)^\Delta$$

for some $\Delta > 0$, then $\sum a_n$ converges to zero.

REFERENCES

1. G. H. Hardy, *An extension of a theorem on the oscillating series*, Proc. London Math. Soc. vol. 12 (1913) pp. 174-180.
2. G. H. Hardy and J. E. Littlewood, *Notes on the theory of series* (XVII): *Some new convergence criteria for Fourier series*, J. London Math. Soc. vol. 7 (1932) pp. 252-256.
3. ———, *Some new convergence criteria for Fourier series*, Annali della Scuola Normale Superiore di Pisa vol. 3 (1934) pp. 1-20.
4. G. H. Hardy and W. W. Rogosinski, *Fourier series*, Cambridge, 1946, pp. 45, 96.
5. F. C. Hsiang, *On a test for the convergence of Fourier series*, Proc. Amer. Math. Soc. vol. 7 (1956) pp. 1036-1039.
6. W. Jurkat, *Zur Convergenztheorie der Fourier-Reihen*, Math. Zeit. vol. 53 (1950) pp. 309-339.
7. K. Kanno, *On the Riesz summability of Fourier series*, Tôhoku Math. J. vol. 8 (1956) pp. 223-234.
8. W. G. Morgan, *On the convergence criteria for Fourier series of Hardy and Littlewood*, Annali della Scuola Normale Superiore di Pisa vol. 4 (1935) pp. 337-382.
9. G. Sunouchi, *Notes on Fourier series: I, On the convergence test of Fourier series*, Mathematica Japonica vol. 1 (1948-49) pp. 41-44.

10. F. T. Wang, *On Riesz summability of Fourier series*, Proc. London Math. Soc. (2) vol. 47 (1942) pp. 308–325.

11. ———, *On Riesz summability of Fourier series* (III), Proc. London Math. Soc. (2) vol. 51 (1949) pp. 215–231.

NATIONAL TAIWAN UNIVERSITY

ON THE IDENTITY OF FUNCTION SPACES ON CARTESIAN PRODUCT SPACES¹

JOHN C. HOLLADAY

For $i = 1, \dots, n$, let S_i be a compact Hausdorff space and F_i be a closed linear subspace of the complex Banach space $C(S_i)$, the set of all continuous functions from S_i to the complex numbers. Let $S_1 \times \dots \times S_n$ be the Cartesian Product of $S_1, \dots, S_n$.

Define $F_1 * \dots * F_n$ as $\{\phi \mid \phi \in C(S_1 \times \dots \times S_n); \text{ for any } i = 1, \dots, n \text{ and } (s_1, \dots, s_{i-1}, s_{i+1}, \dots, s_n), \phi(s_1, \dots, s_{i-1}, \cdot, s_{i+1}, \dots, s_n) \in F_i\}$. Also, define $F_1 \otimes \dots \otimes F_n$ as the closure of the space of linear combinations of functions of the form $\phi(s_1, \dots, s_n) = f_1(s_1) \times \dots \times f_n(s_n)$ where each $f_i \in F_i$, where we base the topology on the norm, $\|\phi\| = \max_{s_1, \dots, s_n} |\phi(s_1, \dots, s_n)|$.

It is easily shown that $F_1 \otimes \dots \otimes F_n$ is a subspace of $F_1 * \dots * F_n$ and that if F_2 is one-dimensional, then $F_1 \otimes F_2 = F_1 * F_2$. Furthermore, by using continuous partitions of unity, one may show that $F_1 \otimes C(S_2) = F_1 * C(S_2)$. Therefore, if all but at most one of the F_i are either one-dimensional or $C(S_i)$, then $F_1 \otimes \dots \otimes F_n = F_1 * \dots * F_n$. However, it is not known whether for all cases $F_1 \otimes \dots \otimes F_n$ will equal $F_1 * \dots * F_n$ or not.² Although this question is not fully answered here, the purpose of this paper is to give a partial answer to this question. The results and arguments of this paper also apply to *real-valued* function spaces.

1. LEMMA. *Let F be a closed linear subspace of $C(S_1)$ and G a closed linear subspace of $C(S_2)$. Let H be a closed subspace of G differing from G by only one dimension. Then $F \otimes G = F * G$ implies that $F \otimes H = F * H$.*

Received by the editors October 22, 1956 and, in revised form, April 16, 1957.

¹ A small part of the work done under an AEC Predoctoral Fellowship at Yale University, year 1952–1953, under the kind and patient guidance of Dr. Charles E. Rickart.

² Under Proposition 37 of his first paper of the Amer. Math. Soc. Memoirs, no. 16, Alexandre Grothendieck discusses a number of conjectures which are equivalent to this one.