

REFERENCES

1. A. B. Simon, *Vanishing algebras*, to appear in Trans. Amer. Math. Soc.
2. E. J. McShane, *Images of sets satisfying the condition of Baire*, Ann. of Math. vol. 51 (1950) pp. 380–386.
3. B. J. Pettis, *On continuity and openness of homomorphisms in topological groups*, Ann. of Math. vol. 52 (1950) pp. 293–308.
4. ———, *Remarks on a theorem of E. J. McShane*, Proc. Amer. Math. Soc. vol. 2 (1951) pp. 166–171.
5. Fred B. Wright, *Semigroups in compact groups*, Proc. Amer. Math. Soc. vol. 7 (1956) pp. 309–311.

TULANE UNIVERSITY

SIMPLE NODAL NONCOMMUTATIVE JORDAN ALGEBRAS

LOUIS A. KOKORIS

1. **Introduction.** Nodal algebras were defined by R. D. Schafer [4] and have also been studied by the author [2; 3]. A noncommutative Jordan algebra is an algebra \mathfrak{A} over a field \mathfrak{F} satisfying (1) the flexible law $(xy)x = x(yx)$ and (2) the condition that \mathfrak{A}^+ is a Jordan algebra. That is, \mathfrak{A}^+ satisfies the identity $(x^2 \cdot y) \cdot x = x^2 \cdot (y \cdot x)$ where we have used the dot to indicate the product of \mathfrak{A}^+ . The algebra \mathfrak{A}^+ is defined to be the same vector space as \mathfrak{A} but with product $x \cdot y = (xy + yx)/2$ where xy and yx are products in \mathfrak{A} . Then \mathfrak{A} is called nodal if it is finite dimensional, if \mathfrak{A} has identity element 1, if \mathfrak{A} can be written as a vector space direct sum $\mathfrak{A} = \mathfrak{F}1 + \mathfrak{N}$ where \mathfrak{N} is the subspace of nilpotent elements of \mathfrak{A} , and if \mathfrak{N} is not a subalgebra of \mathfrak{A} .

Every known nodal algebra \mathfrak{A} has the property that \mathfrak{A}^+ is an associative algebra. The flexible algebras with \mathfrak{A}^+ associative have been described in [3]. In this paper we shall prove the following theorem.

THEOREM 1. *Let \mathfrak{A} be a simple nodal noncommutative Jordan algebra of characteristic $\neq 2$. Then \mathfrak{A}^+ is associative.*

Define \mathfrak{B} to be the subspace of \mathfrak{A} generated by all the associators in \mathfrak{N}^+ . That is, \mathfrak{B} is generated by elements of the form $(x \cdot y) \cdot z - x \cdot (y \cdot z)$ with x, y, z in \mathfrak{N} . The proof of the theorem will be made by showing that the ideal \mathfrak{C} of \mathfrak{A} generated by \mathfrak{B} is not all of \mathfrak{A} and since \mathfrak{A} is simple it will follow that $\mathfrak{C} = 0$ and $\mathfrak{B} = 0$. This is the desired result.

The original proof was not valid when the characteristic is 3. The author thanks Professor R. D. Schafer for suggesting a modification

Presented to the Society, January 28, 1958; received by the editors January 13, 1958.

which makes the proof simpler and also valid when the characteristic is 3.

2. The proof. Let x_1, x_2, x_3 , and y be any elements of \mathfrak{N} . Since \mathfrak{A} is nodal we have:

$$(1) \quad x_i y = \lambda_i 1 + z_i.$$

The proof will depend on relation (8) of Schafer's paper [4] which is

$$(2) \quad \begin{aligned} (x_1 \cdot x_2)y &= \lambda_1 x_2 + \lambda_2 x_1 + x_1 \cdot z_2 + x_2 \cdot z_1 \\ &\quad - (x_1 \cdot y) \cdot x_2 - (x_2 \cdot y) \cdot x_1 + (x_1 \cdot x_2) \cdot y. \end{aligned}$$

In the proof it will also be necessary to use the fact that \mathfrak{N}^+ is a subalgebra of \mathfrak{A}^+ [1].

By (2) $(x_1 \cdot x_2)y$ is in \mathfrak{N} and it follows from (2) that $[(x_1 \cdot x_2) \cdot x_3]y = \lambda_3 x_1 \cdot x_2 + (x_1 \cdot x_2) \cdot z_3 + x_3 \cdot [\lambda_1 x_2 + \lambda_2 x_1 + x_1 \cdot z_2 + x_2 \cdot z_1 - (x_1 \cdot y) \cdot x_2 - (x_2 \cdot y) \cdot x_1 + (x_1 \cdot x_2) \cdot y] - [(x_1 \cdot x_2) \cdot y] \cdot x_3 - (x_3 \cdot y) \cdot (x_1 \cdot x_2) + [(x_1 \cdot x_2) \cdot x_3] \cdot y$. Without bothering to simplify interchange subscripts 1 and 3 to get $[x_1 \cdot (x_2 \cdot x_3)]y = \lambda_1 x_3 \cdot x_2 + (x_3 \cdot x_2) \cdot z_1 + x_1 \cdot [\lambda_3 x_2 + \lambda_2 x_3 + x_3 \cdot z_2 + x_2 \cdot z_3 - (x_3 \cdot y) \cdot x_2 - (x_2 \cdot y) \cdot x_3 - (x_1 \cdot y) \cdot (x_3 \cdot x_2) + [(x_3 \cdot x_2) \cdot x_1] \cdot y$. Using the notation $(a, b, c) = (a \cdot b) \cdot c - a \cdot (b \cdot c)$ for the associator of a, b, c in \mathfrak{A}^+ we have, upon subtracting the second relation from the first, $(x_1, x_2, x_3)y = (x_1, x_2, z_3) + (x_1, z_2, x_3) + (z_1, x_2, x_3) - (x_1 \cdot y, x_2, x_3) - (x_1, x_2 \cdot y, x_3) + (x_3 \cdot y, x_2, x_1) + (x_1, x_2, x_3) \cdot y$.

Now define the set \mathfrak{B} to be the subspace of \mathfrak{A} generated by the associators (a, b, c) with a, b, c in \mathfrak{N} and using the product of \mathfrak{N}^+ . We have proved the following lemma.

LEMMA 1. *Let \mathfrak{A} be a nodal noncommutative Jordan algebra whose characteristic is not 2. Then $\mathfrak{B}\mathfrak{N} \subseteq \mathfrak{B} + \mathfrak{B} \cdot \mathfrak{N}$. Also $\mathfrak{N}\mathfrak{B} \subseteq \mathfrak{B} + \mathfrak{B} \cdot \mathfrak{N}$.*

The last statement follows from the fact that if b is in \mathfrak{B} , n in \mathfrak{N} , then $nb = 2b \cdot n - bn$.

Let $\mathfrak{C}_0 = \mathfrak{B}$, $\mathfrak{C}_1 = \mathfrak{B} + \mathfrak{B} \cdot \mathfrak{N} = \mathfrak{C}_0 + \mathfrak{C}_0 \cdot \mathfrak{N}$, and in general $\mathfrak{C}_{i+1} = \mathfrak{C}_i + \mathfrak{C}_i \cdot \mathfrak{N}$. Equivalently, $\mathfrak{C}_{i+1} = \mathfrak{C}_i + \mathfrak{B}(R_{\mathfrak{N}}^+)^{i+1}$.

LEMMA 2. *The product $(\mathfrak{B} \cdot \mathfrak{N})\mathfrak{N} \subseteq \mathfrak{C}_2$ and $\mathfrak{N}(\mathfrak{B} \cdot \mathfrak{N}) \subseteq \mathfrak{C}_2$. It follows that $\mathfrak{C}_1\mathfrak{N} \subseteq \mathfrak{C}_2$, $\mathfrak{N}\mathfrak{C}_1 \subseteq \mathfrak{C}_2$.*

The proof follows from the flexible law as does (2) which was proved by Schafer. The linearized form of the flexible identity is

$$(3) \quad (xy)z + (zy)x = x(yz) + z(yx).$$

Add $(yx)z + (yz)x$ to both sides of (3) to obtain the equivalent relation

$$(4) \quad (x \cdot y)z + (y \cdot z)x = yz \cdot x + yx \cdot z.$$

If x is in \mathfrak{B} , y, z in \mathfrak{N} , then $(y \cdot z)x$ is in $(\mathfrak{N} \cdot \mathfrak{N})\mathfrak{B} \subseteq \mathfrak{N}\mathfrak{B}$. By Lemma 1, $(y \cdot z)x$ is in \mathfrak{C}_1 . The product yz is in $\mathfrak{F}1 + \mathfrak{N}$ so $yz \cdot x$ is in $\mathfrak{B} + \mathfrak{N} \cdot \mathfrak{B} = \mathfrak{C}_1$. And $yx \cdot z$ is in $\mathfrak{N}\mathfrak{B} \cdot \mathfrak{N} \subseteq \mathfrak{C}_2$. Therefore, $(x \cdot y)z$ is in \mathfrak{C}_2 as desired.

LEMMA 3. *The product $[\mathfrak{B}(R_{\mathfrak{N}}^+)^i]\mathfrak{N} \subseteq \mathfrak{C}_{i+1}$ and $\mathfrak{N}[\mathfrak{B}(R_{\mathfrak{N}}^+)^i] \subseteq \mathfrak{C}_{i+1}$. Or, equivalently, $\mathfrak{C}_i\mathfrak{N} \subseteq \mathfrak{C}_{i+1}$, $\mathfrak{N}\mathfrak{C}_i \subseteq \mathfrak{C}_{i+1}$.*

Assume that $[\mathfrak{B}(R_{\mathfrak{N}}^+)^{i-1}]\mathfrak{N}$ and $\mathfrak{N}[\mathfrak{B}(R_{\mathfrak{N}}^+)^{i-1}]$ are in \mathfrak{C}_i . Take x in (4) to be in $\mathfrak{S} = \mathfrak{B}(R_{\mathfrak{N}}^+)^{i-1}$, and y, z to be in \mathfrak{N} . Then $(y \cdot z)x$ is in $\mathfrak{N}\mathfrak{S} \subseteq \mathfrak{C}_i$, $yz \cdot x$ is in $\mathfrak{S} + \mathfrak{N} \cdot \mathfrak{S} \subseteq \mathfrak{C}_i$, and $yx \cdot z$ is in $(\mathfrak{N}\mathfrak{S}) \cdot \mathfrak{N} \subseteq \mathfrak{C}_{i+1}$. Thus $(x \cdot y)z$ is in \mathfrak{C}_{i+1} .

LEMMA 4. *There exists a positive integer k such that $\mathfrak{C}_k = \mathfrak{C}_{k+1}$ and \mathfrak{C}_k is an ideal of \mathfrak{A} .*

The set \mathfrak{B} is contained in \mathfrak{N} . Since \mathfrak{A}^+ is a Jordan algebra, \mathfrak{N}^+ is nilpotent. Consequently, $\mathfrak{B}(R_{\mathfrak{N}}^+)^{k+1} = 0$ for some k . For this k , $\mathfrak{C}_k = \mathfrak{C}_{k+1}$. By Lemma 3 $\mathfrak{C}_k\mathfrak{N} \subseteq \mathfrak{C}_{k+1} = \mathfrak{C}_k$ and $\mathfrak{N}\mathfrak{C}_k \subseteq \mathfrak{C}_k$.

The ideal \mathfrak{C}_k is contained in $\mathfrak{N} \cdot \mathfrak{N} \cdot \mathfrak{N} = \mathfrak{N}_3$. Since \mathfrak{N}^+ is a subalgebra of \mathfrak{A}^+ , $\mathfrak{N}_3 \subset \mathfrak{N}$ and so $\mathfrak{C}_k \subset \mathfrak{N}$. If \mathfrak{A} is a simple algebra, $\mathfrak{C}_k = 0$ and thus $\mathfrak{B} = 0$. This says that every associator in \mathfrak{N}^+ is zero. Now if a, b, c are any elements in \mathfrak{A} , $a = \alpha 1 + x$, $b = \beta 1 + y$, $c = \gamma 1 + z$ with x, y, z in \mathfrak{N} . Then $(a \cdot b) \cdot c - a \cdot (b \cdot c) = (x \cdot y) \cdot z - x \cdot (y \cdot z)$ so every associator in \mathfrak{A}^+ is an associator in \mathfrak{N}^+ . This completes the proof of the theorem.

Any ideal properly contained in \mathfrak{A} is contained in \mathfrak{N} , hence is a nilideal and is contained in the radical of \mathfrak{A} . This implies the corollary which we state below.

COROLLARY. *Let \mathfrak{A} be a semisimple nodal noncommutative Jordan algebra of characteristic $\neq 2$. Then \mathfrak{A} is simple and \mathfrak{A}^+ is associative.*

REFERENCES

1. N. Jacobson, *A theorem on the structure of Jordan algebras*, Proc. Nat. Acad. Sci. U.S.A. vol. 42 (1956) pp. 140-147.
2. L. A. Kokoris, *Some nodal noncommutative Jordan algebras*, Proc. Amer. Math. Soc. vol. 9 (1958) pp. 164-166.
3. ———, *Nodal noncommutative Jordan algebras*, Canadian J. Math. (to appear).
4. R. D. Schafer, *On noncommutative Jordan algebras*, Proc. Amer. Math. Soc. vol. 9 (1958) pp. 110-117.

WASHINGTON UNIVERSITY