

# ORDER AND COMMUTATIVITY IN BANACH ALGEBRAS

PHILIP C. CURTIS, JR.

S. Sherman has shown [4] that if the self adjoint elements of a  $C^*$  algebra form a lattice under their natural ordering the algebra is necessarily commutative. In this note we extend this result to real Banach algebras with an identity and arbitrary Banach  $*$  algebras with an identity. The central fact for a real Banach algebra  $A$  is that if the positive cone is defined to be the uniform closure of the set of finite sums of squares of elements of  $A$ , and if  $A$  is a lattice under the ordering induced by this cone, then extreme points of the unit sphere of the dual cone are multiplicative linear functionals. A similar situation holds for  $*$  algebras.

**1. Real Banach algebras.** Let  $X$  be a real Banach space, and let  $C$  be a closed cone in  $X$ .<sup>1</sup> For  $x, y \in X$  we define  $x \geq y$  if  $x - y \in C$ . If in addition  $X$  is a lattice under the ordering  $\geq$ , we say  $C$  lattice-orders  $X$ . Let  $C'$  be the dual cone and let  $\Sigma = \{f \in C' : \|f\| \leq 1\}$ . The set of extreme points of  $\Sigma$  will be denoted by  $S$ . For a real linear functional  $f$  let  $I_f = \{x \in X : f(x) = 0\}$  and let  $R = \bigcap_{f \in C'} I_f$ . Lastly if  $X$  is a lattice we define  $x_+ = x \vee 0$ ,  $x_- = x \wedge 0$ , and  $|x| = x_+ - x_-$ . We note  $|x| \geq 0$ .

LEMMA 1. *If  $C$  is a closed cone in a real Banach space  $X$ , then*

- (i)  $R = C \cap -C$ ,
- (ii)  $R = \bigcap_{f \in S} I_f$ ,
- (iii) *If  $C$  lattice-orders  $X$ ,  $R = \{0\}$ .*

PROOF. Obviously  $C \cap -C \subset R$ . For the converse, by the Hahn-Banach theorem  $x \in C$  iff  $f(x) \geq 0$  for each  $f \in C'$ . Therefore  $R \subset C \cap -C$ . For (ii) suppose  $x \in \bigcap_{f \in S} I_f$ ,  $x \notin R$ , then there exists an  $f \in C'$ ,  $\|f\| = 1$ , such that  $|f(x)| = 2\epsilon \neq 0$ . But by the Krein-Milman theorem there exist finitely many  $f_i \in S$  and real numbers  $\alpha_i$  such that  $|f(x) - \sum \alpha_i f_i(x)| < \epsilon$ . Hence for some  $i$ ,  $f_i(x) \neq 0$ , which is a contradiction. Lastly let  $C$  lattice order  $X$ . Then  $x \in C$  implies  $x \geq 0$  or  $x_- = 0$ , and  $x \in -C$  implies  $-x \geq 0$  or  $x_+ = 0$ . Since  $x = x_+ + x_-$ ,  $x \in C \cap -C$  implies  $x = 0$ .

The central tool in both this investigation and that of Sherman is the following result of Krein and Krein [3]. It can be stated in a

---

Received by the editors August 5, 1957.

<sup>1</sup> We refer the reader to [2] for the appropriate definitions of cone, dual cone, etc.

slightly more general fashion, but the following is sufficient for our purposes.

**THEOREM 1.** *Let  $X$  be a real Banach space which is lattice ordered by a cone  $C$ . Suppose in addition that  $C$  contains an element  $e$ ,  $\|e\|=1$ , such that  $\{y: \|e-y\|\leq 1\}\subset C$ . Then  $f\in S$  iff  $|f(x)|=f(|x|)$  for each  $x\in X$ .*

Let us specialize to a real Banach algebra  $A$  with identity such that  $\|1\|=1$ , and let  $C$  be the closure of the set of finite sums of squares of elements of  $A$ . By the familiar binomial series argument (c.f. [2]),  $\{y: \|1-y\|\leq 1\}\subset C$ . Also, for  $f\in C'$  and  $x, y\in A$  we have the Schwartz inequality,  $[f(xy+yx)]^2\leq 4f(x^2)f(y^2)$ , which may be verified by the classical argument. Also useful is the following property of functionals in  $C'$ .

**LEMMA 2.** *Let  $A$  be a real Banach algebra, and let  $x\in C, f\in C'$ . Then  $f(x)=0$  implies  $f(x^2)=0$ .*

**PROOF.** First assume  $x=y^2$ , and let  $f\in C'$ . If  $\|x\|\leq 1$ , the binomial series for  $(1-x)^{1/2}$  converges absolutely. Therefore  $1-x\in C$ . Moreover since  $x=y^2$ ,  $y(1-x)^{1/2}=(1-x)^{1/2}y$  and  $x(1-x)=[y(1-x)^{1/2}]^2\in C$ . Therefore  $x-x^2\geq 0$ . Hence  $f(x)=0$  implies  $f(x^2)=0$ . We proceed now by induction. Let  $x=\sum_{i=1}^{n+1} y_i^2$  and let  $f(x)=0$ . If  $z=\sum_{i=1}^n y_i^2$ , then  $f(y_{n+1}^2)=f(z)=0$ . We assume  $f(z^2)=0$ , and by the above argument  $f(y_{n+1}^4)=0$ . An application of the Schwartz inequality gives us

$$0\leq f(x^2)=f((z+y_{n+1}^2)^2)=f(zy_{n+1}^2+y_{n+1}^2z)\leq 2[f(z^2)f(y_{n+1}^4)]^{1/2}=0.$$

Therefore the result holds for all finite sums of squares, and by continuity it holds for all  $x$  in  $C$ .

**THEOREM 2.** *If  $C$  lattice-order  $A$  then each  $f\in S$  is a homomorphism of  $A$  onto the real numbers, and  $A$  is commutative.*

**PROOF.** For  $x, y\in A$  define the Jordan multiplication  $x\circ y=(xy+yx)/2$ . Thus  $A$  can be considered as a Jordan ring with an identity. We assert that for  $f\in S$ ,  $I_f$  is a Jordan ideal. By Theorem 1  $x\in I_f$  iff  $x_+, x_-\in I_f$ . Therefore let  $x\geq 0, x\in I_f$ . By the Schwartz inequality and Lemma 2, for any  $y\in A$ ,  $[f(xy+yx)]^2\leq 4f(x^2)f(y^2)=0$ . Hence  $xy+yx\in I_f$ , and since  $I_f$  is obviously closed under addition,  $I_f$  is a Jordan ideal.

Now a linear functional of any algebra over a field which takes the identity of the algebra into the identity of the field is a homomorphism if its kernel is a two-sided ideal. Hence  $f$  is a Jordan homo-

morphism of  $A$  onto the reals. On the other hand Jacobson and Rickart [1, Theorem 2] have proved that a Jordan homomorphism of a ring into an integral domain is either a homomorphism or an anti-homomorphism. An application of this result proves that  $f$  is a homomorphism.

Finally for each  $f \in S$  and  $x, y \in A$ ,  $xy - yx \in I_f$ . Since by Lemma 1  $\bigcap_{f \in S} I_f = \{0\}$ ,  $A$  must be commutative.

**2. Banach  $*$  algebras.** Let  $A$  be a Banach  $*$  algebra with a continuous involution and an identity. Let  $C$  be the closure of the set of finite sums of elements  $xx^*$ .  $C$  is a closed cone in the real linear space  $H$  of self adjoint elements of  $A$ . The dual cone of  $C$  (in the conjugate space of  $H$ ) can be identified with the set of those functionals  $f$  on  $A$  for which  $f(xx^*) \geq 0$  for each  $x \in A$  (c.f. [2] for details). Let  $\sum, S$  be as before and for  $f \in C'$  let  $I_f = \{x \in A : f(x) = 0\}$  and  $R = \bigcap_{f \in C'} I_f$ . We also note that  $\{h \in H : \|1 - h\| \leq 1\} \subset C$  and for  $f \in C'$  the familiar Schwartz inequality holds, i.e.  $|f(xy^*)|^2 \leq f(xx^*)f(yy^*)$ ,  $x, y \in A$ .

LEMMA 3.

$$R = C \cap -C + i(C \cap -C),$$

$$R = \bigcap_{f \in S} I_f.$$

PROOF. Let  $T = (C \cap -C) + i(C \cap -C)$ . Obviously  $T \subset R$ . If  $x \in R$ , let  $h = (x + x^*)/2$ ,  $k = (x - x^*)/2i$ . Then  $h, k$  are self adjoint,  $h, k \in I_f$  and  $x = h + ik$ . But a self adjoint element  $y \in C$  iff  $f(y) \geq 0$  for each  $f \in C'$ . Therefore  $h, k \in C \cap -C$  and  $T = R$ . For the second assertion if  $x \in \bigcap_{f \in S} I_f$  and  $x \notin R$ , we may assume  $x$  is self adjoint and apply the argument of Lemma 1.

LEMMA 4. Let  $h \in C$  and  $f \in C'$ . Then  $f(h) = 0$  implies  $f(h^2) = 0$ .

PROOF. If  $h = h^*$ , and  $\|h\| \leq 1$ , then by the familiar series argument  $1 - h = k^2$ , where  $k = k^*$ . Therefore  $1 - h \in C$ . Since  $kh = hk$  and  $khk \in C$ ,  $khk = hk^2 = h - h^2 \in C$ . Therefore  $f(h) = 0$  implies  $f(h^2) = 0$ .

THEOREM 3. If  $C$  lattice-orders  $H$ , then each  $f \in S$  is a homomorphism of  $A$  onto the complex numbers, and  $A$  is commutative.

PROOF. To prove that  $f$  is a homomorphism it suffices to show that for  $f \in S$ ,  $I_f$  is a two-sided ideal. Let  $x \in I_f$ ,  $y \in A$ . We assert  $xy \in I_f$ . First we may assume  $x$  is self adjoint and by Theorem 1 we may assume  $x \geq 0$ . But then applying the Schwartz inequality

$$|f(xy)|^2 \leq f(x^2)f(y^*y) = 0.$$

This proves  $xy \in A$  and similarly  $yx \in A$ . Since  $I_f$  is obviously closed under addition, it is a two sided ideal. An application of Lemma 3 proves that  $A$  is commutative.

### BIBLIOGRAPHY

1. N. Jacobson and C. E. Rickart, *Jordan homomorphism of rings*, Trans. Amer. Math. Soc. vol. 69 (1950) pp. 479–502.
2. J. L. Kelley and R. L. Vaught, *The positive cone in Banach algebras*, Trans. Amer. Math. Soc. vol. 74 (1953) pp. 44–55.
3. M. Krein and S. Krein, *On an inner characteristic of the set of all continuous functions defined on a bicomact Hausdorff space*, Comptes Rendus (Doklady) de l'Academie des Sciences de l'URSS. N. S. vol. 27 (1940) pp. 427–430.
4. S. Sherman, *Order in operator algebras*, Amer. J. Math. vol. 73 (1951) pp. 227–232.

UNIVERSITY OF CALIFORNIA AT LOS ANGELES

---

## A NOTE ON VALUED LINEAR SPACES

PAUL CONRAD

Banaschewski [1] has given a simple and elegant proof of Hahn's embedding theorem for ordered abelian groups. His method can be used to prove the author's generalization of Hahn's theorem [2, p. 11]. In this note we make use of Banaschewski's method to prove a special case of the author's theorem (which is also a generalization of Hahn's theorem) that has been proven by Gravett [3].

Let  $(L, \Delta, d)$  be a valued linear space [3]. That is,  $L$  is a vector space over a division ring  $K$ ,  $\Delta$  is a linearly ordered set with minimum element  $\theta$ , and  $d$  is a mapping of  $L$  onto  $\Delta$  such that for all  $x, y \in L$ ,  $d(x) = \theta$  if and only if  $x = 0$ ,  $d(x) = d(kx)$  for all  $0 \neq k \in K$ , and  $d(x+y) \leq \text{Max } [d(x), d(y)]$ . For each  $\delta \in \Delta$ , let  $C^\delta = \{x \in L: d(x) \leq \delta\}$  and let  $C_\delta = \{x \in L: d(x) < \delta\}$ . Let  $W$  be the vector space of all mappings  $f$  of  $\Delta$  into the join of the spaces  $C^\delta/C_\delta$  for which  $f(\delta) \in C^\delta/C_\delta$  and  $R_f = \{\delta \in \Delta: f(\delta) \neq C_\delta\}$  is an inversely well ordered set.  $W$  is a subspace of the unrestricted direct sum  $V$  of the  $C^\delta/C_\delta$ .  $W$  is also a valued linear space  $(W, \Delta, d')$ , with  $d'(f)$  the largest  $\delta \in R(f)$ .