ON A CLASS OF UNIVALENT, STAR SHAPED MAPPINGS

ALBERT SCHILD¹

1. Introduction. Among all functions $w=f(z)=z+\sum_{n=2}^{\infty} a_n z^n$ regular and univalent in the unit circle, two classes of functions have been discussed extensively: The class of functions mapping the unit circle onto star shaped regions, characterized by $\operatorname{Re}\{zf'(z)/f(z)\} \geq 0$ for $|z| < 1$, and the class of functions mapping the unit circle onto convex regions, characterized by $\operatorname{Re}\{zf''(z)/f'(z)\} + 1 \geq 0$ for $|z| < 1$.

This short paper will examine some of the geometric and analytic properties of a class of functions $w=f(z)=z+\sum_{n=2}^{\infty} a_n z^n$ which map the unit circle onto a region whose geometric nature is somewhat intermediate between star shaped and convex. The functions under consideration are to satisfy $\operatorname{Re}\{zf'(z)/f(z)\} \geq 1/2$ for all $|z| < 1$.

Interest in functions of this type can be traced back to two papers by A. Marx [4] and E. Strohäcker [8] who have shown that for any function $w=f(z)=z+\sum_{n=2}^{\infty} a_n z^n$, which maps the unit circle onto a convex region, we have $\operatorname{Re}\{zf'(z)/f(z)\} \geq 1/2$, and as the function $f(z)=z/(1+z)$ shows, the constant $1/2$ cannot be improved. It is also clear that the converse is not true, i.e. functions for which $\operatorname{Re}\{zf'(z)/f(z)\} \geq 1/2$ need not map the unit circle onto a convex region. An example of this type is given by the function $w=f(z)=z-1/3 z^2$ for which $\operatorname{Re}\{zf'(z)/f(z)\} \geq 1/2$, $|z| \leq 1$, yet the image region is not convex.

Recently, interest in this class of functions was roused again in a paper by R. F. Gabriel [2]. There it is shown that if $p(z)$ is analytic and single valued for $|z| < 1$, and if we denote by $w_1(z)=1+\sum_{n=2}^{\infty} a_n z^n$ and by $w_2(z)=z+\sum_{n=2}^{\infty} b_n z^n$ two linearly independent solutions of $w''+p(z)w=0$, then $f(z)=w_1(z)/w_2(z)=1/z+\dots$ will map $|z| \leq 1$ onto the exterior of a convex region if and only if $\operatorname{Re}\{zf'(z)/f(z)\} \geq 1/2$ for $|z| < 1$.

M. S. Robertson [6] considered also the class of functions $w=f(z)=z+\sum_{n=2}^{\infty} a_n z^n$ for which $\operatorname{Re}\{zf'(z)/f(z)\} \geq \alpha > 0$, $|z| < 1$.

NOTATION. On the following pages we shall denote the class of functions $w=f(z)=z+\sum_{n=2}^{\infty} a_n z^n$ regular and univalent in the unit circle by S .

The subclass of S which map $|z| < 1$ onto a star shaped region, i.e. for which $\operatorname{Re}\{zf'(z)/f(z)\} \geq 0$ by St.

Presented to the Society, January 28, 1958; received by the editors March 7, 1958 and, in revised form, April 21, 1958.

¹ This work was aided by a grant from the Committee on Research and Publications of Temple University.

The subclass of St which map $|z| < 1$ onto a "special" star shaped region, i.e. for which $\operatorname{Re}\{zf'(z)/f(z)\} \geq 1/2$ by St^* .

The subclass of S which map $|z| < 1$ onto a convex region, i.e. for which $\operatorname{Re}\{zf''(z)/f'(z)\} + 1 \geq 0$ by C .

By [4] and [8] we obviously have:

$$1.1 \quad S \supset St \supset St^* \supset C.$$

2. Methods for constructing functions of class St^* . The intermediacy of St^* between St and C suggests, that one might obtain functions of class St^* by "relaxing" the conditions on functions of class C or by "strengthening" conditions on functions of class St . Both methods yield results.

THEOREM 2.1. *If $w=f(z) \in C$, then $g(z)=z(f'(z))^\alpha \in St^*$ for $0 < \alpha \leq 1/2$. The converse is true for $\alpha=1/2$.*

PROOF. $\operatorname{Re}\{zg'(z)/g(z)\} = \operatorname{Re}\{1 + \alpha zf''(z)/f'(z)\} \geq 1 - \alpha \geq 1/2$. For the converse we have: $\operatorname{Re}\{zf''(z)/f'(z)\} + 1 = \operatorname{Re}\{2zg'(z)/g(z)\} - 1 \geq 0$. This should be compared with the well known result that if $f(z) \in C$ then $g(z)=zf'(z) \in St$ and conversely.

COROLLARY 2.2. *Let $|b_i|=1$ and $\sum_{i=1}^n \mu_i \leq 1$, $\mu_i \geq 0$, then $w=g(z) = z \prod_{i=1}^n (1 - b_i z)^{-\mu_i} \in St^*$.*

PROOF. $f(z) = \int_0^z \prod_{i=1}^n (1 - b_i t)^{-v_i} dt \in C$ if $v_i \geq 0$ and $\sum_{i=1}^n v_i \leq 2$, $|b_i|=1$. The corollary follows if we let $\alpha=1/2$ in Theorem 2.1.

THEOREM 2.3. *If $w=f(z) \in St$, then $g(z)=z\{f(z)/z\}^\alpha \in St^*$ for $0 < \alpha \leq 1/2$. The converse is true for $\alpha=1/2$.*

PROOF. $\operatorname{Re}\{zg'(z)/g(z)\} = \operatorname{Re}\{\alpha zf'(z)/f(z) + 1 - \alpha\} \geq 1 - \alpha \geq 1/2$.

For the converse we have: $\operatorname{Re}\{zf'(z)/f(z)\} = \operatorname{Re}\{2zg'(z)/g(z) - 1\} \geq 0$. (Marx [4] has the above theorem for $\alpha=1/2$.)

The method of Theorem 2.3 for constructing functions of class St^* gives some insight into the geometrical nature of the region onto which the unit circle is mapped by functions of class St^* . If $z=re^{i\theta}$ is mapped by $w=f(z) \in St$ into $Re^{i\phi}$, then a comparison of r and R and θ and ϕ tells us about the amount of distortion. We notice that $g(z)=\{zf(z)\}^{1/2} \in St^*$ reduces the amount of distortion effected by $f(z) \in St$ by an "averaging" process, i.e. $z=re^{i\theta}$ will be mapped by $g(z)$ into $\rho e^{i\psi}$ where $\rho=(rR)^{1/2}$ and $\psi=(\theta+\phi)/2$.

3. Distortion theorems for functions of class St^* .

THEOREM 3.1. *For all $g(z) \in St^*$ we have $|z|/(1+|z|) \leq |g(z)| \leq |z|/(1-|z|)$.*

PROOF. By Theorem 2.3 ($\alpha=1/2$) and the "Verzerrungs Satz" we have: $|z|/(1+|z|)^2 \leq |g^2(z)/z| \leq |z|/(1-|z|)^2$ and hence the Theorem follows. These inequalities are sharp for $g(z)=z/(1+z) \in \text{St}^*$, $z = \pm r$.

THEOREM 3.2. For all $g(z) \in \text{St}^*$ the domain of values of $zg'(z)/g(z)$ is the circle with center at $1/(1-|z|^2)$ and radius $|z|/(1-|z|^2)$.

PROOF. Let

$$G(z) = zg'(z)/g(z) - 1/2$$

and

$$H(z) = (2G(z) - 1)/(2G(z) + 1) = \{zg'(z)/g(z) - 1\}/zg'(z)/g(z).$$

Then $H(z)$ is regular for $|z| < 1$, $H(0)=0$ and $|H(z)| < 1$ for $|z| < 1$. Hence the Lemma of Schwarz can be applied and we have for $|z| < 1$

$$\left| \left(\frac{zg'(z)}{g(z)} - 1 \right) / \frac{zg'(z)}{g(z)} \right| < |z| \quad \text{or:} \quad \left| \frac{zg'(z)}{g(z)} - 1 \right| < |z| \left| \frac{zg'(z)}{g(z)} \right|.$$

But the domain defined by this inequality is the interior of the Circle of Apollonius with the line segment from $1/(1+|z|)$ to $1/(1-|z|)$ as a diameter, i.e. the interior of the circle with radius $|z|/(1-|z|^2)$ and center at $1/(1-|z|^2)$. The function $f(z)=z/(1+z) \in \text{St}^*$ shows that the theorem cannot be improved.

4. **Some coefficient relations.** It is well known that if $f(z) \in \text{St}$ then $|a_n| \leq n$. For functions of class St^* we have:

THEOREM 4.1. If $f(z)=z+\sum_{n=2}^{\infty} a_n z^n \in \text{St}^*$, then $|a_n| \leq 1$.

PROOF. Let $p(z)=2zf'(z)/f(z)-1=1+c_1z+c_2z^2+\dots$. Since $p(z)$ is regular and $\text{Re}\{p(z)\} > 0$ for $|z| < 1$, therefore, by a well known lemma we have $|c_n| \leq 2$ for $n=1, 2, 3, \dots$.

Comparing coefficients we obtain $2(n-1)a_n=c_{n-1}+a_2c_{n-2}+\dots+a_{n-1}c_1$, and hence: $|a_n| \leq 1/(n-1)\{1+|a_2|+\dots+|a_{n-1}|\}$. It follows now by induction that $|a_n| \leq 1$ for $n=1, 2, 3, \dots$.

The function $f(z)=z/(1-z)=\sum_{n=1}^{\infty} z^n \in \text{St}^*$ shows that these inequalities are sharp.

For some kind of a converse we have:

THEOREM 4.2. If $g(z)=z+\sum_{n=2}^{\infty} a_n z^n$ and if $\sum_{n=2}^{\infty} (2n-1)|a_n| \leq 1$ then $g(z) \in \text{St}^*$.

PROOF. The proof is based on a method used by A. Goodman [3]. We have:

$$\begin{aligned}
 (4.3) \quad \frac{zg'(z)}{g(z)} - \frac{1}{2} &= \frac{1 + 3a_2z + 5a_3z^2 + \cdots + (2n-1)a_nz^{n-1} + \cdots}{2(1 + a_2z + a_3z^2 + \cdots + a_nz^{n-1} + \cdots)} \\
 &= \frac{1}{2} + \sum_{n=1}^{\infty} b_n z^n
 \end{aligned}$$

where

$$\begin{aligned}
 (4.4) \quad b_1 &= a_2, \\
 b_2 &= 2a_3 - a_2b_1, \\
 b_3 &= 3a_4 - a_2b_2 - a_3b_1, \\
 &\vdots \\
 b_{n-1} &= (n-1)a_n - a_2b_{n-2} - a_3b_{n-3} - \cdots - a_{n-1}b_1, \\
 &\vdots
 \end{aligned}$$

and therefore for $n \geq 2$

$$(4.5) \quad \sum_{k=1}^{n-1} b_k = \sum_{k=2}^n (k-1)a_k - a_2 \sum_{k=1}^{n-2} b_k - a_3 \sum_{k=1}^{n-3} b_k - \cdots - a_{n-1}b_1.$$

The inequality of the theorem implies that $|b_1| = |a_2| \leq 1/3$. It will now be shown by mathematical induction that for all n we have $|\sum_{k=1}^n b_k| \leq 1/2$. Assume that $|\sum_{k=1}^m b_k| \leq 1/2$ for $m=1, 2, 3, \dots, n-2$. Then (4.5) yields

$$\begin{aligned}
 (4.6) \quad \left| \sum_{k=1}^{n-1} b_k \right| &\leq \sum_{k=2}^n (k-1) |a_k| + \frac{1}{2} \sum_{k=2}^{n-1} |a_k| \leq \sum_{k=2}^n (k-1/2) |a_k| \\
 &= \frac{1}{2} \sum_{k=2}^n (2k-1) |a_k| \leq 1/2
 \end{aligned}$$

by the inequality of the theorem, and hence $|\sum_{k=1}^n b_k| \leq 1/2$ for all n . From (4.3) it follows now that

$$(4.7) \quad |zg'(z)/g(z) - 1| \leq 1/2 \text{ for } z = r.$$

But the inequality of Theorem 4.2 and the special starshapedness of the image domain are invariant under rotations of the z and w planes. Hence any point in the unit circle may be placed in the interval $[0, 1]$, and thus (4.7) is valid throughout the unit circle i.e.

$$\operatorname{Re} \{zg'(z)/g(z)\} \geq 1/2 \text{ for } |z| < 1.$$

COROLLARY 4.8. *Let $g(z) = z - \sum_{n=2}^{\infty} a_n z^n$, where all $a_i \geq 0$, then $g(z) \in \text{St}^*$, if and only if $\sum_{n=2}^{\infty} (2n-1)a_n \leq 1$.*

PROOF. The sufficiency of the condition follows from the Theorem. For the necessity we have:

$$(4.9) \quad \operatorname{Re} \left\{ \frac{zg'(z)}{g(z)} - \frac{1}{2} \right\} = \operatorname{Re} \left\{ \frac{1 - \sum_{n=2}^{\infty} (2n-1)a_n z^{n-1}}{2 \left(1 - \sum_{n=2}^{\infty} a_n z^{n-1} \right)} \right\}.$$

If $\sum_{n=2}^{\infty} (2n-1)a_n > 1$, then we could find a positive value of z , r_0 , for which the numerator of (4.9) would be negative and the denominator positive and hence $\operatorname{Re}\{zg'(z)/g(z)\} < 1/2$ for that value of z and all positive values of $z > r_0$. (See also [7].)

5. The radius of special star shapedness. We define the radius of special star shapedness, r^* , as the upper bound of the radii, r , of circles $|z| \leq r$, which are mapped by any function $f(z) \in S$ onto a region of special star shapedness, i.e. that for all functions $f(z) \in S$ we have $\operatorname{Re}\{zf'(z)/f(z)\} \geq 1/2$ for all $|z| \leq r$. From the introduction it follows that r^* is larger than the bound of convexity (Rundungsschranke) and less than the bound of starlikeness. Therefore

$$(5.1) \quad 2 - 3^{1/2} = .268 \dots \leq r^* \leq .65 \dots = \tanh \pi/4.$$

We have:

THEOREM 5.2. $r^* = 1/3$ for all $f(z) \in \operatorname{St}$, and hence $r^* \leq 1/3$.

PROOF. By [5] we have for any $f(z) \in \operatorname{St}$

$$\frac{1 - |z|}{1 + |z|} \leq \operatorname{Re} \left\{ \frac{zf'(z)}{f(z)} \right\} \leq \frac{1 + |z|}{1 - |z|}.$$

Therefore $\operatorname{Re}\{zf'(z)/f(z)\} \geq (1 - |z|)/(1 + |z|) \geq 1/2$ for all $|z| \leq 1/3$. This result is sharp for $f(z) = z(1+z)^{-2} \in \operatorname{St}$ and $z = 1/3$.

THEOREM 5.3. A lower bound for r^* is: $r^* > .301 \dots$.

PROOF. If we let

$$(5.4) \quad \begin{aligned} g(z) &= f^{-1/2}(z^{-2}) \\ &= z + b_1/z + b_3/z^3 + \dots + b_{2n-1}/z^{2n-1} + \dots \end{aligned}$$

then by Bieberbach's Flächensatz [1] we have

$$(5.5) \quad \sum_{n=1}^{\infty} (2n-1) |b_{2n-1}|^2 \leq 1.$$

The condition $\operatorname{Re}\{zf'(z)/f(z)\} \geq 1/2$ is equivalent to

$$(5.6) \quad \left| \frac{zf'(z)}{f(z)} \right| \geq \left| \frac{zf'(z)}{f(z)} - 1 \right|$$

i.e. by (5.4)

$$\left| \frac{z^{-1/2}g'(z^{-1/2})}{g(z^{-1/2})} \right| \geq \left| \frac{z^{-1/2}g'(z^{-1/2})}{g(z^{-1/2})} - 1 \right|$$

or:

$$(5.7) \quad \left| \frac{1 - b_1z - 3b_3z^2 - \dots - (2n-1)b_{2n-1}z^n - \dots}{1 + b_1z + b_3z^2 + \dots + b_{2n-1}z^n + \dots} \right| \geq \left| \frac{1 - b_1z - 3b_3z^2 - \dots - (2n-1)b_{2n-1}z^n + \dots}{1 + b_1z + b_3z^2 + \dots + b_{2n-1}z^n + \dots} - 1 \right|$$

or:

$$\begin{aligned} & |1 - b_1z - 3b_3z^2 - \dots - (2n-1)b_{2n-1}z^n| \\ & \geq 2|b_1z + 2b_3z^2 + 3b_5z^3 + \dots + nb_{2n-1}z^n + \dots|. \end{aligned}$$

A sufficient condition for this inequality to be satisfied is that

$$\begin{aligned} 1 - |b_1|r - 3|b_3|r^2 - \dots - (2n-1)|b_{2n-1}|r^n + \dots \\ \geq 2|b_1|r + 4|b_3|r^2 + \dots \end{aligned}$$

or:

$$\begin{aligned} 3|b_1|r + 7|b_3|r^2 + 11|b_5|r^3 + \dots \\ + (4n-1)|b_{2n-1}|r^n + \dots \leq 1 \end{aligned}$$

or

$$(5.8) \quad 2\{|b_1|r + 3|b_3|r^2 + \dots + (2n-1)|b_{2n-1}|r^n + \dots\} + \{|b_1|r + |b_3|r^2 + \dots + |b_{2n-1}|r^n + \dots\} \leq 1.$$

If in the first parenthesis of inequality (5.8) we let $c_n = (2n-1)^{1/2}|b_{2n-1}|$ and $d_n = (2n-1)^{1/2} \cdot r^n$ then the Inequality of Schwarz

$$\sum c_n \cdot d_n \leq (\sum c_n^2)^{1/2} \cdot (\sum d_n^2)^{1/2}$$

gives

$$(5.9) \quad \begin{aligned} & |b_1|r + 3|b_3|r^2 + \dots + (2n-1)|b_{2n-1}|r^n + \dots \\ & \leq (|b_1|^2 + 3|b_3|^2 + \dots + (2n-1)|b_{2n-1}|^2 + \dots)^{1/2} \\ & \quad \cdot (r^2 + \dots + (2n-1)r^{2n} + \dots)^{1/2} \end{aligned}$$

and therefore by (5.5)

$$(5.10) \quad \begin{aligned} & |b_1| r + 3 |b_3| r^2 + \cdots + (2n-1) |b_{2n-1}| r^n + \cdots \\ & \leq (r^2 + 3r^4 + \cdots + (2n-1)r^{2n} + \cdots)^{1/2} = \frac{r(1+r^2)^{1/2}}{1-r^2}. \end{aligned}$$

Similarly, if in the second parenthesis of inequality (5.8) we let $c_n = (2n-1)^{1/2} |b_{2n-1}|$, $d_n = r^n(2n-1)^{-1/2}$ and apply the Inequality of Schwarz again, we get:

$$(5.11) \quad \begin{aligned} & |b_1| r + |b_3| r^2 + \cdots + |b_{2n-1}| r^n + \cdots \\ & \leq (|b_1|^2 + 3 |b_3|^2 + \cdots + (2n-1) |b_{2n-1}|^2 + \cdots)^{1/2} \\ & \quad \cdot \left(r^2 + \frac{r^4}{3} + \cdots + \frac{r^{2n}}{2n-1} + \cdots \right)^{1/2} \end{aligned}$$

and again by (5.5)

$$(5.12) \quad \begin{aligned} & |b_1| r + |b_3| r^2 + \cdots + |b_{2n-1}| r^n + \cdots \\ & \leq \left(r^2 + \frac{r^4}{3} + \cdots + \frac{r^{2n}}{2n-1} + \cdots \right)^{1/2} = \left(\frac{r}{2} \ln \frac{1+r}{1-r} \right)^{1/2}. \end{aligned}$$

Substituting (5.10) and (5.12) into (5.8) we have:

$$(5.13) \quad \frac{2r(1+r^2)^{1/2}}{1-r^2} + \left(\frac{r}{2} \log \frac{1+r}{1-r} \right)^{1/2} < 1.$$

This will be satisfied for $|z| = r < .301 \cdots$, i.e. $r^* > .301 \cdots$.

BIBLIOGRAPHY

1. L. Bieberbach, *Lehrbuch der Funktionentheorie*, vol. 2, pp. 84-85.
2. R. F. Gabriel, *The Schwarzian derivative and convex functions*, Proc. Amer. Math. Soc. vol. 6 (1955) pp. 58-66.
3. A. Goodman, *Univalent functions and nonanalytic curves*, Proc. Amer. Math. Soc. vol. 8 (1957) pp. 598-601.
4. A. Marx, *Untersuchungen über schlichte Abbildungen*, Math. Ann. vol. 107 (1932-1933) pp. 40-67.
5. Pólya and Szegő, *Aufgaben und Lehrsätze*, vol. 1, problem 287, p. 140.
6. M. S. Robertson, *On the theory of univalent functions*, Ann. of Math. vol. 37 (1936) pp. 374-408.
7. A. Schild, *On a class of functions schlicht in the unit circle*, Proc. Amer. Math. Soc. vol. 5 (1954) pp. 115-120.
8. E. Strohäcker, *Beiträge zur Theorie der schlichten Funktionen*, Math. Z. vol. 37 (1933) pp. 356-380.

TEMPLE UNIVERSITY