COEFFICIENTS IN CERTAIN ASYMPTOTIC FACTORIAL EXPANSIONS

T. D. RINEY

1. Introduction. Let p and q denote integers with $0 \leq p \leq q$, then it is known from [2; 3] and [7] that the function

$$(1.1) \quad g(w) = \prod_{i=1}^p \Gamma(w + \sigma_i) / \prod_{j=0}^q \Gamma(w + \rho_j),$$

where the σ_i and ρ_j are independent of w , admits for large $|w|$ in the sector $|\arg w| < \pi - \epsilon$, an asymptotic expansion of the form

$$(1.2) \quad g(w) = ((2\pi)^{1/2})^{1-\alpha} \alpha^{\alpha w + \beta - 1/2} \left[\sum_{m=0}^M c_m / \Gamma(\alpha w + \beta + m) + O(1/\Gamma(\alpha w + \beta + M + 1)) \right].$$

Here, and in the sequel, M denotes a non-negative integer independent of w , ϵ any positive constant and

$$(1.3) \quad c_0 = 1; \quad \alpha = q + 1 - p; \quad \beta = \sum_{j=0}^q \rho_j - \sum_{i=1}^p \sigma_i + (1 - \alpha)/2.$$

Recently two types of linear recurrence relations satisfied by the c_m have been found. The first [5] is an inductive formula for c_m depending on all the preceding c_n ; the second [4] gives c_m in terms of the preceding q coefficients. An alternative method of obtaining a recursion formula of the latter type has been pointed out by E. M. Wright [6].

J. G. van der Corput [8] has shown that the inverse of $g(w)$ possesses the asymptotic development

$$(1.4) \quad (g(w))^{-1} = ((2\pi)^{1/2})^{\alpha-1} \alpha^{-\alpha w - \beta + 1/2} \left[\sum_{m=0}^M \gamma_m \Gamma(\alpha w + \beta - m) + O(\Gamma(\alpha w + \beta - M - 1)) \right],$$

in the sector $|\arg w| < \pi - \epsilon$. Moreover, van der Corput has extended

Presented to the Society, August 27, 1958, under the title of *Coefficients in certain asymptotic factorial expansions of the second kind*; received by the editors August 7, 1958 and, in revised form, November 6, 1958.

the method of [5] to obtain a linear recursion formula for γ_m , $\gamma_0 = 1$, depending on all the preceding γ_n . The purpose here is to present a general method which when applied to $g(w)$ and $(g(w))^{-1}$ yields a finite recursion formula of fixed length q for the c_m and γ_m respectively.¹

2. General factorial expansion. Let us consider the function

$$(2.1) \quad F(x) = \prod_{j=0}^q \Gamma(x + s_j) / \Gamma(x + r_j),$$

where the s_j and r_j are independent of x . From Stirling's formula it is easy to show that there exist constants A_m , $A_0 = 1$, such that in the sector $|\arg x| \leq \pi - \epsilon$ $F(x) = \sum_{m=0}^M A_m x^{-b-m} + O(x^{-b-M-1})$, where

$$(2.2) \quad b = \sum_{j=0}^q (r_j - s_j).$$

The proof which is similar to the proofs cited for (1.2) is omitted. We immediately have the following result which will be stated in the form of a theorem.

THEOREM 2.1. *Let d_m and a be given complex numbers, $|\arg a| < \pi/2$. Then there exist constants B_m , $B_0 = 1$, such that*

$$(2.3) \quad F(x/a) = a^b \sum_{m=0}^M B_m \Gamma(x + d_m - m) / \Gamma(x + d_m + b) + O(x^{-b-M-1})$$

in the sector $|\arg x| < \pi/2 + \epsilon$ provided $\epsilon < \pi/2 - |\arg a|$.

The problem is to determine an efficient method for computing the coefficients B_m . For this purpose we shall need the general result of the next section.

3. A general theorem. Let $G(x)$ denote a function analytic in some sector $|\arg(x - x_0)| \leq \pi/2 + \epsilon$, ($\epsilon < \pi/2$), such that in this sector

$$(3.1) \quad G(x) = O(x^{-h}),$$

where h denotes a constant. For such a function, we define

$$(3.2) \quad \psi(t) = L\{G\} = (2\pi i)^{-1} \int_C t^{\lambda x} G(x) dx, \quad \begin{aligned} (1 \leq t < \infty \text{ if } \lambda > 0), \\ (0 < t \leq 1 \text{ if } \lambda < 0), \end{aligned}$$

where λ ($\neq 0$) is real and C denotes the path $|\arg(x - x_0)| = \pi/2 + \epsilon$ orientated in the upward direction; clearly, $\psi(t)$ does not depend on

¹ We had originally demonstrated the result for γ_m . This method of giving the general result was suggested by the referee.

the choice of ϵ or x_0 . Further, $\psi(t)$ is infinitely differentiable, in fact

$$(3.3) \quad \psi^{(k)}(t) = \left(\frac{d}{dt}\right)^k L\{G\} = t^{-k} L\{\lambda x(\lambda x - 1) \cdots (\lambda x - k + 1)G(x)\}.$$

Finally, we note the following result:

THEOREM 3.1. $G(x) = O(x^{-h})$ implies that near $t = 1$

$$(3.4) \quad \psi^{(k)}(t) = O(|t - 1|^{h-k-1}),$$

provided $k < \text{Re}(h) - 1$.

PROOF. In the integral representation of $\psi^{(k)}(t)$ replace the curve C by $\text{Re}(x) = u \geq u_0 = \text{Re}(x_0)$ and set $x = u + iv$ to obtain

$$\begin{aligned} \psi^{(k)}(t) = (2\pi)^{-1} t^{\lambda u - k} \int_{-\infty}^{+\infty} t^{i\lambda v} (\lambda u + i\lambda v)(\lambda u - 1 + i\lambda v) \cdots \\ (\lambda u - k + 1 + i\lambda v) G(u + iv) dv. \end{aligned}$$

There exists a constant K , independent of u and v , such that

$$|\psi^{(k)}(t)| < K \int_{-\infty}^{+\infty} t^{\lambda u - k} ((u^2 + v^2)^{1/2})^{k - \text{Re } h} dv.$$

Substitute $v^2 = wu^2$ to write the last relation in the form

$$|\psi^{(k)}(t)| < (K/2) u^{k - \text{Re } h + 1} \int_{-\infty}^{+\infty} t^{\lambda u - k} ((1 + w)^{1/2})^{k - \text{Re } h} w^{-1/2} dw.$$

Since the integral is convergent, the assertion follows immediately upon setting $u = |t - 1|^{-1}$.

4. The function $\phi(t)$. First apply Theorem 3.1 to the order term in (2.3) to find that, near $t = 1$,

$$\begin{aligned} \left(\frac{d}{dt}\right)^h L \left\{ F(x/a) - a^b \sum_{m=0}^M B_m \Gamma(x + d_m - m) / \Gamma(x + d_m + b) \right\} \\ = O(|t - 1|^{b+M-k}), \end{aligned}$$

provided $k < M + \text{Re}(b)$. It now follows upon introducing

$$(4.1) \quad \begin{aligned} \phi(t) = L\{F(x/a)\}, & \quad (1 \leq t < \infty \text{ if } \lambda > 0), \\ & \quad (0 < t \leq 1 \text{ if } \lambda < 0), \end{aligned}$$

and using the well-known relation [1, p. 261]

$$\begin{aligned} L\{\Gamma(x + d_m - m) / \Gamma(x + d_m + b)\} \\ = (\Gamma(m + b))^{-1} t^{-\lambda(d_m + b - 1)} (t^\lambda - 1)^{m + b - 1}, \end{aligned}$$

that

$$(4.2) \quad \phi^{(k)}(t) = a^b \sum_{m=0}^M B_m (\Gamma(m+b))^{-1} \left(\frac{d}{dt} \right)^k [t^{-\lambda(d_m+b-1)} (t^\lambda - 1)^{m+b-1}] \\ + O(|t-1|^{b+M-k}),$$

provided $k < M + \operatorname{Re}(b)$.

Now note that by applying the recurrence relation $\Gamma(x+1) = x\Gamma(x)$ to (2.1), we obtain

$$(4.3) \quad F(x/a + 1) \prod_{j=0}^q (x + ar_j) = F(x/a) \prod_{j=0}^q (x + as_j).$$

There exist constants C_k and E_k ($k=0, 1, \dots, q+1$) such that

$$(4.4) \quad \prod_{j=0}^q (x + as_j) = \sum_{k=0}^{q+1} C_k \lambda x (\lambda x - 1) \cdots (\lambda x - k + 1), \\ \prod_{j=0}^q (x + ar_j) = \sum_{k=0}^{q+1} E_k (\lambda x + \lambda a) (\lambda x + \lambda a - 1) \cdots (\lambda x + \lambda a - k + 1).$$

Relations (4.3) and (4.4) are used to prove the assertion which follows.

THEOREM 4.1. *The function $\phi(t)$ defined by (4.1) satisfies the linear differential equation*

$$(4.5) \quad \sum_{k=0}^{q+1} (t^{k+\lambda a} C_k - t^k E_k) \phi^{(k)}(t) = 0,$$

where

$$(4.6) \quad C_k = (k!)^{-1} \prod_{j=0}^q (as_j + k/\lambda) - \sum_{n=0}^{k-1} C_n / (k-n)!, \\ E_k = (k!)^{-1} \prod_{j=0}^q (ar_j - a + k/\lambda) - \sum_{n=0}^{k-1} E_n / (k-n)!.$$

In particular

$$(4.7) \quad C_{q+1} = \lambda^{-q-1}; \quad C_q = \lambda^{-q} \left[a \sum_{j=0}^q s_j + q(q+1)(2\lambda)^{-1} \right], \\ E_{q+1} = \lambda^{-q-1}; \quad E_q = \lambda^{-q} \left[a \sum_{j=0}^q r_j + (q-2a\lambda)(q+1)(2\lambda)^{-1} \right].$$

PROOF. From (3.3) and (4.1) we may write

$$\begin{aligned}
\sum_{k=0}^{q+1} E_k t^k \phi^{(k)}(t) &= L \left\{ F(x/a) \sum_{k=0}^{q+1} E_k \lambda x (\lambda x - 1) \cdots (\lambda x - k + 1) \right\} \\
&= t^{\lambda a} L \left\{ F(x/a + 1) \sum_{k=0}^{q+1} E_k (\lambda x + \lambda a) (\lambda x + \lambda a - 1) \right. \\
&\quad \left. \cdots (\lambda x + \lambda a - k + 1) \right\} \\
&= t^{\lambda a} L \left\{ F(x/a) \sum_{k=0}^{q+1} C_k \lambda x (\lambda x - 1) \cdots (\lambda x - k + 1) \right\} \\
&= t^{\lambda a} \sum_{k=0}^{q+1} C_k t^k \phi^{(k)}(t),
\end{aligned}$$

which proves assertion (4.5). Equations (4.6) are obtained immediately upon substituting the appropriate values of x into the identities (4.4). Relations (4.7) are obtained by comparing the coefficients of x^{q+1} and x^q in the identities (4.4).

5. Recursion formulae. A general technique for obtaining the constants B_m in (2.3) is to substitute the asymptotic expansions (4.2) in (4.5), expand all the resulting terms about $t=1$, and equate the coefficients of like powers. This procedure will first be applied to find a finite recursion formula for the constants γ_m which occur in (1.4).

The method requires that we write (1.4) in the form of (2.3) with $b > q+1$. We choose $b = q+3$ to be consistent with the notation of [4]. In (1.4) set $x = \alpha w + \beta + q + 2$ and use the identity

$$((2\pi)^{1/2})^{\alpha-1} \Gamma(x+1) = \alpha^{x+1/2} \prod_{\mu=1}^{\alpha} \Gamma(x/2 + \mu/2)$$

to find

$$(5.1) \quad F_1(x/\alpha) = \alpha^{q+3} \sum_{m=0}^M \gamma_m \Gamma(x - q - 2 - m) / \Gamma(x+1) + O(x^{-q-4-M}),$$

where

$$\begin{aligned}
F_1(x) &= \prod_{j=0}^q \Gamma(x + \rho_j - \alpha^{-1}(\beta + q + 2)) \\
&\quad \cdot \left(\prod_{i=1}^p \Gamma(x + \sigma_i - \alpha^{-1}(\beta + q + 2)) \prod_{\mu=1}^{\alpha} \Gamma(x + \alpha^{-1}\mu) \right)^{-1}.
\end{aligned}$$

Equation (5.1) is in the form of (2.3) with

$$(5.2) \quad a = \alpha; \quad b = q + 3; \quad d_m = -q - 2 (m = 0, 1, \dots),$$

$$s_j = \rho_j - \alpha^{-1}(\beta + q + 2) \quad (j = 0, 1, \dots, q),$$

$$(5.3) \quad r_j = \sigma_{j+1} - \alpha^{-1}(\beta + q + 2) \quad (j = 0, 1, \dots, p-1),$$

$$= \alpha^{-1}(j - p + 1) \quad (j = p, \dots, q).$$

If now we choose $\lambda = 1$ in (4.2) we have simply

$$(5.4) \quad \phi^{(k)}(t) \sim \alpha^{q+3} \sum_{m=0}^{\infty} \frac{\gamma_m (t-1)^{m+q+2-k}}{\Gamma(m+q+3-k)} \quad (1 \leq t < \infty).$$

Substitution of (5.4) into (4.5) yields the desired recurrence relation for the constants γ_m .

THEOREM 5.1. *The coefficients γ_m ($m = 0, 1, \dots$), occurring in the expansion (1.4), satisfy the following recursion formula of length q :*

$$(5.5) \quad \gamma_m = (\alpha m)^{-1}(m+2)! \cdot \left\{ \sum_{i=1}^q \sum_{k=q-i}^{q+1} C_k \binom{k}{q-i} \gamma_{m-i} / (m+q-i+2-k)! \right. \\ \left. - \sum_{i=1}^{p-1} \sum_{k=q-i}^{q+1} E_k \binom{k-\alpha}{q-i-\alpha} \gamma_{m-i} / (m+q-i+2-k)! \right\},$$

($\gamma_0 = 1$ and $\gamma_{-1} = \gamma_{-2} = \dots = 0$), where

$$(5.6) \quad C_k = (k!)^{-1} \prod_{j=0}^q (\alpha \rho_j - \beta - q - 2 + k) - \sum_{n=0}^{k-1} C_n / (k-n)!,$$

$$E_k = (k!)^{-1} \prod_{j=0}^{p-1} (\alpha \sigma_{j+1} - \beta - q - 2 - \alpha + k) \prod_{j=p}^q (j - q + k) \\ - \sum_{n=0}^{k-1} E_n / (k-n)!.$$

In particular,

$$(5.7) \quad C_{q+1} = 1; \quad C_q = \alpha \sum_{j=0}^q \rho_j - (q+1)(\beta + q + 2) + q(q+1)/2,$$

$$E_{q+1} = 1; \quad E_q = \alpha \sum_{i=1}^p \sigma_i - p(\alpha + \beta + q + 2) \\ + q(q+1)/2 - \alpha(\alpha-1)/2.$$

PROOF. Equations (5.6) and (5.7) are immediately obtained by the substitution of (5.2) and (5.3) into (4.6) and (4.7). To find the recursion formula (5.5) set $\lambda = 1$ and employ the binomial theorem to write (4.5) in the form

$$\sum_{j=0}^{q+1} \sum_{k=j}^{q+1} C_k \binom{k}{j} (t-1)^{k-i} \phi^{(k)}(t) - \sum_{j=\alpha}^{q+1} \sum_{k=j}^{q+1} E_k \binom{k-\alpha}{j-\alpha} (t-1)^{k-i} \phi^{(k)}(t) = 0.$$

By (5.4) the coefficient of t^{m+2} in the asymptotic expansion for $(t-1)^{k-i} \phi^{(k)}(t)$ is

$$\alpha^{q+3} \gamma_{m-q+j} ((m+j+2-k)!)^{-1},$$

so that, for each integer m ,

$$\sum_{j=0}^{q+1} \sum_{k=j}^{q+1} C_k \binom{k}{j} \gamma_{m-q+j} ((m+j+2-k)!)^{-1} - \sum_{j=\alpha}^{q+1} \sum_{k=j}^{q+1} E_k \binom{k-\alpha}{j-\alpha} \gamma_{m-q+j} ((m+j+2-k)!)^{-1} = 0,$$

provided we agree that $\gamma_m = 0$ for $m < 0$. From (5.7), the coefficients of γ_{m+1} and γ_m are 0 and $\alpha m / (m+2)!$ respectively. The above formula therefore reduces to (5.5) upon solving for γ_m and setting $i = q - j$.

The general method may also be applied to obtain a recursion formula for the constants c_m which appear in (1.2). In this case set $x = \alpha w + \beta - q - 3$ and write (1.2) as

$$(5.8) \quad F_2(x/\alpha) = \alpha^{q+3} \sum_{m=0}^M c_m \Gamma(x) / \Gamma(x + q + 3 + m) + O(x^{-q-4-M}),$$

where

$$F_2(x) = \prod_{i=1}^p \Gamma(x + \sigma_i - \alpha^{-1}(\beta - q - 3)) \cdot \prod_{\mu=1}^{\alpha} \Gamma(x + \alpha^{-1}(\mu - 1)) \bigg/ \prod_{j=0}^q \Gamma(x + \rho_j - \alpha^{-1}(\beta - q - 3)).$$

Now (5.8) is in the form of (2.3) with

$$(5.9) \quad a = \alpha; \quad b = q + 3; \quad d_m = m,$$

$$(5.10) \quad s_j = \sigma_{j+1} - \alpha^{-1}(\beta - q - 3) \quad (j = 0, 1, \dots, p-1)$$

$$= \alpha^{-1}(j - p) \quad (j = p, \dots, q),$$

$$r_j = \rho_j - \alpha^{-1}(\beta - q - 3) \quad (j = 0, 1, \dots, q).$$

If we choose $\lambda = -1$, then (4.2) and (4.5) become simply

$$(5.11) \quad \phi^{(k)}(t) \sim \alpha^{q+3} \sum_{m=0}^{\infty} (-1)^{m+q} c_m (t-1)^{m+q+2-k} / \Gamma(m+q+3-k) \quad (0 < t \leq 1)$$

and

$$(5.12) \quad \sum_{k=0}^{q+1} (t^{k-\alpha} C_k - t^k E_k) \phi^{(k)}(t) = 0$$

respectively. The recursion formula for the c_m obtained by substituting (5.11) into (5.12) is precisely the result given in [4].

6. Remark. When Theorem (5.1) is applied to the special case of

$$(6.1) \quad (g(w))^{-1} = \Gamma(w + \rho_0) \Gamma(w + \rho_1)$$

the result is

$$(6.2) \quad \gamma_m (8m)^{-1} [4(\rho_1 - \rho_0)^2 - (2m - 1)^2] \gamma_{m-1}.$$

On the other hand, van der Corput's formula [8, Equation 8] gives

$$\gamma_1 = (8)^{-1} [4(\rho_1 - \rho_0)^2 + 71] \gamma_0$$

for the second coefficient. This discrepancy has been resolved by H. O. Pollak who has computed γ_1 for this special case by substituting Stirling's formula into (6.1) directly. The result is in agreement with (6.2).

BIBLIOGRAPHY

1. A. Erdelyi et al., *Tables of integral transforms*, vol. 1, McGraw-Hill Book Company, Inc., 1954.
2. W. B. Ford, *The asymptotic developments of functions defined by Maclaurin series*, University of Michigan Studies, Scientific Series, vol. 11, 1936.
3. H. K. Hughes, *The asymptotic developments of a class of entire functions*, Bull. Amer. Math. Soc. vol. 51 (1945) pp. 456-461.
4. T. D. Riney, *A finite recursion formula for the coefficients in asymptotic expansions*, Trans. Amer. Math. Soc. vol. 88 (1958) pp. 214-226.
5. ———, *On the coefficients in asymptotic factorial expansions*, Proc. Amer. Math. Soc. vol. 7 (1956) pp. 245-249.
6. E. M. Wright, *A recursion formula for the coefficients in an asymptotic expansion*, to be published in Proc. Glasgow Math. Assoc.
7. ———, *The asymptotic expansion of the generalized hypergeometric function*, Proc. London Math. Soc. (2) vol. 46 (1940) pp. 389-408.
8. J. G. van der Corput, *On the coefficients in certain asymptotic factorial expansions*, Akad. Wet. (Amsterdam) Proc. Series A vol. 60 no. 4 (1957) pp. 337-351.

BELL TELEPHONE LABORATORIES, INC., ALLENTOWN, PA.