ON THE NOTION OF ANALYTICITY

C. F. MOPPERT

Let E_n be a normed n -dimensional vector space over a field Γ . We denote by $a, b, \dots$ the elements of E_n , with $\alpha, \beta, \dots$ the elements of Γ ; $a = \alpha_1 e_1 + \dots + \alpha_n e_n \sim \{\alpha_1, \dots, \alpha_n\}$ where $e_1, \dots, e_n$ is a basis of E_n . The norm in E_n we denote by $|\cdot|_n$.

By $a = L \circ b$ we denote a linear homogeneous transformation on E_n to E_n , i.e. $a \in E_n$ is defined for every $b \in E_n$; $L \circ (\lambda b) = \lambda L \circ b$; $L \circ (b_1 + b_2) = L \circ b_1 + L \circ b_2$ for every λ, b, b_1, b_2 . Naturally, given a basis $e_1, \dots, e_n$ of E_n , L can be represented as a square matrix: $L \sim [\lambda_{\mu\nu}]$ such that $\alpha_\mu = \lambda_{\mu 1} \beta_1 + \dots + \lambda_{\mu n} \beta_n$ ($\mu = 1, \dots, n$) for $a = L \circ b$, $a \sim \{\alpha_1, \dots, \alpha_n\}$, $b = \{\beta_1, \dots, \beta_n\}$.

Let $y = f(x)$ be a function on E_n to E_n , i.e. $y \in E_n$ for some $x \in E_n$. $f(x)$ is called *differentiable* (in the sense of *Fréchet*) at $x = x_0$ if for $h \in E_n$, $|h|_n \rightarrow 0$ there is a $L = L(x_0)$ such that

$$(1) \quad f(x_0 + h) = f(x_0) + L(x_0) \circ h + r(h); r(h) \in E_n, |r(h)|_n = o(|h|_n).$$

If, with respect to the basis $e_1, \dots, e_n$ it is $f \sim \{\phi_1, \dots, \phi_n\}$, $x \sim \{\xi_1, \dots, \xi_n\}$, $L \sim [\lambda_{\mu\nu}]$ then obviously

$$(2) \quad \lambda_{\mu\nu} = \partial \phi_\mu / \partial \xi_\nu \quad (x = x_0).$$

We call $\otimes$ a *product on E_n* if to every $a \in E_n$, $b \in E_n$ there exists $c = a \otimes b \in E_n$ such that $a \otimes b = L_a \circ b = L_b \circ a$. For some fixed basis $e_1, \dots, e_n$ with $a \sim \{\alpha_1, \dots, \alpha_n\}$, $b \sim \{\beta_1, \dots, \beta_n\}$, $c \sim \{\gamma_1, \dots, \gamma_n\}$ such a product can always be realized by n^3 constants $\pi_{\rho\mu\nu}$ such that

$$(3) \quad \gamma_\rho = \sum_{\mu, \nu=1}^n \pi_{\rho\mu\nu} \alpha_\mu \beta_\nu \quad (\rho = 1, \dots, n).$$

Naturally, by introducing a product in a vector space this vector space becomes an algebra.

If $f(x)$ is a function on E_n to E_n we call it *analytic with respect to the product $\otimes$* at $x = x_0$ if it is there differentiable and if there is an $l = l(x_0) \in E_n$ such that

$$(4) \quad L(x_0) \circ h = l(x_0) \otimes h,$$

Received by the editors December 21, 1957 and, in revised form, November 10, 1958.

where $L(x_0)$ is defined by (1). Without referring to the definition of differentiability this means: there is an $l(x_0)$ such that

$$(5) \quad \begin{aligned} f(x_0 + h) &= f(x_0) + l(x_0) \otimes h + r(h), & r(h) &\in E_n, \\ |r(h)|_n &= o(|h|_n) & \text{for } |h|_n &\rightarrow 0. \end{aligned}$$

It is obvious that this definition of analyticity contains the one for functions of a complex variable.

Let $l(x_0) \sim \{\lambda_1(x_0), \dots, \lambda_n(x_0)\}$, $h \sim \{\eta_1, \dots, \eta_n\}$ for a basis $e_1, \dots, e_n$. Putting $l(x_0) \otimes h = k$, $k \sim \{\kappa_1, \dots, \kappa_n\}$ we have $\kappa_\rho = \sum_{\mu, \nu=1}^n \pi_{\rho\mu\nu} \lambda_\mu(x_0) \eta_\nu$ ($\rho = 1, \dots, n$). On the other hand, the ρ th component of $L(x_0) \circ h$ is given by

$$\lambda_{\rho 1} \eta_1 + \dots + \lambda_{\rho n} \eta_n = \partial \phi_\rho / \partial \xi_1 \cdot \eta_1 + \dots + \partial \phi_\rho / \partial \xi_n \cdot \eta_n.$$

Instead of (4) we have therefore

$$(6) \quad \partial \phi_\rho / \partial \xi_\nu = \pi_{\rho 1\nu} \lambda_1(x_0) + \dots + \pi_{\rho n\nu} \lambda_n(x_0) \quad (\rho, \nu = 1, \dots, n).$$

We put

$$\Pi_\nu = \text{Det } \pi_{\rho\mu\nu} = \begin{vmatrix} \pi_{11\nu} & & \\ & \ddots & \\ & & \pi_{nn\nu} \end{vmatrix} \quad (\nu = 1, \dots, n)$$

and denote by $\Pi_{\nu\rho}$ the determinant we get from Π_ν by replacing its ρ th column by $\partial \phi_1 / \partial \xi_\nu, \dots, \partial \phi_n / \partial \xi_\nu$. It follows

$$(7) \quad \Pi_{1\rho} / \Pi_1 = \dots = \Pi_{n\rho} / \Pi_n (= \lambda_\rho) \quad (\rho = 1, \dots, n),$$

provided $\Pi_\nu \neq 0$ ($\nu = 1, \dots, n$). If some of these determinants vanish, $\Pi_1 = \dots = \Pi_k = 0$, $k \leq n$ say, then (7) has to be replaced by

$$(8) \quad \begin{aligned} \Pi_{1\rho} &= \dots = \Pi_{k\rho} = 0, \\ \Pi_{k+1, \rho} / \Pi_{k+1} &= \dots = \Pi_{n\rho} / \Pi_n \end{aligned} \quad (\rho = 1, \dots, n).$$

The equations (7) viz (8) form a necessary condition for the analyticity of $f(x)$ at $x = x_0$. We are therefore entitled to consider them as a generalization of the Cauchy-Riemann differential equations.

In the case $n=2$ and complex multiplication, the equations (7) yield the Cauchy-Riemann equations. In the case $n=4$, E_n being a real space and the product being defined according to the multiplication of quaternions, these equations read

$$\begin{aligned}
\phi_{11} &= \phi_{22} = \phi_{33} = \phi_{44}, \\
\phi_{21} &= -\phi_{12} = \phi_{43} = -\phi_{34}, \\
\phi_{31} &= -\phi_{42} = -\phi_{13} = \phi_{24}, \\
\phi_{41} &= \phi_{32} = -\phi_{23} = -\phi_{14}
\end{aligned}$$

where we have abbreviated $\partial\phi_\nu/\partial\xi_\mu$ by $\phi_{\nu\mu}$. In the case $n=3$, E_n being real and the product being the ordinary vector product, the function $f(\xi_1, \xi_2, \xi_3) = \{\xi_2 - \xi_3, \xi_3 - \xi_1, \xi_1 - \xi_2\}$ turns out to be everywhere analytic.

UNIVERSITY OF MELBOURNE, MELBOURNE, AUSTRALIA