MEROMORPHIC FUNCTIONS OF FINITE ORDER

S. M. SHAH

1. Introduction. Let $f(z)$ be an entire function of finite order ρ . If the genus p of the canonical product (c.p.) formed with zeros of $f(z)$ is equal to the genus of $f(z)$, then it is known¹ that [9; 10]

$$\liminf_{r \rightarrow \infty} \frac{\log M(r)}{n(r, 0)\phi(r)} = 0$$

where $\phi(r)$ is any positive nondecreasing function such that

$$(1) \quad \int_1^\infty dt/t\phi(t)$$

is convergent. Another proof of this theorem is given by Boas [2] where he assumes only that $\phi(r)$ is positive and (1) exists. Following the method of Boas, Shanker Hari has shown in a paper under publication that if $f(z)$ is a c.p. of order and genus ρ then

$$\liminf_{r \rightarrow \infty} \frac{\log M(r)}{N(r, 0)\phi(r)} = 0.$$

In this paper we prove a more general theorem by the method given in [9; 10].

Let G , p_1 , p_2 denote respectively the genus of a meromorphic function $f(z)$, the genus of the c.p. P_1 formed with zeros of f and the genus of the c.p. P_2 formed with poles of f . Write² $S = \max(p_1, p_2)$, $n(t) = n(t, f-0) + n(t, f-\infty)$, $n_1(t) = n(t, P_1-0) + n(t, P_2-0)$. Let k denote any constant ≥ 1 , $\phi(x)$ any positive nondecreasing function such that (1) exists, $(a_i(z))$, $i=0, 1, \dots, l$ any meromorphic functions (including rational functions or finite constants) such that

$$(2) \quad T(r, a_i(z)) = o(T(r, f)), \quad i = 0, 1, \dots, l$$

and $g_0(z) = g(z)$, $g_1(z)$, $g_2(z)$ any meromorphic functions (including rational functions or constants, finite or infinite) such that

$$(3) \quad T(r, g_i(z)) = o(T(r, f)), \quad i = 0, 1, 2.$$

Presented to the Society, November 29, 1958; received by the editors December 2, 1958.

¹ For notations see 7 [7; 8].

² The function $n(r, f-g)$ denotes the number of zeros of $f-g$ in $|z| \leq r$. If g is infinite constant, then $N(r, 1/(f-g)) = N(r, f)$.

Write

$$\psi(f) = \sum_0^l a_i(z) f^{(i)}(z), \quad \psi(f - g_1) = \sum_0^l a_i(z) \{f^{(i)}(z) - g_1^{(i)}(z)\}$$

when g_1 is not infinite constant; we suppose that (a_i) are so chosen that neither $\psi(f)$ nor $\psi(f - g_1)$ reduces identically to a constant.

THEOREM 1. *Let $f(z)$ be a meromorphic function of integer order ρ .*

(a) *If $G = S$, then*

$$(4) \quad \liminf_{r \rightarrow \infty} \frac{T(kr, f)}{\{n(r, 0) + n(r, \infty)\} \phi(r)} = 0,$$

$$(5) \quad \liminf_{r \rightarrow \infty} \frac{T(kr, f)}{\{N(r, 0) + N(r, \infty)\} \phi(r)} = 0.$$

(b) *If $G > S$, then*

$$(6) \quad \bar{N}\left(r, \frac{1}{\psi(f) - 1}\right) \sim T(r, f) \sim Ar^\rho \quad (0 < A < \infty).$$

THEOREM 2. *Let $f(z)$ be a meromorphic function of noninteger order $\rho > 0$. Then*

(a)

$$(7) \quad \liminf_{r \rightarrow \infty} \frac{T(kr, f)}{n(r, f - g_1) + n(r, f - g_2)} \leq k^{1+S} B(S),$$

$$(8) \quad \liminf_{r \rightarrow \infty} \frac{T(kr, f)}{N(r, 1/(f - g_1)) + N(r, 1/(f - g_2))} \leq k^{1+S} (S + 1) B(S)$$

where

$$\begin{aligned} S = [\rho], \quad A(S) &= 2 & \text{if } S = 0, \\ &= 2e(2 + \log S)(S + 1)^2/S & \text{if } S \geq 1, \end{aligned}$$

and

$$B(S) = A(S)/(\rho - S)(1 + S - \rho)$$

and g_1, g_2 are any two distinct functions satisfying (3).

COROLLARY.

$$(9) \quad \liminf_{r \rightarrow \infty} \frac{T(r, f)}{n(r, f - g)} \leq 2B(S),$$

$$(10) \quad \liminf_{r \rightarrow \infty} \frac{T(r, f)}{N(r, 1/(f - g))} \leq 2(S + 1)B(S)$$

except possibly for one g in each relation (9), (10).

(b) If (10) does not hold for $g = g_1$, then for all $r > r_0$,

$$(11) \quad T(r, f) < \left\{ \frac{2(S + 1)B(S)}{2(S + 1)B(S) - 1} \right\} \{N(r, f) + \overline{N}(r, 1/(\psi(f - g_1) - 1))\}$$

if $g_1 \neq \infty$ and

$$(12) \quad T(r, f) < \left\{ \frac{2(S + 1)B(S)}{2(S + 1)B(S) - 1} \right\} \left\{ N\left(r, \frac{1}{f}\right) + \overline{N}\left(r, \frac{1}{\psi(f) - 1}\right) \right\}$$

if $g_1 \equiv \infty$.

(c) If (9) does not hold for $g = g_1$, then we have for a sequence of values of $r \uparrow \infty$,

$$(13) \quad T(r, f) < \left\{ \frac{2\rho B(S)}{2\rho B(S) - 1} \right\} \{N(r, f) + \overline{N}(r, 1/(\psi(f - g_1) - 1))\}$$

if $g_1 \neq \infty$ and

$$(14) \quad T(r, f) < \left\{ \frac{2\rho B(S)}{2\rho B(S) - 1} \right\} \{N(r, 1/f) + \overline{N}(r, 1/(\psi(f) - 1))\}$$

if $g_1 \equiv \infty$.

THEOREM 3. Let $f(z)$ be an entire function of order ρ and let, when $\rho > 0$, $h(\rho) = (\rho + (1 + \rho^2)^{1/2})(1 + (1 + \rho^2)^{1/2})^\rho / \rho^\rho$. If $\rho > 0$, then

$$(15) \quad \liminf_{r \rightarrow \infty} \frac{\log M(kr)}{n(r, f - g)} \leq \frac{2k^\rho h(\rho)}{\rho},$$

$$(16) \quad \liminf_{r \rightarrow \infty} \frac{\log M(kr)}{N(r, 1/(f - g))} \leq 2k^\rho h(\rho)$$

for every entire function $g(z)$ (including a polynomial or a finite constant) satisfying (3), with one possible exception. If $\rho = 0$, then

$$(17) \quad \liminf_{r \rightarrow \infty} \frac{\log M(kr)}{N(r, 1/(f - g))} = 1$$

for every entire function $g(z)$ such that $M(r, g) = o(M(r, f))$.

THEOREM 4. Let $f(z)$ be a meromorphic function of order ρ . If $\rho > 0$, then

$$(18) \quad \liminf_{r \rightarrow \infty} \frac{T(kr, f)}{n(r, f - g)} \leq \frac{3k^p}{\rho},$$

$$(19) \quad \liminf_{r \rightarrow \infty} \frac{T(kr, f)}{N(r, 1/(f - g))} \leq 3k^p$$

for every meromorphic function $g(z)$ satisfying (3), with two possible exceptions. If $\rho = 0$ then

$$(20) \quad \liminf_{r \rightarrow \infty} \frac{T(kr, f)}{N(r, 1/(f - g_1)) + N(r, 1/(f - g_2))} \leq 1$$

where g_1, g_2 are any two distinct functions satisfying (3).

2. Remarks. (i) Let $\delta(\alpha)$ denote Nevanlinna defect. If $\sum \delta(\alpha) < 2$, then $G = S$ and (4) holds. If $\sum \delta(\alpha) = 2$, then (4) may or may not hold. (See [13, pp. 590–591] for an entire function of which the genus is greater than the genus of its c.p. and for which $\sum \delta(\alpha) = 2$ and (4) holds.) Conversely if (4) holds, then $\sum \delta(\alpha)$ may have its maximum value 2 or $\sum \delta(\alpha) < 2$ ($f(z) = \sin z$). If (4) does not hold then $G > S$, $\delta(0) = \delta(\infty) = 1$.

(ii) For entire functions of order ρ , $0 \leq \rho < 1$, it is known that [1; 11]

$$\liminf_{r \rightarrow \infty} \frac{\log M(r)}{N(r, a)} \leq \frac{1}{1 - \rho}$$

and for meromorphic functions of order ρ , $0 \leq \rho < 1$, it is known that

$$\liminf_{r \rightarrow \infty} \frac{T(r)}{N(r, a) + N(r, b)} \leq \frac{1}{1 - \rho}.$$

More precise results in this direction have been given by Edrei and Fuchs [4].

(iii) For entire functions of noninteger order $\rho > 0$, it is known that

$$L(f, a) = \liminf_{r \rightarrow \infty} \frac{\log M(r)}{n(r, f - a)} < \infty$$

for every finite a . A sharp upper bound for $L(f, a)$ when $\rho > 1$, is not known but by (15) $L(f, a) \leq 2h(\rho)/\rho$ except possibly for one value of a . This exceptional value may exist. Consider [7, pp. 18–19]

$$f(z) = \prod_2^\infty E\left(\frac{z}{a_n}, p\right), \quad a_n = -n^{1/\rho}, \quad p < \rho < p + 1.$$

$f(z)$ is an entire function of order ρ and

$$\log M(r)/n(r, 0) \sim \lambda/\sin \lambda(\rho - p)$$

which can be made greater than $2h(\rho)/\rho$ by choosing $\rho - p$ sufficiently small.

(iv) If (15), (16), (18) or (19) do not hold for $g = g_1$ then we can obtain relations similar to (11)–(14). For instance if (15) or (16) does not hold when $g = g_1$, then

$$\liminf_{r \rightarrow \infty} \frac{T(r, f)}{N(r, 1/(\psi(f - g_1) - 1))} \leq 2.$$

3. Theorem 1.

LEMMA 1. If $f(z) = z^v \exp(Q(z))P_1(z)/P_2(z)$ is a proper meromorphic function³ of order $\rho \geq 0$ such that $G = S$, then for all $r > r_0$

$$T(kr, f) \leq A(S)k^{1+S}J(r, S)$$

where⁴

$$\begin{aligned} A(S) &= 2 && \text{if } S = 0, \\ &= 2e(2 + \log S)(S + 1)^2/S && \text{if } S \geq 1, \end{aligned}$$

and

$$J(r, S) = r^{1+S} \int_0^\infty \frac{n_1(t)dt}{t^{1+S}(t+r)}.$$

PROOF. We have

$$T(kr, f) = O(r^S + \log r) + \log M(kr, P_1) + \log M(kr, P_2)$$

and when $p_1 \geq 1$ [8, pp. 225–226]

$\log M(kr, P_1)$

$$\leq (2 + 1/p_1)e(2 + \log p_1)(1 + p_1)k^{1+p_1}r^{1+p_1} \int_0^\infty \frac{n(t, P_1 - 0)dt}{t^{1+p_1}(t + kr)}.$$

Hence, when $p_1 = p_2 \geq 1$,

$$T(kr, f) \leq \frac{2e(2 + \log S)(1 + S)^2}{S} k^{1+S}J(r, S).$$

If $p_1 = p_2 = 0$, then

$$T(kr, f) < O(\log r) + kr \int_0^\infty \frac{n_1(t)dt}{t(t + kr)}.$$

³ f is not a rational function.

⁴ We can take $A(0) = 1 + c$, where c is any positive number.

If $p_1 > p_2$ then $\log M(kr, P_2) = o(r^{1+p_2}) = o(r^{p_1})$ and a similar result if $p_1 < p_2$. Hence the lemma is proved.

LEMMA 2. If $f(z)$ is a proper meromorphic function of finite order and

$$\psi(f) = \psi(z) = \sum_0^l a_i(z)f^{(i)}(z)$$

then

$$(21) \quad (1 + o(1))T(r, f) < \overline{N}(r, f) + \overline{N}\left(r, \frac{1}{\psi(f) - 1}\right) + N\left(r, \frac{1}{f}\right).$$

This lemma is substantially contained in Theorem 7 of [5].

LEMMA 3. If $f(z) = z^v \exp(Q(z))P_1(z)/P_2(z)$ and $G > S$ then $Q(z) = a_p z^p + \dots$, $|a_p| = A \neq 0$ and

- (i) $T(r, f) \sim Ar^p$,
- (ii) $T(r, f') \sim Ar^p$,
- (iii) $T(r, \psi) = T(r, \sum_0^l a_i f^{(i)}) \sim Ar^p$.

PROOF. Write $f = b_0 \exp(Q)$ where $b_0 = z^v P_1 P_2^{-1}$,

$$T(r, f) \leq T(r, \exp Q) + T(r, b_0),$$

$$T(r, \exp Q) \leq T(r, f) + T(r, b_0) + O(1)$$

and since $T(r, \exp Q) \sim Ar^p$, $T(r, b_0) = o(r^p)$, (i) follows.

$$(ii) \quad f' = e^Q \{ k z^{k-1} P_1 P_2^{-1} + z^k Q' P_1 P_2^{-1} + z^k P_1' P_2^{-1} - Z^k P_1 P_2^{-2} P_2' \} \\ = e^Q b_1 \text{ (say).}$$

Then $T(r, b_1) = o(r^p)$ and so $T(r, f') \sim Ar^p$.

(iii) $f'' = (\exp Q)b_2$ where $T(r, b_2) = o(r^p)$ and so on. Hence

$$\begin{aligned} T(r, \psi(f)) &= T(r, (\exp Q) \sum a_i b_i) \\ &\leq T(r, \exp Q) + T(r, \sum a_i b_i) \\ &\leq (A + o(1))r^p + \sum T(r, a_i) + \sum T(r, b_i) \\ &= (A + o(1))r^p. \end{aligned}$$

Also

$$\exp Q = \psi(f) / \sum a_i b_i, \quad T(r, \exp Q) \leq T(r, \psi(f)) + o(r^p)$$

and hence $T(r, \psi(f)) \sim Ar^p$.

(a) PROOF OF⁵ (4). Since $\rho \geq 1$ and $G = S$, $n(x)$ tends to infinity with

⁵ (4) can also be proved by the method given in [12, pp. 696-697].

x . Further $T(kr, f) \sim T(kr, fz^v)$ where v is any integer and so we may suppose $f(0) \neq 0, \neq \infty$. Then $f(z) = \exp(Q(z))P_1(z)/P_2(z)$. Since $G = S$, we have $S = \rho$ or $\rho - 1$. Consider first when $S = \rho$. If $n(x) = O(x^S)$ we have

$$J(r, S) < c_1 r^S \int_0^r \frac{n(x) dx}{x^{1+S}}$$

and hence from Lemma 1 of [9, p. 24] we get

$$\liminf_{r \rightarrow \infty} J(r, S)/n(r)\phi(r) = 0.$$

If $\limsup_{x \rightarrow \infty} n(x)/x^S = \infty$, we write $y = \log x$ and $\log(n(x)/x^S) = \log \mu(x) = \psi(y)$. Then $\limsup_{y \rightarrow \infty} \psi(y) = \infty$, and since $f(z)$ is of order $\rho = S$, $\limsup_{y \rightarrow \infty} \psi(y)/y = 0$. Apply Lemma 2 of [9, p. 25]. There exists a sequence $r_n \uparrow \infty$ for which

$$\begin{aligned} \mu(x) &< \mu(r_n), & \Delta < x < r_n, \\ \frac{\log \mu(x)}{\log x} &< \frac{\log \mu(r_n)}{\log r_n}, & x > r_n. \end{aligned}$$

Hence

$$J(r_n, S) < c_2 n(r_n) \log r_n.$$

Consider now $S = \rho - 1$. Given $\phi(x)$ we can find [10] a function $\phi_1(x)$ such that for all large x : (i) $\phi_1(x) \leq \phi(x)$; (ii) $\phi_1(x)/x^\alpha$ is nonincreasing, α being a positive number less than unity; (iii) $\int_1^\infty dt/t\phi_1(t)$ is convergent. If now

$$\limsup_{x \rightarrow \infty} n(x)\phi_1(x)/x^{1+S} > 0,$$

then for a sequence $R_n \uparrow \infty$, $J(R_n, S) = o(n(R_n)\phi_1(R_n))$. If $n(x)\phi_1(x) = o(x^{1+S})$ then we choose in Lemma 3 of [10, p. 184] $0 < \beta < 1 - \alpha$, $\theta(x) = x^\beta$, $\Psi(x) = n(x)\phi_1(x)/x^{S+1-\beta}$ and obtain for $x_n \uparrow \infty$, $J(x_n, S) = o(\phi(x_n)n(x_n))$ and (2) is proved.

Since $n(r, a) \leq N(2r, a)/\log 2$, (5) follows from (4).

(b) PROOF OF (6). By hypothesis

$$N(r, f) = o(r^\rho), \quad N(r, 1/f) = o(r^\rho),$$

and so from (21)

$$(1 + o(1))T(r, f) < \overline{N}(r, 1/(\psi(f) - 1)) < T(r, \psi(f)) + O(1)$$

and so (6) follows from Lemma 3(i) and (iii).

4. Theorems 2, 3, 4.

LEMMA 4. If $f(z)$ is a meromorphic function of order $\rho > 0$, then

$$I_1(r) = \int_1^r \frac{T(kt, f)}{t} dt < \frac{k^\rho}{\rho} (1 + \epsilon) T(r, f)$$

for a sequence of values of r tending to infinity.

LEMMA 5. If $f(z)$ is an entire function of order $\rho > 0$, then

$$I_2(r) = \int_1^r \frac{\log M(kt)}{t} dt < \frac{k^\rho}{\rho} h(\rho) (1 + \epsilon) T(r, f)$$

where $h(\rho)$ has been defined in the statement of Theorem 3, for a sequence of values of r tending to infinity.

The proof of Lemma 4 is straightforward (cf. [14, p. 321]) and the proof of Lemma 5 depends on the relation

$$\log M(r) \leq \frac{h+1}{h-1} T(hr), \quad h > 1.$$

We obtain for a sequence of values of $r \rightarrow \infty$,

$$I_2(r) \leq \frac{h+1}{h-1} \frac{h^\rho k^\rho}{\rho} (1 + \epsilon) T(r).$$

Choose $h = (1 + (1 + \rho^2)^{1/2})/\rho$.

(a) PROOF OF (7). We have $S = [\rho]$

$$T(kr, f) < A(S) k^{1+S} J(r, S), \quad r > r_0.$$

We choose in Lemma 3 of [10, p. 184] (see also [12, p. 69]) $\Psi(x) = n_1(x)/x^{\rho-\epsilon}$ and obtain that for $r = r_n \uparrow \infty$

$$T(kr_n, f) < B(S)(1 + \epsilon') k^{1+S} n(r_n).$$

Let $H(z) = (f(z) - g_1(z))/(f(z) - g_2(z))$, $g_1 \not\equiv \infty$, $g_2 \not\equiv \infty$. Then $T(r, H) \sim T(r, f)$. Also for a sequence $R_n \uparrow \infty$,

$$T(kR_n, H) < B(S)(1 + \epsilon') k^{1+S} \{n(R_n, H - 0) + n(R_n, H - \infty)\}.$$

Now

$$n(R_n, H - 0) + n(R_n, H - \infty) < \sum_1^2 n(R_n, f - g_i) + \sum_1^2 n(R_n, g_i - \infty).$$

Suppose now $k > 1$. Then

$$n(R_n, g_i - \infty) < N(kR_n, g_i)/\log k = o(T(kR_n, f)).$$

Hence for $n > n_1$

$$(22) \quad T(kR_n, f) < B(S)(1 + \epsilon')k^{1+S} \sum_1^2 n(R_n, f - g_i).$$

If we take $H(z) = f(z) - g_2(z)$, $g_2 \not\equiv \infty$, we get for another $\{R_n\} \uparrow \infty$,

$$(23) \quad T(kR_n, f) < B(S)(1 + \epsilon')k^{1+S} \{n(R_n, f - \infty) + n(R_n, f - g_2)\}$$

and so from (22) and (23)

$$(24) \quad \liminf_{r \rightarrow \infty} T(kr, f) \Big/ \sum_1^2 n(r, f - g_i) \leq B(S)k^{1+S}$$

and (7) is proved when $k > 1$. Further k can be chosen arbitrarily near to 1 in the relation

$$\liminf_{r \rightarrow \infty} T(r, f) \Big/ \sum_1^2 n(r, f - g_i) \leq B(S)k^{1+S}$$

and so (7) is true when $k \geq 1$.

PROOF OF (8). Write

$$N_1(t) = \int_0^r \{n_1(t)/t\} dt; \quad N(t) = N(t, f) + N(t, 1/f).$$

Then

$$\begin{aligned} J(r, S) &= \int_0^\infty \frac{r^{1+S} n_1(t) dt}{t^{1+S}(t+r)} \\ &< (S+1) \left\{ r^S \int_0^r \frac{N_1(t) dt}{t^{1+S}} + r^{1+S} \int_r^\infty \frac{N_1(t) dt}{t^{2+S}} \right\} \\ &< (S+1)(1+\epsilon) \left\{ r^S \int_{x_0}^r \frac{[N(t)] dt}{t^{1+S}} + r^{1+S} \int_r^\infty \frac{[N(t)] dt}{t^{S+2}} \right\} \end{aligned}$$

if $r > x_1(\epsilon)$. Hence by Lemma 3 of [10, p. 184] we get

$$\liminf_{r \rightarrow \infty} \frac{T(kr, f)}{N(r)} \leq B(S)(1+S)k^{1+S}$$

and the rest of the argument is the same as for (7). Hence (8) is proved. The Corollary follows from (7) and (8).

(b) If (10) is false for $g = g_1 \not\equiv \infty$, then

$$\liminf_{r \rightarrow \infty} \frac{T(r, f)}{N(r, 1/(f - g_1))} = \beta > 2(S+1)B(S).$$

Hence for all $r > r_0(\epsilon)$,

$$T(r, f) > (\beta - \epsilon)N(r, 1/(f - g_1))$$

We apply Lemma 2 to $f - g_1$ and obtain

$$\begin{aligned} \{1 + o(1)\} T(r, f - g_1) \\ \leq \overline{N}(r, f - g_1) + \overline{N}\left(r, \frac{1}{\psi(f - g_1) - 1}\right) + N\left(r, \frac{1}{f - g_1}\right), \end{aligned}$$

and for $r > r_1(\epsilon)$

$$\left(1 - \frac{1}{\beta - \epsilon} - \epsilon\right) T(r, f) \leq N(r, f) + \overline{N}\left(r, \frac{1}{\psi(f - g_1) - 1}\right).$$

Hence for $r > r_2$

$$T(r, f) < \left\{ \frac{2(S+1)B(S)}{2(S+1)B(S) - 1} \right\} \left\{ N(r, f) + \overline{N}\left(r, \frac{1}{\psi(f - g_1) - 1}\right) \right\}.$$

The proof of (12) is similar.

(c) We omit the proofs of (13)–(14) which can be proved with the help of Lemma 4.

(d) The first part of Theorem 3 follows from Lemma 5 and (17) follows from a result of Boas [1, pp. 6–7; 3, p. 48]. We omit the proofs of Theorems 3 and 4.

5. Example. We show that (4), (5) are best possible in the sense that k cannot be replaced by a function $\alpha(r)$ tending to infinity with r . We prove that *given $\alpha(r) \rightarrow \infty$ with r , there is a function $\phi(x)$ such that (1) exists, and an entire function (c.p.) $f(z)$ of integer order ρ and genus $\rho - 1$ such that*

$$\begin{aligned} (25) \quad \liminf_{r \rightarrow \infty} \frac{T(r\alpha(r), f)}{N(r, 0)\phi(r)} &= \liminf_{r \rightarrow \infty} \frac{T(r\alpha(r), f)}{n(r, 0)\phi(r)} \geq \frac{1}{|(1 + 2^{1/2})^2|} \quad \text{if } \rho = 1 \\ &= \infty \quad \text{if } \rho > 1. \end{aligned}$$

(a) We suppose that $\alpha(r)$ satisfies the following:

(i) $\alpha(r) > 0$ for $r \geq x_0 > 0$ and tends to infinity with r .

(ii) $\alpha'(r) > 0$ and $r\alpha'(r) \downarrow$ for $r \geq x_0$.

(iii) $r^2\alpha'(r)/\alpha^2(r)$ is strictly increasing for $r \geq x_0$. For instance we can take $\alpha(r) = l_k r$, $k = 1, 2, \dots$, ($l_1 r = \log r$; $l_k(r) = \log(l_{k-1}r)$). A function of slower growth and satisfying (i)–(iii) is given by the functional equation $\alpha(e^x) = e\alpha(x)$. Let

$$\begin{aligned} \phi(x) &= \alpha^2(x)/x\alpha'(x) & x \geq x_0, \\ &= \phi(x_0) & 0 \leq x < x_0, \end{aligned}$$

$$\begin{aligned} n(x, 0) &= [x/\phi(x)] & x \geq x_0, \\ &= 0 & 0 \leq x < x_0. \end{aligned}$$

From (ii), (iii) it is seen that $\phi(x)$ and $x/\phi(x) \uparrow$ with x . Further

$$(26) \quad \int_0^X \frac{n(t, 0)}{t^2} dt < \int_{x_0}^X \frac{dt}{t\phi(t)} \rightarrow \frac{1}{\alpha(x_0)}$$

as $X \rightarrow \infty$. Hence (1) is convergent. Write $R = r\alpha(r)$. Then

$$\alpha(R) - \alpha(r) = \int_r^R \alpha'(t) dt = o(\alpha(R)).$$

Hence $\alpha(R) \sim \alpha(r)$ and so $\alpha(r)$ is "slowly increasing" function [6] and

$$\alpha(r) = o(r^\delta), \quad \delta > 0, \quad \int_{x_0}^r \frac{dt}{\alpha(t)} \sim \frac{r}{\alpha(r)}.$$

From (ii) we have $\alpha(r) = o(\phi(r))$. Let

$$f(z) = \prod_1^\infty \left(1 + \frac{z}{r_n}\right)$$

where $r_n > 0$, $n = 1, 2, \dots$, and are given by $n(x, 0)$. (The first few zeros may be equal; cf. [8, p. 228].) From (26) we see that the series $\sum 1/r_n$ is convergent and $f(z)$ is an entire function (c.p.) of genus $p = 0$. Now

$$\log M(r) < r \int_0^r \frac{n(t, 0)}{t} dt + r \int_r^\infty \frac{n(t, 0)}{t^2} dt \leq \{1 + o(1)\} \frac{r}{\alpha(r)},$$

$$\begin{aligned} \log M(r) &> r \int_R^\infty \frac{n(t, 0) dt}{t(t+r)} \\ &> r \int_R^\infty \left\{ \frac{t}{\phi(t)} - 1 \right\} \frac{dt}{t(t+r)} = \frac{r}{\alpha(r)} \{1 + o(1)\}. \end{aligned}$$

Hence

$$\begin{aligned} \log M(r) &\sim r/\alpha(r), \quad \log M(R) \sim r, \quad n(r, 0)\phi(r) \sim r \sim N(r, 0)\phi(r), \\ T(R) &\geq \frac{\beta - 1}{\beta + 1} \log M\left(\frac{R}{\beta}\right), & \beta > 1, \\ (27) \quad &\sim \frac{\beta - 1}{\beta + 1} \frac{R}{\beta\alpha(r)} = \frac{\beta - 1}{\beta + 1} \frac{r}{\beta}. \end{aligned}$$

Choose $\beta = 1 + 2^{1/2}$ and (25) follows when $\rho = 1$.

(b) If $\rho > 1$ then let

$$f(z) = \prod_1^{\infty} \left\{ 1 + \frac{z^{\rho}}{r_n} \right\}$$

where r_n have been defined in (a). $f(z)$ is an entire function (c.p.) of order ρ and genus $\rho - 1$. We have for all $r > r_0$

$$\log M(kr, f) < c_3 n(r, f - 0) \phi(r) / \alpha(r),$$

$$\log M(R, f) \geq \frac{1}{\rho} (1 + o(1)) (\alpha(r))^{\rho-1} n(r, f - 0) \phi(r),$$

$$T(R, f) > c_4 (\alpha(r))^{\rho-1} n(r, f - 0) \phi(r),$$

$$T(R, f) > c_4 (\alpha(r))^{\rho-1} N(r, 1/f) \phi(r)$$

and (25) is proved for $\rho > 1$.

REFERENCES

1. R. P. Boas, *Rend. Circ. Mat. Palermo* vol. 1 (1952) pp. 1-9.
2. ———, *J. London Math. Soc.* vol. 28 (1953) pp. 194-196.
3. ———, *Entire functions*, New York, Academic Press, 1954.
4. A. Edrei and W. H. J. Fuchs, Abstract 546-63, *Notices Amer. Math. Soc.* vol. 6 (1958) p. 371.
5. W. K. Hayman, *Rend. Circ. Mat. Palermo* vol. 2 (1953) pp. 346-392.
6. G. H. Hardy and W. W. Rogosinski, *Quart. J. Math. Oxford Ser. (2)* vol. 16 (1945) pp. 49-58.
7. R. Nevanlinna, *Le theoreme de Picard-Borel et la theorie des fonctions meromorphes*, Paris, 1929.
8. ———, *Eindeutige Analytische Funktionen*, Berlin, 1953.
9. S. M. Shah, *J. London Math. Soc.* vol. 15 (1940) pp. 23-31.
10. ———, *J. Indian Math. Soc.* vol. 5 (1941) pp. 179-188.
11. ———, *Math. Student* vol. 12 (1944) pp. 67-70.
12. ———, *Proc. Amer. Math. Soc.* vol. 2 (1951) pp. 694-698.
13. ———, *Duke Math. J.* vol. 19 (1952) pp. 585-593.
14. ———, *J. Indian. Math. Soc.* vol. 20 (1956) pp. 315-327.

UNIVERSITY OF WISCONSIN AND

MUSLIM UNIVERSITY, ALIGARH, INDIA