ON THE ABSOLUTE HARMONIC SUMMABILITY OF A SERIES RELATED TO A FOURIER SERIES

O. P. VARSHNEY

1.1. Let $\sum a_n$ be a given infinite series with the sequence of partial sums $\{s_n\}$. Let the sequence $\{t_n\}$ be defined by

$$t_n = \frac{(n+1)^{-1}s_0 + n^{-1}s_1 + \cdots + 1 \cdot s_n}{P_n},$$

$$\left(P_n = 1 + \frac{1}{2} + \cdots + \frac{1}{n+1}\right).$$

The series $\sum a_n$ is defined to be summable by harmonic means if the sequence $\{t_n\}$ tends to a limit as $n \rightarrow \infty$ [4]. If the series $\sum_n |t_n - t_{n-1}|$ is convergent, we say that the series is absolutely harmonic summable. It is known that the method of summability is absolutely regular and implies absolute Cesàro summability of every positive order [2].

1.2. Let $f(t)$ be a periodic function, with period 2π , and integrable (L) over $(-\pi, \pi)$. We assume without any loss of generality that the Fourier series of $f(t)$ is given by

$$(1.21) \quad \sum_1^{\infty} (a_n \cos nt + b_n \sin nt) = \sum_1^{\infty} A_n(t),$$

and that $\int_{-\pi}^{\pi} f(t) dt = 0$. We write

$$\phi(t) = \frac{1}{2} \{f(x+t) + f(x-t)\}.$$

Mohanty [3] has considered the absolute Riesz summability of the series

$$(1.22) \quad \sum_{n=1}^{\infty} A_n(t) / \log(n+1).$$

2. In this paper we establish the following theorem:

THEOREM. *If $\phi(t)$ is of bounded variation in $(0, \pi)$ then the series (1.22) is absolutely summable by harmonic means.*

We require the following lemmas for the proof of our theorem:

Received by the editors December 10, 1958.

LEMMA 1 [5]. Uniformly for $0 < t < \pi$

$$\left| \sum_m^n \frac{\sin \nu t}{\nu} \right| \leq K$$

where m and n are any positive integers.

LEMMA 2 [1]. If $0 < t < \pi$, then

$$\left| \sum_{k=0}^m \frac{\cos(k+1)t}{k+1} \right| = O\left(1 + \log \frac{1}{t}\right).$$

With the help of Lemmas 1 and 2, we may easily deduce

LEMMA 3. If $0 < t < \pi$, then for all positive integers m and m'

$$\sum_{k=m}^{m'} \frac{\sin(n-k)t}{k+1} = O\left(1 + \log \frac{1}{t}\right).$$

LEMMA 4. If $P_n = 1 + 1/2 + \dots + 1/(n+1)$, then

$$\begin{aligned} \text{(i)} \quad & \sum_{k=0}^{[n/2]-2} \left| \Delta \left(\frac{P_n(n+1) - P_k(k+1)}{(n-k) \log(n-k+1)} \right) \right| = O(1); \\ \text{(ii)} \quad & \sum_{k=[n/2]}^{n-2} \left| \Delta \left(\frac{P_k}{k+1} \cdot \frac{1}{\log(n-k+1)} \right) \right| = O\left(\frac{P_n}{n}\right). \end{aligned}$$

For proving (i) we observe that

$$\begin{aligned} & \sum_{k=0}^{[n/2]-2} \left| \Delta \left(\frac{P_n(n+1) - P_k(k+1)}{(n-k) \log(n-k+1)} \right) \right| \\ & \leq \sum_{k=0}^{[n/2]-2} \frac{P_n(n+1) - P_k(k+1)}{(n-k)(n-k-1) \log(n-k+1)} \\ & \quad + \sum_{k=0}^{[n/2]-2} \frac{P_{k+1}(k+2) - P_k(k+1)}{(n-k) \log(n-k+1)} \\ & \quad + O\left(\sum_{k=0}^{[n/2]-2} \frac{P_n(n+1) - P_k(k+1)}{(n-k)^2 \log^2(n-k+1)} \right) \\ & = O\left(\sum_{k=0}^{[n/2]-2} \frac{1}{n-k} \right) + O\left[\sum_{k=0}^{[n/2]-2} \frac{P_{k+1}}{(n-k) \log(n-k+1)} \right] \\ & \quad + O\left[\sum_{k=0}^{[n/2]-2} \frac{1}{(n-k) \log(n-k+1)} \right] \\ & = O\left(\sum_{k=0}^{[n/2]-2} \frac{1}{(n-k)} \right) = O(1). \end{aligned}$$

Again

$$\begin{aligned}
 & \sum_{k=[n/2]}^{n-2} \left| \Delta \left(\frac{P_k}{k+1} \cdot \frac{1}{\log(n-k+1)} \right) \right| \\
 &= O \left(\sum_{k=[n/2]}^{n-2} \frac{P_k}{k+1} \cdot \frac{1}{(n-k) \log^2(n-k+1)} \right) \\
 &\quad + O \left(\sum_{k=[n/2]}^{n-2} \frac{P_k}{(k+1)^2 \log(n-k+1)} \right) \\
 &= O(P_n/n) \left(\sum_{k=[n/2]}^{n-2} \frac{1}{(n-k) \log^2(n-k+1)} \right) + O\left(\frac{P_n}{n^2}\right) \cdot (n) \\
 &= O(P_n/n).
 \end{aligned}$$

This proves the lemma completely.

3. Proof of the theorem. Since

$$t_n = \frac{P_n u_0 + P_{n-1} u_1 + \cdots + P_0 u_n}{P_n}, \quad \left(u_n = \frac{A_n(t)}{\log(n+1)} \right),$$

we have

$$\begin{aligned}
 t_n - t_{n-1} &= \sum_{\nu=0}^{n-1} \left(\frac{P_\nu}{P_n} - \frac{P_{\nu-1}}{P_{n-1}} \right) u_{n-\nu} \\
 &= \frac{1}{P_n P_{n-1}} \sum_{\nu=0}^{n-1} \left(\frac{P_n}{\nu+1} - \frac{P_\nu}{n+1} \right) u_{n-\nu}.
 \end{aligned}$$

For the Fourier series of $f(t)$ at $t=k$,

$$A_n = \frac{2}{\pi} \int_0^\pi \phi(t) \cos ntdt$$

so that

$$t_n - t_{n-1} = \frac{2}{\pi} \int_0^\pi \phi(t) \left(\frac{1}{P_n P_{n-1}} \sum_{k=0}^{n-1} \left(\frac{P_n}{k+1} - \frac{P_k}{n+1} \right) \frac{\cos(n-k)t}{\log(n-k+1)} \right) dt.$$

Thus in order to prove the theorem, we have to establish that

$$\sum_n \left| \int_0^\pi \phi(t) g(n, t) dt \right| < \infty,$$

where

$$g(n, t) = \frac{1}{P_n P_{n-1}} \sum_{k=0}^{n-1} \left(\frac{P_n}{k+1} - \frac{P_k}{n+1} \right) \frac{\cos(n-k)t}{\log(n-k+1)}.$$

We observe that

$$\int_0^\pi \phi(t)g(n, t)dt = - \int_0^\pi \left(\int_0^t g(n, u)du \right) d\phi(t),$$

and

$$\sum_n \left| \int_0^\pi \left(\int_0^t g(n, u)du \right) d\phi(t) \right| \leq \int_0^\pi |d\phi(t)| \left\{ \sum_n \left| \int_0^t g(n, u)du \right| \right\}.$$

Since, by hypothesis, $\int_0^\pi |d\phi(t)| < \infty$, it suffices for our purpose to show that, uniformly for $0 < t < \pi$,

$$\sum_n \left| \int_0^t g(n, u)du \right| < \infty,$$

or, what is the same thing,

$$\Sigma \equiv \sum_n \left| \frac{1}{P_n P_{n-1}} \sum_{k=0}^{n-1} \left(\frac{P_n}{k+1} - \frac{P_k}{n+1} \right) \frac{\sin(n-k)t}{(n-k) \log(n-k+1)} \right| < \infty.$$

Denoting $\tau = [1/t]$, we have

$$\begin{aligned} \Sigma &\leq \sum_1^\tau \frac{1}{P_n P_{n-1}} \left| \sum_{k=0}^{n-1} \left(\frac{P_n}{k+1} - \frac{P_k}{n+1} \right) \frac{\sin(n-k)t}{(n-k) \log(n-k+1)} \right| \\ &\quad + \sum_{\tau+1}^\infty \frac{1}{P_n P_{n-1}} \left| \sum_{k=0}^{[n/2]-1} \left(\frac{P_n}{k+1} - \frac{P_k}{n+1} \right) \frac{\sin(n-k)t}{(n-k) \log(n-k+1)} \right| \\ &\quad + \sum_{\tau+1}^\infty \frac{1}{P_n P_{n-1}} \left| \sum_{[n/2]}^{n-1} \left(\frac{P_n}{k+1} - \frac{P_k}{n+1} \right) \frac{\sin(n-k)t}{(n-k) \log(n-k+1)} \right|. \end{aligned}$$

Now since $|\sin(n-k)t| \leq (n-k)t$ and $P_n(n+1) \geq P_k(k+1)$ for $k \leq n$, we have

$$\begin{aligned} \Sigma &= \sum_1^\tau \frac{1}{P_n P_{n-1}} \left| \sum_{k=0}^{n-1} \left(\frac{P_n}{k+1} - \frac{P_k}{n+1} \right) \frac{\sin(n-k)t}{(n-k) \log(n-k+1)} \right| \\ &\leq At \sum_1^\tau \frac{1}{P_n P_{n-1}} \sum_{k=0}^{n-1} P_n/k + 1 \\ (3.1) \quad &= At \sum_1^\tau P_n P_{n-1} / P_n P_{n-1} \\ &= O(1). \end{aligned}$$

By Abel's transformation and taking τ to be a fixed number $> \pi$,

$$\begin{aligned}
\sum_2 &= \sum_{\tau+1}^{\infty} \frac{1}{P_n P_{n-1}} \left| \sum_{k=0}^{[n/2]-1} \left(\frac{P_n}{k+1} - \frac{P_k}{n+1} \right) \frac{\sin(n-k)t}{(n-k) \log(n-k+1)} \right| \\
&= \sum_{\tau+1}^{\infty} \frac{1/(n+1)}{P_n P_{n-1}} \left| \sum_{k=0}^{[n/2]-1} \frac{P_n(n+1) - P_k(k+1)}{(n-k) \log(n-k+1)} \cdot \frac{\sin(n-k)t}{k+1} \right| \\
(3.2) \quad &= O \left[\sum_{\tau+1}^{\infty} \frac{1/(n+1)}{P_n P_{n-1}} \left(\log \frac{r}{t} \right) \frac{P_n(n+1) - P_{[n/2]-1} \left(\left\lceil \frac{n}{2} \right\rceil \right)}{\left(\left\lceil \frac{n}{2} \right\rceil + 1 \right) \log \left(\left\lceil \frac{n}{2} \right\rceil + 2 \right)} \right] \\
&\quad + O \left[\sum_{\tau+1}^{\infty} \frac{1/(n+1)}{P_n P_{n-1}} \left(\log \frac{r}{t} \right) \sum_{k=0}^{[n/2]-2} \left| \Delta \frac{P_n(n+1) - P_k(k+1)}{(n-k) \log(n-k+1)} \right| \right] \\
&= O \left(\log \frac{r}{t} \right) \sum_{\tau+1}^{\infty} \frac{1/(n+1)}{P_n P_{n-1}} = O \left(\frac{\log r/t}{P_r} \right) \\
&= O(1),
\end{aligned}$$

by using Lemmas 3 and 4 (i).

Since $1/(n+1)(n-k) = 1/(k+1)(n-k) - 1/(n+1)(k+1)$, we have

$$\begin{aligned}
\sum_3 &= \sum_{\tau+1}^{\infty} \frac{1}{P_n P_{n-1}} \left| \sum_{k=[n/2]}^{n-1} \left(\frac{P_n}{k+1} - \frac{P_k}{n+1} \right) \frac{\sin(n-k)t}{(n-k) \log(n-k+1)} \right| \\
&\leq \sum_{\tau+1}^{\infty} \frac{1}{P_n P_{n-1}} \left| \sum_{k=[n/2]}^{n-1} \frac{P_n - P_k}{(k+1)} \cdot \frac{\sin(n-k)t}{(n-k) \log(n-k+1)} \right| \\
&\quad + \sum_{\tau+1}^{\infty} \frac{1/(n+1)}{P_n P_{n-1}} \left| \sum_{k=[n/2]}^{n-1} \frac{P_k}{k+1} \cdot \frac{\sin(n-k)t}{\log(n-k+1)} \right| \\
&= \sum_{31} + \sum_{32}, \text{ say.}
\end{aligned}$$

Now since for $k \geq [n/2]$, $P_n - P_k = O(1)$, we obtain

$$\begin{aligned}
\sum_{31} &= \sum_{\tau+1}^{\infty} \frac{1}{P_n P_{n-1}} \left| \sum_{k=[n/2]}^{n-1} \frac{(P_n - P_k)}{k+1} \cdot \frac{\sin(n-k)t}{(n-k) \log(n-k+1)} \right| \\
(3.3) \quad &= O \left[\sum_{\tau+1}^{\infty} \frac{1}{P_n P_{n-1}} \left| \sum_{k=[n/2]}^{n-1} \frac{P_n - P_k}{(k+1)} \cdot \frac{1}{(n-k) \log(n-k+1)} \right| \right] \\
&= O \left(\sum_{\tau+1}^{\infty} \frac{1}{n P_n P_{n-1}} \sum_{k=[n/2]}^{n-1} \frac{1}{(n-k) \log(n-k+1)} \right) \\
&= O \left(\sum_{\tau+1}^{\infty} \frac{\log \log n}{n \log^2 n} \right) = O(1).
\end{aligned}$$

Finally since $\sum_a^b \sin nt = O(1/t)$ for all a and b , we have by Abel's transformation

$$\begin{aligned}
 \sum_{32} &= \sum_{\tau+1}^{\infty} \frac{1/(n+1)}{P_n P_{n-1}} \left| \sum_{k=[n/2]}^{n-1} \frac{P_k}{k+1} \cdot \frac{\sin(n-k)t}{\log(n-k+1)} \right| \\
 &= O \left(\sum_{\tau+1}^{\infty} \frac{1/(n+1)}{P_n P_{n-1}} \sum_{k=[n/2]}^{n-2} \frac{1}{t} \left| \Delta \left(\frac{P_k}{k+1} \cdot \frac{1}{\log(n-k+1)} \right) \right| \right) \\
 &\quad + O \left(\frac{1}{t} \right) \left(\sum_{\tau+1}^{\infty} \frac{1/(n+1)}{P_n P_{n-1}} \cdot \frac{P_{n-1}}{n} \right) \\
 (3.4) \quad &= O \left(\frac{1}{t} \right) \left(\sum_{\tau+1}^{\infty} \frac{1}{n^2 \log n} \right) \\
 &= O \left(\frac{1}{t} \right) \left(\frac{1}{\tau \log \tau} \right) \\
 &= O(1),
 \end{aligned}$$

using Lemma 4 (ii).

Collecting (3.1), (3.2), (3.3), and (3.4), we see that the theorem is completely proved.

I am very much indebted to Professor Misra for his kind interest and advice in the preparation of this paper.

REFERENCES

1. G. H. Hardy and W. W. Rogosinski, *Notes on Fourier series*, IV. *Summability* (R_2), Proc. Cambridge Philos. Soc. vol. 43 (1947) pp. 10-25.
2. Leonard McFadden, *Absolute Nörlund summability*, Duke Math. J. vol. 9 (1942) pp. 168-207.
3. R. Mohanty, *On the absolute Riesz summability of a Fourier series and allied series*, Proc. London Math. Soc. ser. 2 vol. 52 (1951) pp. 295-320.
4. M. Riesz, *Sur l'équivalence de certaines méthodes de sommation*, Proc. London Math. Soc. vol. 22 (1924) pp. 412-419.
5. E. C. Titchmarsh, *The theory of functions*, Oxford, 1939, p. 440.

UNIVERSITY OF SAUGAR, SAGAR, INDIA