DIVISIBLE MODULES

EBEN MATLIS

Introduction. Let R be an integral domain with quotient field Q , and let A be a module over R . A is said to be a *divisible* R -module, if $rA = A$ for every $r \neq 0 \in R$. An element $x \in A$ is said to be a *torsion element* of A , if there exists $r \neq 0 \in R$ such that $rx = 0$. The set of torsion elements of A is a submodule of A called the *torsion submodule* of A , and we will consistently denote it by A_T . We will let $E(A)$ denote the *injective envelope* of A (see [3]); and $\text{hd}_R A$ will denote the *homological dimension* of A as an R -module.

We will study conditions, some necessary, some sufficient, for the torsion submodule of a divisible module to be a direct summand. These will be related to the condition that $\text{hd}_R Q = 1$, where Q is the quotient field of R . We will apply these conditions to show that, if R is a Noetherian integral domain in which prime ideals different from zero are maximal, and if D is a divisible module over R , then D is a homomorphic image of an injective R -module and D_T is a direct summand of D . The same conclusions hold, if we merely assume for an arbitrary integral domain that its quotient field is countably generated as a module over the ring.

1. The torsion submodule. It is easy to see that, if C is an injective module over an integral domain R , then its torsion submodule C_T is also an injective R -module, and therefore a direct summand. Namely, any homomorphism of an ideal of R into C_T can be extended to a homomorphism of R into C ; but this extension must in fact map R into C_T . The following theorem is a generalization of this fact.

THEOREM 1.1. *Let R be an integral domain and H a homomorphic image of an injective R -module. Then H_T is a direct summand of H .*

PROOF. The mapping of a free R -module F onto an injective R -module C can be extended to a mapping of $E(F)$ onto C . We can thus assume that there exists a torsion-free, divisible R -module U and an epimorphism $f: U \rightarrow H$. Now H/H_T , being torsion-free and divisible, is a direct sum of R -modules Q_i , where Q_i is isomorphic to Q the quotient field of R . Let S_i be the inverse image of Q_i under the canonical map $H \rightarrow H/H_T$. We will prove that H_T is a direct summand of each S_i , and by [2, Lemma 2] this will complete the proof of the theorem.

Presented to the Society, September 3, 1959; received by the editors May 23, 1959.

Let $y \in S_j - H_T$; then Ry is a torsion-free submodule of S_j . Choose $x \in U$ such that $f(x) = y$. Since U is a vector space over Q , there exists an R -submodule T_j of U such that $T_j \cong Q$ and $x \in T_j$. If $u \neq 0 \in T_j$, there exist $r, s \in R$ such that $ru = sx \neq 0$. Since $rf(u) = sf(x) = sy \neq 0$, we have $f(u) \neq 0$, and thus $f(T_j) \cong T_j \cong Q$.

Since $H = \sum_i S_i$, $f(u) = w + z$, where $w \in \sum_{i \neq j} S_i$ and $z \in S_j$. Hence $rw + rz = rf(u) = sy \in S_j$. Thus $rw \in \sum_{i \neq j} S_i \cap S_j = H_T$. Therefore, $w \in H_T \subset S_j$, and so $f(u) \in S_j$. This shows that $f(T_j) \subset S_j$. Since $f(T_j) \cong Q$, we have $f(T_j) \cap H_T = 0$ and $f(T_j)$ maps onto Q_j under the canonical map $H \rightarrow H/H_T$. Thus $S_j = H_T \oplus f(T_j)$; and so H_T is a direct summand of H .

THEOREM 1.2. *Let R be an integral domain with quotient field $Q \neq R$. Suppose that D_T is a direct summand of D for every divisible R -module D . Then $\text{hd}_R Q = 1$.*

PROOF. Let A be any R -module and let $E = E(A)$. Since E is an essential extension of A , E/A is a torsion R -module. Let G be any R -module extension of E/A by Q . Since both Q and E/A are divisible, G is also divisible. Clearly E/A is the torsion submodule of G ; and thus by assumption E/A is a direct summand of G . Thus $\text{Ext}_R^1(Q, E/A) = 0$ [1, Theorem 14.1.1]. We also have $\text{Ext}_R^n(Q, E) = 0$ for $n > 0$, since E is injective. Therefore, from the exact sequence:

$$\text{Ext}_R^1(Q, E/A) \rightarrow \text{Ext}_R^2(Q, A) \rightarrow \text{Ext}_R^2(Q, E)$$

we deduce that $\text{Ext}_R^2(Q, A) = 0$. Hence $\text{hd}_R Q \leq 1$. Since Q is not R -projective, $\text{hd}_R Q = 1$.

Over an arbitrary integral domain it is not true that D_T is a direct summand for every divisible module D . For by the above theorem this would imply that $\text{hd}_R Q = 1$. However, I. Kaplansky has shown (unpublished) that $\text{hd}_R Q = 1$ for a valuation ring R if and only if Q is a countably generated R -module.

THEOREM 1.3. *Let R be an integral domain with quotient field $Q \neq R$, and suppose that Q is countably generated as a R -module. Then every divisible R -module D is a homomorphic image of an injective R -module. Thus D_T is a direct summand of D , and $\text{hd}_R Q = 1$.*

PROOF. There exists a countable set of generators $\{q_n\}$ for Q over R , and elements $\{a_{n+1}\}$ of R such that $q_1 = 1$ and $a_{n+1}q_{n+1} = q_n$. Let D be a divisible R -module, and let $x \neq 0 \in D$. We define a mapping f from the generators $\{q_n\}$ to D . Let $f(1) = x$; then there exists $x_2 \in D$ such that $a_2 x_2 = x$, and we define $f(q_2) = x_2$. There exists $x_3 \in D$ such that $a_3 x_3 = x_2$, and we define $f(q_3) = x_3$. We continue in this way and

define f on all the generators $\{q_n\}$. It is easily verified that f induces an R -homomorphism from Q into D such that the image contains x . It is now clear that by taking a big enough direct sum G of copies of Q we can define an R -homomorphism of G onto D .

It should be remarked that if R is any integral domain and S a countable, multiplicatively closed subset of R , then it can be easily shown that if F is a countably generated free R -module and f a suitably chosen mapping of F onto R_S , then the kernel of f is free; and thus $\text{hd}_R R_S \leq 1$.

2. $\text{hd}_R Q = 1$.

PROPOSITION 2.1. *Let R be an integral domain with quotient field Q such that $\text{hd}_R Q = 1$. Let H be an R -module. Then the following statements are equivalent:*

- (1) $\text{Ext}_R^1(Q/R, H) = 0$.
- (2) Every R -homomorphism from R into H can be extended to an R -homomorphism from Q into H .
- (3) H is a homomorphic image of an injective R -module.

PROOF. That (1) implies (2) follows immediately from the exact sequence:

$$\text{Hom}_R(Q, H) \rightarrow \text{Hom}_R(R, H) \rightarrow \text{Ext}_R^1(Q/R, H).$$

That (2) implies (3) is trivial. That (3) implies (1) follows from the fact that $\text{hd}_R Q/R = 1$.

PROPOSITION 2.2. *Let R be an integral domain with quotient field Q such that $\text{hd}_R Q = 1$. Let H be an R -module. Then:*

- (1) If H is a homomorphic image of an injective R -module, so is H_T .
- (2) If B is a submodule of H and if B and H/B are homomorphic images of injective modules, then so is H .

PROOF.

(1) If H is a homomorphic image of an injective R -module, then H_T is a direct summand of H by Theorem 1.1. Hence H_T is also a homomorphic image of an injective R -module.

(2) Suppose that B and H/B are homomorphic images of injective R -modules. We have an exact sequence:

$$\text{Ext}_R^1(Q/R, B) \rightarrow \text{Ext}_R^1(Q/R, H) \rightarrow \text{Ext}_R^1(Q/R, H/B).$$

By Proposition 2.1 the two end modules are zero, and thus $\text{Ext}_R^1(Q/R, H) = 0$. Hence by Proposition 2.1 again, H is a homomorphic image of an injective R -module.

DEFINITION. Let B be a module over an integral domain. Then we will say that B is *h-reduced*, if B has no nonzero submodules which are homomorphic images of injective modules.

COROLLARY 2.3. *Let A be a module over an integral domain R with quotient field Q such that $\text{hd}_R Q = 1$. Then A has a unique largest submodule H which is a homomorphic image of an injective R -module, and A/H is *h-reduced*.*

PROOF. Let H be the sum of all submodules of A which are homomorphic images of injective R -modules. It is clear that H is the unique largest submodule of A which is a homomorphic image of an injective R -module. Suppose that B/H is a homomorphic image of an injective R -module, where B is a submodule of A containing H . Then by Proposition 2.2 B is a homomorphic image of an injective R -module. Therefore, $B = H$ and $B/H = 0$.

PROPOSITION 2.4. *Let R be an integral domain with quotient field Q such that $\text{hd}_R Q = 1$. Let D be a divisible module over R , and let H be a submodule of D_T such that H is a homomorphic image of an injective R -module. Then D_T is a direct summand of D if and only if D_T/H is a direct summand of D/H .*

PROOF. Suppose that D_T is a direct summand of D , and let S be a complementary summand of D_T in D . Then $D/H \cong D_T/H \oplus S$, and since D_T/H is the torsion submodule of D/H , D_T/H is a direct summand of D/H . Conversely, suppose that $D/H = D_T/H \oplus G/H$, where G is a submodule of D containing H . Now G/H is torsion-free and divisible, hence injective. Thus by Proposition 2.2 and Theorem 1.1 H is a direct summand of G . Let L be a complementary summand of H in G . Then it is clear that $D = D_T \oplus L$.

PROPOSITION 2.5. *Let R be an integral domain with quotient field Q , and let T be an *h-reduced* torsion R -module. Then $\text{Ext}_R^1(Q, T) = 0$ if and only if $T \cong \text{Ext}_R^1(Q/R, T)$.*

PROOF. Since $\text{Hom}_R(Q, T) = 0$, we have an exact sequence:

$$0 \rightarrow \text{Hom}_R(R, T) \rightarrow \text{Ext}_R^1(Q/R, T) \rightarrow \text{Ext}_R^1(Q, T) \rightarrow 0.$$

It follows that if $\text{Ext}_R^1(Q, T) = 0$, then $T \cong \text{Ext}_R^1(Q/R, T)$. Conversely, if $T \cong \text{Ext}_R^1(Q/R, T)$, then the above exact sequence shows that $\text{Ext}_R^1(Q, T)$ is a torsion module. However, $\text{Ext}_R^1(Q, T)$ is torsion-free, and thus $\text{Ext}_R^1(Q, T) = 0$.

COROLLARY 2.6. *Let R be an integral domain with quotient field Q .*

Then the torsion submodule of a divisible R -module is always a direct summand if and only if the following two conditions hold:

- (1) $\text{hd}_R Q = 1$.
- (2) $T \cong \text{Ext}_R^1(Q/R, T)$, whenever T is an h -reduced, torsion, divisible R -module.

PROOF. The necessity follows from Theorem 1.2 and Proposition 2.5; the sufficiency follows from Corollary 2.3 and Propositions 2.4 and 2.5.

PROPOSITION 2.7. Let R be an integral domain with quotient field Q such that $\text{hd}_R Q = 1$ and $\text{gl. dim. } R \leq 2$. Let S be any torsion-free R -module. Then $\text{hd}_R S \leq 1$. Thus if A is an R -module such that A_T is a homomorphic image of an injective R -module, then A_T is a direct summand of A .

PROOF. Let B be any R -module. Then from the exact sequence:

$$0 \rightarrow S \rightarrow Q \otimes_R S \rightarrow Q/R \otimes_R S \rightarrow 0$$

we derive the exact sequence:

$$\text{Ext}_R^2(Q \otimes_R S, B) \rightarrow \text{Ext}_R^2(S, B) \rightarrow \text{Ext}_R^3(Q/R \otimes_R S, B).$$

Since $\text{hd}_R Q = 1$ and $\text{gl. dim. } R \leq 2$, the two end modules are zero. Thus $\text{hd}_R S \leq 1$, and the rest of the theorem follows immediately.

2. Krull dimension = 1.

Throughout this section R will be a Noetherian integral domain with the property that nonzero prime ideals are maximal. We will let Q be the quotient field of R and $K = Q/R$.

DEFINITION. Let A be an R -module and M a prime ideal of R . We will say that A is M -primary, if for any $x \neq 0 \in A$, the order ideal of x is an M -primary ideal. If B is any R -module, and A is the set of all elements of B whose order ideal is M -primary (together with the element 0), then A is an M -primary R -module which we will call the M -primary component of B .

LEMMA 3.1. Let B be any torsion R -module. Then B is the direct sum of its M -primary components, M ranging over the prime ideals of R . Furthermore, $B \otimes_R R_M$ is the M -primary component of B .

PROOF. Let $\{M_a\}$ be the collection of nonzero prime ideals of R . By [3, Theorem 3.3] $E(B) = \sum_a \oplus E_a$, where E_a is the M_a -component of $E(B)$. Let $B_a = B \cap E_a$; then B_a is the M_a -component of B . Let $x \neq 0 \in B$; then $x = x_1 + \cdots + x_n$, where $x_i \in E_i$. Now $\bigcap_{i=2}^n M_i \not\subset M_1$;

hence there exists $s \in \bigcap_{i=1}^n M_i$ such that $s \in M_1$. Then there exists an integer $k > 0$ such that $s^k x_i = 0$ for $i = 2, \dots, n$. Hence $s^k x = s^k x_1$. There are elements $m \in M$ and $t \in R$ such that $1 = m + ts^k$. There is an integer $q > 0$ such that $m^q x_1 = 0$. Since $1 = m^q + rs^k$, $r \in R$, we have $x_1 = rs^k x_1 = rs^k x \in B_1$. Similarly $x_i \in B_i$ for $i = 2, \dots, n$. Thus $B = \sum_a B_a$.

Let M_v be a prime ideal of R . Clearly $B_a \otimes_R R_{M_v} = 0$, if $M_a \neq M_v$. Thus $B \otimes_R R_{M_v} = B_v \otimes_R R_{M_v}$. It is easily seen that the canonical map $B_v \rightarrow B_v \otimes_R R_{M_v}$ is an epimorphism. However, since B_v is M_v -primary, the kernel of this map is zero. Thus $B_v = B_v \otimes_R R_{M_v}$, and so $B \otimes_R R_{M_v} = B_v$.

LEMMA 3.2. $\text{hd}_R Q = 1$.

PROOF. It is sufficient to prove that $\text{hd}_R K = 1$. By Lemma 3.1 $K = \sum_a \oplus K_{M_a}$; thus it is sufficient to prove that $\text{hd}_R K_{M_v} = 1$. For this it is sufficient to prove that if D is any divisible R -module, then $\text{Ext}_R^1(K_{M_v}, D) = 0$. Let A be any extension of D by K_{M_v} ; then A is a divisible R -module. We have $D = \sum_a \oplus D_{M_a}$ and $A = \sum_a \oplus A_{M_a}$. Clearly $D_{M_a} = D \cap A_{M_a}$. Hence for $a \neq v$, we have $D_{M_a} = A_{M_a}$. Thus we have an exact sequence:

$$0 \rightarrow D_{M_v} \rightarrow A_{M_v} \rightarrow K_{M_v} \rightarrow 0,$$

and all of the modules and mappings of this sequence are R_{M_v} -modules and mappings. Thus we can assume that R is a local ring with a single nonzero prime ideal M .

Take $s \neq 0 \in M$, and let S be the multiplicatively closed set consisting of the powers of s . Now $R_S \subset Q$; on the other hand, the prime ideals of R_S and the prime ideals of R not meeting S are in 1-1 correspondence. Thus R_S is a field, and $R_S = Q$. Therefore, Q is a countably generated R -module; and thus $\text{hd}_R Q = 1$ by Theorem 1.3, or the remark following it.

THEOREM 3.3. *Every divisible R -module D is a homomorphic image of an injective R -module; and thus D_T is a direct summand of D .*

PROOF. Since $\text{hd}_R Q = 1$ by Lemma 3.2, it follows from Proposition 2.2 that we only need to prove that D_T is a homomorphic image of an injective R -module. By Lemma 3.1 $D_T = \sum_a \oplus D_{T_a}$, where $D_{T_a} = D_T \otimes_R R_{M_a}$ is a divisible R_{M_a} -module. As we have seen in Lemma 3.2, Q is a countably generated R_{M_a} -module; and thus by Theorem 1.3, D_{T_a} is a homomorphic image of an injective R_{M_a} -module. Thus D_{T_a} is a homomorphic image of an injective R -module; and therefore, the same is true of D_T .

REFERENCES

1. H. Cartan and S. Eilenberg, *Homological algebra*, Princeton University Press, 1956.
2. I. Kaplansky, *Modules over Dedekind rings and valuation rings*, Trans. Amer. Math. Soc. vol. 72 (1952) pp. 327-340.
3. E. Matlis, *Injective modules over Noetherian rings*, Pacific J. Math. vol. 8 (1958) pp. 511-528.

NORTHWESTERN UNIVERSITY

A CHARACTERIZATION OF ALGEBRAIC NUMBER FIELDS WITH CLASS NUMBER TWO¹

L. CARLITZ

Let $Z=R(\theta)$ denote an algebraic number field over the rationals with class number h . It is familiar that $h=1$ if and only if unique factorization into prime holds for the integers of Z . For fields with $h \leq 2$ we have the following criterion.

THEOREM. *The algebraic number field Z has class number ≤ 2 if and only if for every nonzero integer $\alpha \in Z$ the number of primes π_j in every factorization*

$$(1) \quad \alpha = \pi_1 \pi_2 \cdots \pi_k$$

depends only on α .

Suppose first that $h=2$ and consider the factorization into prime ideals

$$(2) \quad (\alpha) = \mathfrak{p}_1 \cdots \mathfrak{p}_s \mathfrak{r}_1 \cdots \mathfrak{r}_t,$$

where the $\mathfrak{p}_j$ are principal ideals while the $\mathfrak{r}_j$ are not. Then

$$\mathfrak{p}_j = (\pi_j) \quad (j = 1, \cdots, s).$$

Since $h=2$, it follows that

$$\mathfrak{r}_i \mathfrak{r}_j = (\rho_{ij}) \quad (i, j = 1, \cdots, t);$$

moreover t must be even, $=2u$, say. Thus every factorization into primes implied by (2), for example

Received by the editors August 3, 1959.

¹ Research sponsored by National Science Foundation grant NSF G-9425.