ZERO SETS FOR CONTINUOUS FUNCTIONS¹

R. C. BUCK

1. Introduction. Let D be a closed set in m -space, and let V be a linear space of real valued continuous functions defined on D . For any $f \in V$, let $Z(f)$ be its zero-set, the set of $p \in D$ with $f(p) = 0$. If $m = 1$, and V is a class of well behaved functions, then $Z(f)$ will be discrete except for f identically zero. This behavior depends strongly upon m . When $m \geq 2$, then—except in special cases—one expects $Z(f)$ to be a nowhere dense set which is locally of dimension $m - 1$.

In this note, we shall be concerned with certain distribution-properties of the zero-sets for functions of m variables which do not seem to have been explicitly stated before; as will be seen, they do not depend upon the nature of the functions in V but only upon the dimension of V . If $\dim(V) = N$, then there exists a system of N nonempty disjoint relatively open sets $D_1, D_2, \dots, D_N$ in D such that if f and g are in V , and if they agree at some point in each of the sets D_j , then $f = g$. Otherwise stated, if $f \in V$ and $Z(f)$ intersects each of the sets D_j , then $f \equiv 0$.

If D is the plane, and V the class of functions $f(x, y) = ax + by + c$, then the assertion is merely that there are three open sets D_j which cannot all be cut by a single straight line. Again, choose V as the space of polynomials in x, y, z of degree at most 2. If $f \in V$ but $f \neq 0$ then $Z(f)$ is a quadric surface in 3-space. Any set of nine points in 3-space lies in some set $Z(f)$; most sets of ten points do not. If we choose a set of points $p_1, \dots, p_{10}$ through which no quadric passes, and then surround each by a sufficiently small open sphere, we obtain ten open sets $D_1, \dots, D_{10}$ which form a system of uniqueness domains for the space V .

2. Uniqueness domains. Let V be any subspace of $C[D]$ of dimension N , and let $\phi_1, \phi_2, \dots, \phi_N$ be a basis for V .

THEOREM 1. *There exist disjoint nonempty sets $D_1, D_2, \dots, D_N$, relatively open in D , and a constant B such that for any $f \in V$, and $p \in D$, and any choice of $p_i \in D_i$*

$$(1) \quad |f(p)| \leq B \{ |f(p_1)| + |f(p_2)| + \dots + |f(p_N)| \}.$$

PROOF. Let S be the set of all points in N -space of the form $x = \langle \phi_1(p), \phi_2(p), \dots, \phi_N(p) \rangle$ for $p \in D$. If S were to lie in a proper

Received by the editors August 10, 1959 and, in revised form, September 23, 1959.

¹ This work was carried out under OOR Contract DA-11-022-ORD-2059.

subspace of N -space, there would exist constants $b_1, b_2, \dots, b_N$ such that $\sum b_j x_j = 0$ for all $x \in S$. This would imply that $\sum b_j \phi_j = 0$ so that the functions ϕ_j would not have been independent. We can therefore assume that S generates all of N -space. There then exist points $p_1, p_2, \dots, p_N$ in D such that the N points $\langle \phi_1(p_i), \phi_2(p_i), \dots, \phi_N(p_i) \rangle$, for $i=1, 2, \dots, N$, are linearly independent. Accordingly, the N by N matrix

$$\Delta = \begin{pmatrix} \phi_1(p_1) & \phi_2(p_1) & \dots & \phi_N(p_1) \\ \phi_1(p_2) & \phi_2(p_2) & \dots & \phi_N(p_2) \\ \vdots & \vdots & & \vdots \\ \phi_1(p_N) & \phi_2(p_N) & \dots & \phi_N(p_N) \end{pmatrix}$$

is nonsingular.

For any point $p \in D$, let L_p be the functional defined on $C[D]$ by $L_p(f) = f(p)$. The functionals L_{p_i} , for $i=1, 2, \dots, N$, are linearly independent; for, if $\sum c_i L_{p_i} = 0$, then for each $j=1, 2, \dots, N$, we would have

$$0 = \sum c_i L_{p_i}(\phi_j) = \sum c_i \phi_j(p_i)$$

and by the nonsingularity of Δ , $c_1 = c_2 = \dots = 0$. Since $\dim(V) = N$, its dual space V' is also of dimension N . Accordingly, $L_{p_1}, L_{p_2}, \dots, L_{p_N}$ constitutes a basis for V' . In particular then, if $f \in V$ and $L_{p_i}(f) = f(p_i) = 0$ for $i=1, 2, \dots, N$, then $L(f) = 0$ for every $L \in V'$, and $f \equiv 0$. This argument rests upon the nonsingularity of Δ ; since the functions $\phi_1, \phi_2, \dots, \phi_N$ are continuous on D , $d(p_1, p_2, \dots, p_N) = \det(\Delta)$ is a continuous function on $D^N = D \times D \times \dots \times D$. In particular, we can choose nonempty open sets $D_1, D_2, \dots, D_N$ in D so that $d(p_1, p_2, \dots, p_N) \neq 0$ for $p_i \in D_i$. As before, the functionals $L_{p_1}, L_{p_2}, \dots, L_{p_N}$ will form a basis for V' for any choices of the points p_i , with $p_1 \in D_1, p_2 \in D_2, \dots, p_N \in D_N$. Hence, if $f \in V$, and $Z(f)$ touches each of the sets D_i , $i=1, 2, \dots, N$, then $f \equiv 0$. This is part of the statement (1). To obtain the more complete formulation, we observe that the sets D_i can be taken so that $\inf |d(p_1, \dots, p_N)| = \delta > 0$, for all $p_i \in \bar{D}_i$, $i=1, 2, \dots, N$. Let $P = (p_1, p_2, \dots, p_N)$, and let $C_j(r, P)$ be the unique numbers defined by

$$\sum_{j=1}^N C_j(r, P) \phi_j(p_k) = \begin{cases} 1 & \text{if } k = r, \\ 0 & \text{if } k \neq r \end{cases}$$

for $r=1, 2, \dots, N$, and any P in $D_1 \times D_2 \times \dots \times D_N$. For each j and r , $C_j(r, P)$ is continuous in P , and since $|d(P)| \geq \delta$, there exists a constant B_0 , determined by the functions $\{\phi_j\}$ and the sets D_i , such that

$$|C_j(r, P)| \leq B_0 \quad \text{all } j, r, p_i \in D_i.$$

Setting $\psi_r = \sum_{j=1}^N C_j(r, P)\phi_j$, we obtain a new basis for V such that for any $f \in V$,

$$(2) \quad f = \sum_{r=1}^N f(p_r)\psi_r.$$

Moreover, for any r ,

$$\|\psi_r\| \leq \sum_1^N |C_j(r, P)| \|\phi_j\| \leq B_0 \sum_1^N \|\phi_j\| \leq B$$

where we have used $\|g\|$ to denote $\sup_{q \in D} |g(q)|$, for any $g \in C[D]$. Note that B again is determined only by the $\{\phi_j\}$ and the D_i . Returning to (2), we see that

$$\|f\| \leq B \sum_1^N |f(p_i)|$$

holding for any choice of $p_i \in D_i$, for $i=1, 2, \dots, N$.

The conclusion of Theorem 1 can be restated in an interesting way. Introduce a special functional M on $C[D]$:

$$M(f) = \sum_1^N \min_{p_i \in D_i} |f(p_i)|.$$

If we were to replace "min" by "max," this would define a seminorm on $C[D]$. On the finite dimensional subspace V , it would in fact be a norm, and would therefore have to define there the unique locally convex linear topology. As defined above, however, M is not a seminorm, for it is not true in general that $M(f+g) \leq M(f) + M(g)$. However, Theorem 1 implies that M still defines a Hausdorff topology on V ; if $M(f) = 0$, then f must take the value zero somewhere in each of the sets $\bar{D}_i$, and $f \equiv 0$. The essence of the more general formulation (1) is that this topology again coincides with the unique linear topology in V , and there must exist a constant B such that $\|f\| \leq BM(f)$, for all $f \in V$.

Another familiar instance of this general principle is the fact that in a finite dimensional function space, uniform convergence and L^1 (or L^p) convergence coincide. This is usually formulated as an inequality: there is a constant A such that

$$|f(p)| \leq A \int_D |f|$$

for all $p \in D$, and $f \in V$. This follows at once from (1). Let δ be the smallest of the measures $\mu(D_i)$. Then,

$$\int_D |f| \geq \sum_1^N \int_{D_i} |f| \geq \sum_1^N \mu(D_i) \min_{p_i \in D_i} |f(p_i)| \geq \delta M(f) \geq \frac{\delta}{B} \|f\|.$$

3. Generalizations. It is natural to ask if there are any other necessary restrictions upon the sets $\{D_i\}$ if they are to be uniqueness domains. This is answered by the next statement.

THEOREM 2. *If $D_1, D_2, \dots, D_N$ is any collection of open disjoint nonempty subsets of D , then there is a subspace V of dimension N in $C[D]$ such that if $f \in V$, and if $Z(f)$ touches each of the sets D_i , then $f \equiv 0$.*

Choose functions ϕ_i so that $\phi_i(p)$ is the distance from p to the closed complement of D_i . Then, ϕ_i vanishes off D_i , but is strictly positive on D_i . Let V be the set of functions $f = \sum c_i \phi_i$. If f takes the value 0 at some point of D_i , then it follows that $c_i = 0$; accordingly, if $Z(f)$ contains at least one point of each D_i , then $f \equiv 0$.

From this, by a method suggested by the referee, other examples can be constructed. Let θ be any continuous mapping of D onto another set D' , and let V' be an N dimensional subspace of $C[D']$. Suppose that $\{D'_1, \dots, D'_N\}$ is a collection of uniqueness domains in D' , and let $D_i = \theta^{-1}(D'_i)$. For any $\phi' \in V'$, define a function ϕ on D by $\phi(p) = \phi'(\theta p)$. Then, this yields a linear space $V \subset C[D]$ having $\{D_1, \dots, D_N\}$ for uniqueness domains. In particular, the sets D_i can be selected so that their union exhausts all but a nowhere dense subset of D , and this can be achieved with many different choices of V .

This is no longer possible in general if we wish to retain the more precise inequality (1) given in Theorem 1. Suppose that we have a space V of dimension N , and a collection of sets $\{D_i\}_1^N$ such that for any $f \in V$,

$$\|f\| \leq B \sum_1^N \inf_{p_i \in D_i} |f(p_i)|.$$

Suppose that there is a point q which is a common boundary point for all the sets D_i . Then $N = 1$. For, if $f \in V$ and $f(q) = 0$, then by the inequality and the continuity of f , $f \equiv 0$. Suppose ϕ_1 and ϕ_2 are two independent functions in V ; set $f = \phi_2(q)\phi_1 - \phi_1(q)\phi_2$ and have $f(q) = 0$.