A PROOF OF A FUNCTIONAL EQUATION RELATED TO THE THEORY OF PARTITIONS

SHÔ ISEKI

Introduction. The purpose of this note is to give a simple alternative proof for the following functional equation:

$$(1) \quad \sum_{l=0}^{\infty} \{ \lambda((l + \alpha)z - i\beta) + \lambda((l + 1 - \alpha)z + i\beta) \} + \pi z(\alpha^2 - \alpha + 1/6) \\ = \sum_{l=0}^{\infty} \{ \lambda((l + \beta)z^{-1} + i\alpha) + \lambda((l + 1 - \beta)z^{-1} - i\alpha) \} \\ + \pi z^{-1}(\beta^2 - \beta + 1/6) + 2\pi i(\alpha - 1/2)(\beta - 1/2),$$

where $\lambda(t) = -\log(1 - e^{-2\pi t})$ (the principal value), $0 \leq \alpha \leq 1$, $0 < \beta < 1$ (or $0 < \alpha < 1$, $0 \leq \beta \leq 1$), and $\Re(z) > 0$. This formula was first proved in [2], and its applications to the theory of partitions have also been investigated (see [2; 3]).

The method we employ in this paper is essentially that of Rademacher [4]. Although our method is still based on the Mellin transform technique and the theory of the Hurwitz zeta-functions, we have made a thorough revision of Rademacher's original method, and thus a very short and direct proof of the functional equation can be given.

It may be noted that if we define a function $\Lambda(z, \alpha, \beta)$ by

$$\Lambda(z, \alpha, \beta) = \sum_{l=0}^{\infty} \{ \lambda((l + \alpha)z - i\beta) + \lambda((l + 1 - \alpha)z + i\beta) \} \\ + \pi z(\alpha^2 - \alpha + 1/6) - \pi i(\alpha - 1/2)(\beta - 1/2),$$

then (1) is written in the form

$$\Lambda(z, \alpha, \beta) = \Lambda(z^{-1}, 1 - \beta, \alpha).$$

On the other hand, it is clear from the definition that

$$\Lambda(z, \alpha, \beta) = \Lambda(z, 1 - \alpha, 1 - \beta).$$

Proof of the functional equation. To prove (1), we first assume that $0 < \alpha < 1$, $0 < \beta < 1$, $\Re(z) > 0$. Using the expansion

$$\lambda(x) = -\log(1 - e^{-2\pi x}) = \sum_{n=1}^{\infty} n^{-1} e^{-2\pi x n}$$

and applying the Mellin formula

Received by the editors May 13, 1960 and, in revised form, June 21, 1960.

$$e^{-y} = (2\pi i)^{-1} \int_{c-i\infty}^{c+i\infty} \Gamma(s) y^{-s} ds \quad (\Re(y) > 0)$$

with $c=3/2$, it follows that

$$\begin{aligned} \sum_{l=0}^{\infty} \lambda((l+\alpha)z - i\beta) &= \sum_{l=0}^{\infty} \sum_{n=1}^{\infty} \frac{1}{n} e^{-2\pi((l+\alpha)z - i\beta)n} \\ &= \sum_{l=0}^{\infty} \sum_{n=1}^{\infty} \frac{e^{2\pi i\beta n}}{2\pi i n} \int_{3/2-i\infty}^{3/2+i\infty} \Gamma(s) \{2\pi(l+\alpha)zn\}^{-s} ds \\ &= \frac{1}{2\pi i} \int_{3/2-i\infty}^{3/2+i\infty} \frac{\Gamma(s)}{(2\pi z)^s} \left(\sum_{l=0}^{\infty} \frac{1}{(l+\alpha)^s} \right) \left(\sum_{n=1}^{\infty} \frac{e^{2\pi i\beta n}}{n^{1+s}} \right) ds. \end{aligned}$$

Therefore

$$(2) \quad \sum_{l=0}^{\infty} \lambda((l+\alpha)z - i\beta) = \frac{1}{2\pi i} \int_{3/2-i\infty}^{3/2+i\infty} \frac{\Gamma(s)}{(2\pi z)^s} \zeta(s, \alpha) \zeta_{\beta}(1+s) ds,$$

where $\zeta(s, \alpha) = \sum_{l=0}^{\infty} (l+\alpha)^{-s}$ is the Hurwitz zeta-function, and $\zeta_{\beta}(s) = \sum_{n=1}^{\infty} n^{-s} e^{2\pi i\beta n}$ ($\Re(s) > 0$).

Now the function $\zeta_a(s)$ ($0 < a < 1$) can be expressed in terms of Hurwitz zeta-functions, i.e.,

$$(3) \quad \zeta_a(s) = \Gamma(1-s)(2\pi)^{s-1} \left\{ e^{\pi i(1-s)/2} \zeta(1-s, a) + e^{\pi i(s-1)/2} \zeta(1-s, 1-a) \right\}.$$

This relation is an easy consequence of the famous formula:

$$\begin{aligned} \zeta(s, a) &= \frac{2\Gamma(1-s)}{(2\pi)^{1-s}} \left\{ \sin(\pi s/2) \sum_{n=1}^{\infty} \frac{\cos(2\pi a n)}{n^{1-s}} \right. \\ &\quad \left. + \cos(\pi s/2) \sum_{n=1}^{\infty} \frac{\sin(2\pi a n)}{n^{1-s}} \right\} \quad (\Re(s) < 0). \end{aligned}$$

From (3) we see that $\zeta_a(s)$ is an integral function of s (cf. Apostol [1]).

Returning to (2), we obtain further that

$$\begin{aligned} (4) \quad \sum_{l=0}^{\infty} \{ \lambda((l+\alpha)z - i\beta) + \lambda((l+1-\alpha)z + i\beta) \} \\ = \frac{1}{2\pi i} \int_{3/2-i\infty}^{3/2+i\infty} z^{-s} F(s; \alpha, \beta) ds \end{aligned}$$

with

$$(5) \quad F(s; \alpha, \beta) = \Gamma(s)(2\pi)^{-s} \{ \zeta(s, \alpha) \zeta_\beta(1+s) + \zeta(s, 1-\alpha) \zeta_{1-\beta}(1+s) \}.$$

The function $F(s; \alpha, \beta)$ satisfies the equation

$$(6) \quad F(s; \alpha, \beta) = F(-s; 1-\beta, \alpha),$$

since we have, by (5) and (3),

$$(7) \quad \begin{aligned} F(s; \alpha, \beta) &= \Gamma(s)\Gamma(-s) [e^{-\pi i s/2} \{ \zeta(s, \alpha) \zeta(-s, \beta) + \zeta(s, 1-\alpha) \zeta(-s, 1-\beta) \} \\ &\quad + e^{\pi i s/2} \{ \zeta(s, \alpha) \zeta(-s, 1-\beta) + \zeta(s, 1-\alpha) \zeta(-s, \beta) \}], \end{aligned}$$

which is unchanged when s, α, β are replaced by $-s, 1-\beta, \alpha$ respectively.

Now in the integral on the right of (4), we move the path of integration from $\Re(s)=3/2$ to $\Re(s)=-3/2$, and obtain

$$(8) \quad \begin{aligned} \frac{1}{2\pi i} \int_{3/2-i\infty}^{3/2+i\infty} z^{-s} F(s; \alpha, \beta) ds \\ = \frac{1}{2\pi i} \int_{-3/2-i\infty}^{-3/2+i\infty} z^{-s} F(s; \alpha, \beta) ds + R_1 + R_0 + R_{-1}, \end{aligned}$$

where R_1, R_0, R_{-1} are the corresponding residues of the integrand at its simple poles $s=1, 0, -1$ respectively. The displacement of the path of integration may easily be justified by considering the order of magnitude of the integrand. Actually, we have, writing $\Im(s)=t$,

$$\begin{aligned} z^{-s} &= O(e^{(\pi/2-\delta)|t|}) \quad \text{for } |\arg z| \leq \pi/2 - \delta < \pi/2, \\ \Gamma(s)\Gamma(-s) &= -\pi(s \sin \pi s)^{-1} = O(|t|^{-1} e^{-\pi|t|}), \\ \zeta(s, \alpha) &= O(|t|^\epsilon) \quad (\text{see [5, p. 276]}), \end{aligned}$$

and hence, by (7),

$$z^{-s} F(s; \alpha, \beta) = O(|t|^{2\epsilon-1} e^{-\delta|t|}).$$

The values of the residues are calculated as follows:

$$\begin{aligned} R_1 &= (2\pi z)^{-1} \{ \zeta_\beta(2) + \zeta_{1-\beta}(2) \} \\ &= (2\pi z)^{-1} \sum_{n=1}^{\infty} n^{-2} (e^{2\pi i \beta n} + e^{-2\pi i \beta n}) \\ &= (\pi z)^{-1} \sum_{n=1}^{\infty} n^{-2} \cos(2\pi \beta n) = \pi z^{-1} (\beta^2 - \beta + 1/6), \end{aligned}$$

$$\begin{aligned}
 R_0 &= \zeta(0, \alpha)\zeta_\beta(1) + \zeta(0, 1 - \alpha)\zeta_{1-\beta}(1) \\
 &= (1/2 - \alpha) \sum_{n=1}^{\infty} n^{-1} (e^{2\pi i \beta n} - e^{-2\pi i \beta n}) \\
 &= 2i(1/2 - \alpha) \sum_{n=1}^{\infty} n^{-1} \sin(2\pi \beta n) \\
 &= 2\pi i(1/2 - \alpha)(1/2 - \beta),
 \end{aligned}$$

and, using (6),

$$R_{-1} = -\pi z(\alpha^2 - \alpha + 1/6).$$

Inserting these values and the relation (6) into the right member of (8), changing the variable s to $-s$ and using (4), it is found that the required equation (1) holds for $0 < \alpha < 1$, $0 < \beta < 1$, and $\Re(z) > 0$. The validity of (1) for the end-points of the interval of α or β may then be established by means of letting $\alpha \rightarrow +0$, $1-0$ or $\beta \rightarrow +0$, $1-0$ in (1). This completes the proof of the functional equation.

BIBLIOGRAPHY

1. T. M. Apostol, *On the Lerch zeta function*, Pacific J. Math. vol. 1 (1951) pp. 161-167.
2. S. Iseki, *The transformation formula for the Dedekind modular function and related functional equations*, Duke Math. J. vol. 24 (1957) pp. 653-662.
3. ———, *A partition function with some congruence condition*, Amer. J. Math. vol. 81 (1959) pp. 939-961.
4. H. Rademacher, *Zur Theorie der Modulfunktionen*, J. Reine Angew. Math. vol. 167 (1932) pp. 312-336.
5. E. T. Whittaker and G. N. Watson, *A course of modern analysis*, Cambridge, University Press, 1945.

DEFENSE ACADEMY, YOKOSUKA, JAPAN