A NOTE ON MATRIX RICCATI SYSTEMS

W. J. COLES

1. Let $A(t)$ be an $n \times n$ matrix, continuous on an interval I ; let $n_1, \dots, n_k$ be positive integers such that $\sum_{i=1}^k n_i = n$. Let A be partitioned into submatrices A_{ij} which are $n_i \times n_j$, ($i, j = 1, \dots, k$); let E_m be the identity matrix of order m ; and let $A_m = (A_{m1} \dots A_{mk})$. In this note we consider the matrix Riccati system with side condition

$$(1) \quad Y' = -YA_mY + AY, \quad Y_m(t_0) = E_{n_m},$$

where $Y = \text{col}(Y_1, \dots, Y_k)$ and Y_i is $n_i \times n_m$. This equation is derived in a natural way as a generalization of the so-called Riccati system [1; 2]. Mainly we generalize some results of Levin [3], who treats the equation

$$(2) \quad \Gamma' = -\Gamma G_3 \Gamma - \Gamma G_4 + G_1 \Gamma + G_2,$$

where G_1, G_2, G_3 , and G_4 are $n_1 \times n_1, n_1 \times n_2, n_2 \times n_1$, and $n_2 \times n_2$ respectively, and Γ is $n_1 \times n_2$. To see that (2) is a case of (1), take $m = 2$ and

$$A = \begin{pmatrix} G_1 & G_2 \\ G_3 & G_4 \end{pmatrix}, \quad Y(t) = \begin{pmatrix} \Gamma(t) \\ E_{n_2} \end{pmatrix}.$$

For other related results see [4].

2. Let X be $n \times n$ and such that

$$(3) \quad X' = AX.$$

The partitioning of A induces a partitioning of X into submatrices X_{ij} ($i, j = 1, \dots, k$). Let $X_m = \text{col}(X_{1m} \dots X_{km})$. Then

$$(4) \quad X'_m = AX_m,$$

and, at least formally,

$$(5) \quad (X_{mm}^{-1})' = -X_{mm}^{-1}A_mX_{mm}^{-1}.$$

Thus, if X_{mm}^{-1} exists on some interval $I_0 \subset I$, $X_m X_{mm}^{-1}$ is a solution on I_0 of (1). Further, if Y is a solution of the differential equation of (1), Y_m satisfies an equation of the form $P' + HP = H$; thus from classical

Presented to the Society, January 25, 1961 under the title *The cross-ratio property for the matrix Riccati equation*; received by the editors June 13, 1960 and, in revised form, September 12, 1960.

existence and uniqueness theorems (1) has a local existence and uniqueness theorem and $Y_m(t) \equiv E_{n_m}$. Gathering together these remarks, we have

THEOREM 1. *Let $X(t)$ be the solution of (3) such that $X(t_0) = E_n$. Then the general solution near t_0 of (1) is $X(t)C(\sum_{i=1}^k X_{mi}C_i)^{-1}$, where C is constant and arbitrary, $C = \text{col}(C_1 \cdots C_k)$, C_i is $n_i \times n_m$, and C_m is nonsingular.*

3. Now let $n = rk$ and $n_i = r$ ($i = 1, \dots, k$). We proceed to write the general solution of (1) in terms of solutions of (1) rather than solutions of (3). Let $U_1, \dots, U_k$ denote solutions of (1); let $U = (U_1 \cdots U_k)$; let $Z = \text{diag}(Z_j)$, where Z_j is $r \times r$ and

$$(6) \quad Z'_j = A_m U_j Z_j \quad (j = 1, \dots, k).$$

THEOREM 2. *Let $U(t_0)$ and $Z(t_0)$ be nonsingular; let $C = \text{col}(C_j)$, the C_j being $r \times r$, constant, and arbitrary except that $\sum_{j=1}^k Z_j(t_0)C_j$ is nonsingular. Then the general solution of (1) can be written near t_0 as*

$$(7) \quad Y = UZC \cdot \left(\sum_{j=1}^k U_{mj} Z_j C_j \right)^{-1}.$$

PROOF. It is easily verified that UZ satisfies (3) and that, if $U(t_0)$ is nonsingular, $U(t)$ is nonsingular where it exists; the conclusion then follows.

Now let $U_{hij} = U_{hi} - U_{hj}$ and $V_{hij} = U_{hij}^{-1}$. It is easily verified that $(U_i - U_j)V_{hij}$ satisfies an equation of the form (1). Thus, if

$$(8) \quad [(U_i - U_q)V_{hij} - (U_i - U_j)V_{hij}]_{t=t_0} = 0,$$

it is identically zero near t_0 . Hence, if (8) is satisfied, $V_{hij}U_{hij}$ satisfies

$$(9) \quad W' = A_m U_j W - W A_m U_q.$$

Now fix q ; we may as well take $q = 1$.

THEOREM 3. *Let $|U(t_0)| \neq 0$. Let $i = k + j - 1$ ($j = 2, \dots, k$). Let U_i be a solution of (1) such that, for some r_j ($1 \leq r_j \leq k$, $r_j \neq m$), $U_{r_j ij}$ and $U_{r_j i1}$ are nonsingular at t_0 and (8) is satisfied for $q = 1$ and $h = r_j$. Let $|Z_1(t_0)| \neq 0$. Then the general solution of (1) can be written in the form (7), where $Z_j = V_{r_j ij} U_{r_j i1} Z_1$ ($j = 2, \dots, k$).*

PROOF. Note that $Z'_1 = A_m U_1 Z_1$ and that $V_{r_j ij} U_{r_j i1}$ satisfies (9) with $q = 1$; then apply Theorem 2.

We remark that, given q , j , and $h \neq k$, there does exist a set of initial values for the U_i which satisfies (8). However, if for U_j and

U_p ($p \neq j$) the same initial values for U_i satisfy (8), there is a linear combination of columns of U_i , U_j , and U_p which is zero. Thus, since $|U(t_0)| \neq 0$, each U_j requires a distinct U_i . Thus, although we need not eliminate all of the Z_j 's ($j=2, \dots, k$) in the general solution, we do need a distinct U_i for each one we do eliminate.

For $k=2$ we have Theorem 5 of [3]. For $r=1$ and general k we have a form for Y different from that given in [2].

BIBLIOGRAPHY

1. A. Chiellini, *Sui sistemi di Riccati*, Rend. Sem. Fac. Sci. Univ. Cagliari vol. 18 (1948) pp. 44-58.
2. W. J. Coles, *Linear and Riccati systems*, Duke Math. J. vol. 22 (1955) pp. 333-338.
3. J. J. Levin, *On the matrix Riccati-equation*, Proc. Amer. Math. Soc. vol. 10 (1959) pp. 519-524.
4. W. T. Reid, *A matrix differential equation of Riccati-type*, Amer. J. Math. vol. 68 (1946) pp. 237-246.

UNIVERSITY OF UTAH