INTEGRALITY OF A CERTAIN KIND OF GENUS

YASURÔ TOMONAGA

Introduction. The author introduced in [1] a new genus which is a generalization of index and A -genus. In this paper we shall give the complete proof of the integrality of our genus. However the integrality of A -genus has been proved in [2]. The author wishes to express his gratitude to Professor F. Hirzebruch for many valuable suggestions.

1. Let X^{4k} be a compact orientable differentiable $4k$ -manifold. We defined the new genus by

$$(1.1) \quad \sum_{i=0}^{\infty} p_i = \prod_i (1 + \gamma_i),$$

$$(1.2) \quad \prod_i \frac{(\gamma_i)^{1/2}}{\operatorname{tgh}(\gamma_i)^{1/2}} (1 + y \operatorname{tgh}^2(\gamma_i)^{1/2}) = \sum_{i=0}^{\infty} \Gamma_i(y, p_1, \dots, p_i)$$

where p_i denotes the Pontryagin class of the dimension $4i$. The actual values of Γ_i 's are as follows:

$$\begin{aligned} \Gamma_0 &= 1, & \Gamma_1(y, p_1) &= \left(y + \frac{1}{3}\right) p_1, \\ \Gamma_2(y, p_1, p_2) &= p_2 y^2 + \frac{1}{3} (4p_2 - p_1^2) y + \frac{1}{45} (7p_2 - p_1^2), \\ \Gamma_3(y, p_1, p_2, p_3) &= p_3 y^3 + \frac{1}{3} (6p_3 - p_1 p_2) y^2 \\ &\quad + \frac{1}{15} (17p_3 - 8p_1 p_2 + 2p_1^3) y \\ &\quad + \frac{1}{3^3 \cdot 5 \cdot 7} (62p_3 - 13p_1 p_2 + 2p_1^3). \end{aligned} \quad (1.1)$$

We put

$$(1.4) \quad \Gamma_i(y, p_1, \dots, p_i) = \sum_{j=0}^i \Gamma_{ij}(p_1, \dots, p_i) y^j.$$

Our purpose is to prove that

$$(1.5) \quad \Gamma_{kj}(p_1, \dots, p_k)[X^{4k}] = \text{integer}$$

Received by the editors September 29, 1960.

for every k and j . It suffices to show that

(i) $\Gamma_{kj}(p_1, \dots, p_k)[X^{4k}]$ does not contain the factor 2 in its denominator when it is written as a quotient of relative prime integers.

(ii) $2^\alpha \Gamma_{kj}(p_1, \dots, p_k)[X^{4k}]$ becomes an integer for a suitable integer α .

First of all let us prove (i). It is well known that

$$(1.6) \quad \frac{z^{1/2}}{\operatorname{tgh} z^{1/2}} = 1 + \sum_{k=1}^{\infty} (-1)^{k-1} \frac{2^{2k}}{(2k)!} B_k z^k \quad [3, \text{p.13}]$$

and

$$(1.7) \quad \operatorname{tgh} z = \sum_{n=1}^{\infty} (-1)^{n+1} \frac{2^{2n}(2^{2n} - 1)}{(2n)!} B_n z^{2n-1}.$$

Moreover it is known that

- (1.8) a. B_k (Bernoulli number) contains the factor 2 exactly to the first power in its denominator.
 b. $(2k)!$ is not divisible by 2^{2k} [2, p. 341].

The statement (i) easily follows from these facts and (1.2).

2. Next let us prove (ii). It suffices to show this for the complex algebraic manifold [4]. In this case we have

$$(2.1) \quad \sum_{i=0}^{\infty} p_i = \prod_i (1 + \delta_i^2), \quad \sum_{i=0}^{\infty} c_i = \prod_i (1 + \delta_i)$$

and

$$(2.2) \quad \begin{aligned} & \sum_{i=0}^{\infty} \Gamma_i(y, p_1, \dots, p_i) \\ &= \prod_i \frac{\delta_i}{\operatorname{tgh} \delta_i} (1 + y \operatorname{tgh}^2 \delta_i) \\ &= \prod_i \frac{\delta_i}{1 - e^{-\delta_i}} \left(\frac{1}{\operatorname{tgh} \delta_i} + y \operatorname{tgh} \delta_i \right) (1 - e^{-\delta_i}) \\ &= \prod_i \frac{\delta_i}{1 - e^{-\delta_i}} \left\{ \frac{\frac{1}{2} (1 + e^{-2\delta_i})}{1 - \frac{1 - e^{-\delta_i}}{2}} + y \frac{\frac{1}{2} (1 - e^{-2\delta_i}) (1 - e^{-\delta_i})}{1 - \frac{1 - e^{-2\delta_i}}{2}} \right\}. \end{aligned}$$

Hence we have

$$\begin{aligned}
 & \Gamma_k(y, p_1, \dots, p_k) \\
 &= \kappa_{4k} \left[\prod_i \frac{\delta_i}{1 - e^{-\delta_i}} \prod_i \left[\frac{1}{2} (1 + e^{-2\delta_i}) \right. \right. \\
 & \quad \cdot \left\{ 1 + \frac{1 - e^{-\delta_i}}{2} + \left(\frac{1 - e^{-\delta_i}}{2} \right)^2 + \dots + \left(\frac{1 - e^{-\delta_i}}{2} \right)^{2k} \right\} \\
 (2.3) \quad & \quad + \frac{1}{2} y (1 - e^{-2\delta_i}) (1 - e^{-\delta_i}) \left\{ 1 + \frac{1 - e^{-2\delta_i}}{2} + \left(\frac{1 - e^{-2\delta_i}}{2} \right)^2 \right. \\
 & \quad \left. \left. + \dots + \left(\frac{1 - e^{-2\delta_i}}{2} \right)^{2k-2} \right\} \right] \\
 &= \kappa_{4k} \left[\left(\sum_i y^i \sum_{a_1, \dots, a_m} A_{a_1 \dots a_m}^i \sum_{i_1 \neq \dots \neq i_m} \exp(a_1 \delta_{i_1} + \dots + a_m \delta_{i_m}) \right) \right. \\
 & \quad \left. \times \prod_i \frac{\delta_i}{1 - e^{-\delta_i}} \right]
 \end{aligned}$$

where A denotes some rational number which becomes an integer when it is multiplied by some power of 2 and a_i denotes some integer. Putting

$$(2.4) \quad \prod_{i_1 \neq \dots \neq i_m} (1 + a_1 \delta_{i_1} + \dots + a_m \delta_{i_m}) = \sum_i d_i$$

we see that d_i is an integral cohomology class, i.e., $d_i \in H^{2i}(X^{4k}, \mathbb{Z})$. Hence we have

$$(2.5) \quad \Gamma_k(p_1, \dots, p_k)[X^{4k}] = \sum_i b_i T(X^{4k}, W_i) = \sum_i b_i \chi(X^{4k}, W_i)$$

where b_i denotes some rational number which becomes an integer by multiplying some power of 2 and $T(X^{4k}, W_i)$ denotes the Todd genus with regard to W_i and W_i denotes a complex-analytic vector bundle whose Chern class is $\sum d_i$ and $\chi(X^{4k}, W_i)$ denotes the Riemann-Roch number with regard to W_i [3, p.154]. Thus we have proved (ii).

REFERENCES

1. Y. Tomonaga, *A generalization of Hirzebruch polynomial and cobordism decomposition*, Proc. Amer. Math. Soc. vol. 12 (1961) pp. 573-576.
2. A. Borel and F. Hirzebruch, *Characteristic classes and homogeneous spaces. II*, Amer. J. Math. vol. 31 (1959) pp. 315-382.
3. F. Hirzebruch, *Neue topologische Methoden in der algebraischen Geometrie*, Berlin, Springer-Verlag, 1956.
4. J. Milnor, *On the cobordism ring Ω^* and a complex analogue*, Amer. J. Math., to appear.

UTSUNOMIYA UNIVERSITY, JAPAN