SURFACE AREA OF A CONVEX BODY UNDER AFFINE TRANSFORMATIONS

C. M. PETTY

1. Introduction. Let K be a nondegenerate convex body in E^n . We will seek requirements on K such that its surface area is minimal compared to that of its affine transforms of the same volume. This problem has been studied for $n=2$ by Behrend [1], Green [2], and Gustin [3].

2. Necessary and sufficient conditions. Let S denote the surface of K , $V(K)$ its volume and $S(K)$ its surface area.

THEOREM 1. *Necessary and sufficient conditions that a nondegenerate convex body K in E^n have minimal surface area among its affine transforms of volume $V(K)$ are that*

$$\int_s u_i u_j dS_u = 0 \quad \text{for } i \neq j$$

and

$$\int_s u_i^2 dS_u = \frac{1}{n} S(K)$$

for $i, j=1, \dots, n$ where $(u_1, \dots, u_n)$ is the outer unit vector normal to S at the point with surface area element dS_u .

PROOF. We need only consider central affine transformations of the type (x, y are column vectors of length n)

$$(1) \quad y = Bx, \quad \det(B) = 1.$$

Suppose that the convex body K_i ($i=1, \dots, n$) is transformed into K_i^* by (1), then the linear combination $\lambda_1 K_1 + \dots + \lambda_n K_n$ is transformed into $\lambda_1 K_1^* + \dots + \lambda_n K_n^*$. Since volume is preserved, it follows from the definition of mixed volumes [4, p. 38] that

$$(2) \quad V(K_{p_1}, \dots, K_{p_n}) = V(K_{p_1}^*, \dots, K_{p_n}^*).$$

Let E be an ellipsoid with supporting function $E(u)$, then

$$(3) \quad nV(K, \dots, K, E) = \int_s E(u) dS_u, \quad |u| = 1.$$

Received by the editors September 15, 1960.

It is clear, then, that our problem is equivalent to keeping K fixed and minimizing $V(K, \dots, K, E)$ over all ellipsoids with center at the origin and volume

$$(4) \quad \kappa_n = \frac{\pi^{n/2}}{\Gamma(n/2 + 1)}.$$

To show the existence of a minimizing ellipsoid, let U be a solid sphere, of radius $r > 0$, contained in K and let $\bar{a}$, of length a , be a segment of maximum length contained in E . By the inclusion property of mixed volumes [4, p. 41, (5)] and the representation [4, p. 45] for the area of the projection of U , we have $V(K, \dots, K, E) \geq V(U, \dots, U, \bar{a}) = ar^{n-1}n^{-1}\kappa_{n-1}$. Consequently, a sequence of ellipsoids E_i of the above type, for which $V(K, \dots, K, E_i)$ tends to the g.l.b. of (3), is uniformly bounded and by Blaschke's selection theorem [4, p. 34] there exists a subsequence of the E_i which converges to a convex body E^* . The body E^* is necessarily an ellipsoid and, by the continuity of mixed volumes, E^* is a minimizing ellipsoid.

Now, let A^{-1} be a positive definite symmetric matrix with $\det(A^{-1}) = 1$. If E is the ellipsoid with surface

$$(5) \quad x^T A^{-1} x = 1$$

(the superscript T meaning transpose) then the polar reciprocal of E with respect to the origin has distance function

$$(6) \quad E(u) = \{u^T A u\}^{1/2}.$$

The above is clear if A^{-1} is a diagonal matrix and the general case follows by considering an orthogonal transformation. Consequently, (6) is the supporting function of E , see [4, p. 28], and if E is transformed into E^* by (1) then $E^*(u)$ is found by replacing A in (6) by BAB^T .

Let $F(B)$ be the function of n^2 variables b_{ij} given by

$$(7) \quad F(B) = \int_S E_A(u) dS_u, \quad |u| = 1,$$

where $E_A(u)$ is given by (6) and $A = BB^T$. Since BB^T is positive semi-definite, (7) is defined for all B . We wish to minimize (7), with the constraint

$$(8) \quad \det(B) = 1.$$

If we equate to zero the partial derivatives of $F(B) + \lambda(1 - \det(B))$ and use (8) we have

$$(9) \quad \left[\frac{\partial F}{\partial b_{ij}} \right] = \lambda (B^T)^{-1}.$$

Also

$$(10) \quad \left[\frac{\partial F}{\partial b_{ij}} \right] = \left[\int_S u_i u_j E_A^{-1}(u) dS_u \right] B.$$

Consequently,

$$(11) \quad A^{-1} = \frac{1}{\lambda} \left[\int_S u_i u_j E_A^{-1}(u) dS_u \right], \quad \lambda = \frac{1}{n} \int_S E_A(u) dS_u.$$

If A is any positive definite symmetric matrix with $\det(A) = 1$ which satisfies (11) then the ellipsoid with supporting function (6) will be called a critical ellipsoid of K . It is clear that the unit matrix satisfies (11) if and only if the conditions in Theorem 1 are satisfied. Consequently, a proof of the following result completes the proof of Theorem 1.

THEOREM 2. *If K is a nondegenerate convex body with surface S , then there exists one and only one positive definite symmetric matrix A with $\det(A) = 1$ such that*

$$A^{-1} = \frac{n}{\int_S E_A(u) dS_u} \left[\int_S u_i u_j E_A^{-1}(u) dS_u \right]$$

where $E_A(u)$ is given by (6).

PROOF. Suppose C is the matrix of an affine transformation of type (1) which carries K into K^* . If $F^*(D)$ is the corresponding function (7) for K^* then

$$(12) \quad F(B) = F^*(CB)$$

for all B . By (2) and (3), this is the case for B with $\det(B) = 1$ and by the special form of (7) this may be extended to all B . We have then

$$(13) \quad \left[\frac{\partial F}{\partial b_{ij}} \right] = C^T \left[\frac{\partial F^*}{\partial d_{ij}} \right], \quad D = CB.$$

Consequently, by (11), (13), (10) and (12) we conclude that a critical ellipsoid of K is carried by C into a critical ellipsoid of K^* . We may, therefore, assume that the solid unit sphere is a critical ellipsoid of K and consequently that the conditions in Theorem 1 hold. Let

$$(14) \quad E(a, u) = \left[\sum_1^n a_i^2 u_i^2 \right]^{1/2}$$

and

$$(15) \quad F(a) = \frac{1}{S(K)} \int_s E(a, u) dS_u.$$

The function $F(a)$ is defined for all points a in E^n and has the following properties:

- (1) $F(a) > 0$ for $a \neq 0$ and $F(0) = 0$,
- (2) $F(\lambda a) = \lambda F(a)$, $\lambda > 0$,
- (3) $F(a+b) \leq F(a) + F(b)$.

To prove property (3), we have by Cauchy's inequality

$$\sum a_i b_i u_i^2 \leq \left[\left(\sum a_i^2 u_i^2 \right) \left(\sum b_i^2 u_i^2 \right) \right]^{1/2}$$

and consequently

$$E(a+b, u) \leq E(a, u) + E(b, u).$$

By [4, pp. 21-22], $F(a)$ is the distance function of a convex body and, by the conditions in Theorem 1, this convex body lies entirely in the closed half-space containing the origin and bounded by the tangent plane $\sum_1^n x_i/n^{1/2} = n^{1/2}$. Now let M be the set of all points a such that $a_i > 0$ and $a_1 \cdots a_n = 1$. By the inequality between the geometric and arithmetic means we have $\sum_1^n a_i/n^{1/2} \geq n^{1/2}$ for all points in M with equality only if $a_i = 1$ ($i = 1, \cdots, n$). Consequently, $F(a)$ evaluated over all points in M has a unique absolute minimum for $a = (1, \cdots, 1)$. This completes the proof of Theorem 2 since the conditions in Theorem 1 are invariant under an orthogonal transformation.

3. The projection body. Green [2] has shown that for $n=2$ the conditions in Theorem 1 may be replaced by the requirement that the second Fourier coefficients of the supporting function $H(\cos \theta, \sin \theta) = h(\theta)$ of K vanish. We may generalize this to n -dimensions if in place of the supporting function of K we use the supporting function $\sigma(u)$ of the projection body of K , see [4, p. 45]. This may be viewed as a generalization since for $n=2$, $\sigma(\cos \theta, \sin \theta) = h(\theta + \pi/2) + h(\theta - \pi/2)$.

THEOREM 3. *If K is a nondegenerate convex body whose projection body has supporting function $\sigma(u)$, then necessary and sufficient conditions that the surface area of K be minimal among its affine transforms of volume $V(K)$ are that*

$$\int_{\Omega} u_i u_j \sigma(u) d\omega_u = 0 \quad \text{for } i \neq j$$

and

$$\int_{\Omega} u_i^2 \sigma(u) d\omega_u = \frac{\kappa_{n-1}}{n} S(K), \quad |u| = 1$$

for $i, j = 1, \dots, n$ where the integration is extended over the surface Ω of the solid unit sphere.

PROOF. The supporting function $\sigma(u)$ is given, see [4, pp. 48–49], by

$$(16) \quad \sigma(u) = \frac{1}{2} \int_S |u \cdot \tau| dS_{\tau}, \quad |\tau| = 1$$

and

$$(17) \quad S(K) = \frac{1}{\kappa_{n-1}} \int_{\Omega} \sigma(u) d\omega_u, \quad |u| = 1.$$

Let $S_2(u)$ be any surface harmonic of degree 2, then by the Funk-Hecke theorem [5, pp. 247–248] we have for unit vectors u and τ

$$\int_{\Omega} |u \cdot \tau| S_2(u) d\omega_u = \frac{\pi^{(n-1)/2}}{\Gamma((n+3)/2)} S_2(\tau).$$

Consequently, by (16),

$$\int_{\Omega} S_2(u) \sigma(u) d\omega_u = \frac{1}{2} \frac{\pi^{(n-1)/2}}{\Gamma((n+3)/2)} \int_S S_2(\tau) dS_{\tau}$$

and Theorem 3 follows directly from Theorem 1 and (17).

REFERENCES

1. F. Behrend, *Über einige Affinvarianten konvexer Bereiche*, Math. Ann. vol. 113 (1937) pp. 713–747.
2. J. W. Green, *Length and area of a convex curve under affine transformation*, Pacific J. Math. vol. 3 (1953) pp. 393–402.
3. W. Gustin, *An isoperimetric minimax*, Pacific J. Math. vol. 3 (1953) pp. 403–405.
4. T. Bonnesen and W. Fenchel, *Theorie der konvexen Körper*, New York, Chelsea Publishing Co., 1948.
5. A. Erdélyi et al., *Higher transcendental functions*, Bateman Manuscript Project, vol. 2, New York, McGraw-Hill, 1953, pp. 247–248.

LOCKHEED MISSILES AND SPACE DIVISION