STABILITY FOR INHOMOGENEOUS DIFFERENCE SCHEMES¹

T. SEIDMAN

PART I

Preliminaries. We consider a PDE

$$(1) \quad u_t - Au = f,$$

where u is a (possibly vector-valued) unknown function of a real "time" variable t and an N -dimensional real vector "space" variable x . Here A is a linear operator, constant² in t , operating on u , where u is considered a function of x alone (i.e., A acts on elements of a linear space $\mathfrak{B}$ and, for each value of t , $u(\cdot, t) \in \mathfrak{B}$). The function f is a known function of x and t . Thus u is to satisfy (1) for $t > 0$ and $x \in \mathfrak{R}$. The initial condition

$$(2) \quad u(x, 0) = \bar{u}(x) \quad (x \in \mathfrak{R})$$

and the boundary condition

$$(3) \quad \Gamma u(\cdot, t) = g(t) = g^*(\cdot, t) \quad (t > 0)$$

are assumed to hold. The operator Γ is a linear transformation from $\mathfrak{B}$ to another linear space $\mathfrak{D}$ of functions [$g(t) \in \mathfrak{D}$ for each $t > 0$] defined on $\partial \mathfrak{R}$, the boundary of $\mathfrak{R}$, such that Γu depends only on the values of u near the boundary.

We assume this problem is well-posed. That is, we assume the existence, for each fixed $T > 0$, of a one-parameter family of operators $\hat{S}_t$ acting on a space Σ with elements of the form

$$w = \begin{pmatrix} \bar{u} \\ \gamma \\ \phi \end{pmatrix},$$

where $\bar{u}$ is a function³ of $x \in \mathfrak{R}$, γ is a function of $x \in \partial \mathfrak{R}$ and $0 < t$

Received by the editors September 21, 1960.

¹ This work was performed under the auspices of the U. S. Atomic Energy Commission.

² As in [1], this is not a necessary requirement but is imposed only to avoid notational clumsiness. The modification necessary to the general case should be clear. The operator Γ appearing in equation (3) may also depend on t .

³ The values of $\bar{u}$ and ϕ will be the same sort of vectors as u and f ; those of γ will be as g .

$\leq T$, and ϕ is a function of $x \in \mathcal{R}$ and $0 < t \leq T$. We also assume that each $w \in \Sigma$ satisfies

$$(4) \quad \Gamma \bar{u} = \gamma(0).$$

The operators $\hat{S}_t$ are to send Σ into $\mathcal{R}$ in such a way that for each $w \in \Sigma$

$$(a) \quad \|\hat{S}_t w - \bar{u}\| \rightarrow 0, \quad \text{as } t \rightarrow 0+,$$

$$(b) \quad \Gamma \hat{S}_t w = \gamma(t), \quad \text{for } 0 < t \leq T,$$

$$(5) \quad (c) \quad \|(\hat{S}_{t+\delta} w - \hat{S}_t w)/\delta - A \hat{S}_t w - \phi(t)\| \rightarrow 0, \text{ as } \delta \rightarrow 0, \quad \text{for } 0 < t < T,$$

$$(d) \quad \|\hat{S}_t\| \leq M = M(T), \quad \text{for } 0 < t \leq T.$$

It is convenient to define the translation operators⁴ T_t and T_t by

$$(6) \quad \begin{aligned} [T_t \gamma](s, \tau) &= \begin{cases} \gamma(x, \tau + t), & \tau + t \leq T, x \in \partial \mathcal{R}, \\ 0, & \tau + t > T, x \in \partial \mathcal{R}; \end{cases} \\ [T_t \phi](x, \tau) &= \begin{cases} \phi(x, \tau + t), & \tau + t \leq T, x \in \mathcal{R}, \\ 0, & \tau + t > T, x \in \mathcal{R}. \end{cases} \end{aligned}$$

Clearly, $\hat{S}_t$ can be written as

$$(7) \quad \hat{S}_t = (U_t, G_t, F_t),$$

where

$$U_t \bar{u} = \hat{S}_t \begin{pmatrix} \bar{u} \\ 0 \\ 0 \end{pmatrix}, \quad G_t \gamma = \hat{S}_t \begin{pmatrix} 0 \\ \gamma \\ 0 \end{pmatrix}, \quad F_t \phi = \hat{S}_t \begin{pmatrix} 0 \\ 0 \\ \phi \end{pmatrix},$$

and it is convenient to define the semi-group $\{S_t\}$ by

$$(8) \quad S_t = \begin{pmatrix} U_t & G_t & F_t \\ 0 & T_t & 0 \\ 0 & 0 & T_t \end{pmatrix}.$$

It follows from (5) that $\{S_t\}$ is a bounded, strongly continuous⁵ semi-group, and provides the solution of the problem determined by (1), (2), and (3).

⁴ There should be no confusion caused by using the same letter for these two different operators, distinguishing them only by their arguments.

⁵ We assume that ϕ and γ are restricted so that the operators T_t are strongly continuous. For example, we may assume them to be continuous in t if the maximum norm is used. Actually, a finite number of discontinuities may be handled also without difficulty.

If we now consider, for each $\delta > 0$, a mesh μ_δ consisting of a finite number of points in $\mathcal{R}$, each taken at $t=0, \delta, 2\delta, \dots$, then (sufficiently smooth) functions in Σ may be approximated by functions determined by their values on μ_δ . We identify the approximating functions with functions on μ_δ in such a way as to approximate norms.

By a difference scheme, we mean a family of operators $\tilde{S}_\delta$ such that $(\tilde{S}_\delta)^n$ approximates $S_{n\delta}$ in an appropriate sense. We will write $\tilde{S}_\delta$ in the form

$$(9) \quad \tilde{S}_\delta = \begin{pmatrix} \tilde{u}_\delta & \tilde{G}_\delta & \tilde{F}_\delta \\ 0 & \tilde{T}_\delta & 0 \\ 0 & 0 & \tilde{T}_\delta \end{pmatrix},$$

and, if $\tilde{w}$ is the function on μ_δ corresponding to an arbitrary but sufficiently smooth function $w \in \Sigma$ of the type described above [let $\tilde{w}_1 = \tilde{u}$, $\tilde{w}_2 = \tilde{\gamma}$, $\tilde{w}_3 = \tilde{\phi}$, where $\tilde{w} = (\tilde{u}, \tilde{\gamma}, \tilde{\phi})^T$], we will ask that

$$(10) \quad \|[(\tilde{S}_\delta \tilde{w})_1 - \tilde{u}]/\delta - Au - \phi(0)\| \rightarrow 0 \quad \text{as } \delta \rightarrow 0.$$

If (10) is satisfied, we say that $\tilde{S}_\delta$ is *consistent* with the problem. If there is a constant $\tilde{M} = \tilde{M}(T)$ such that

$$(11) \quad \|\tilde{S}_\delta^n\| \leq \tilde{M} \quad \text{for } 0 \leq n\delta \leq T,$$

then we say the scheme is *stable*. If $\tilde{S}_\delta^n$ converges strongly to S_t as $\delta \rightarrow 0$ with $(n\delta) \rightarrow t$, we say the scheme is *convergent*.

Convergence implies stability. The proof here is identical with that in [1].

Suppose the scheme is convergent. Then, since $[0, T]$ is closed, either $\|\tilde{S}_\delta^n w_0\|$ is uniformly bounded for each $w_0 \in \Sigma$ and $n\delta \leq T$, or, for some fixed $t_0 \in [0, T]$ and fixed $w_0 \in \Sigma$, there would be a sequence $\delta_j \rightarrow 0$ and a sequence n_j such that $n_j \delta_j \rightarrow t_0$ for which $\|(\tilde{S}_{\delta_j})^{n_j} w_0\|$ becomes arbitrarily large. This, however, would contradict the convergence; thus, for each $w_0 \in \Sigma$ there is an $m = m(T, w_0)$ such that $\|\tilde{S}_\delta^n w_0\| \leq m$ for $0 \leq n\delta \leq T$. Since, for each δ , $\tilde{S}_\delta$ is effectively finite-dimensional, it is obviously bounded as is $\tilde{S}_\delta^n$. Hence, by the Principle of Uniform Boundedness, the operators $\tilde{S}_\delta^n$ are uniformly bounded for $n\delta \leq T$ and the scheme is stable.

Stability implies convergence. This again is substantially as presented in [1].

Let $w_0 \in \Sigma$ and $w(t) = S_t w_0$. From (5) the "first component" of $w(t)$ is the desired solution of the problem. By (5c), we have, for any $\epsilon > 0$, a δ_1 such that, for $0 < \delta < \delta_1$,

$$(12) \quad \|[\tilde{S}_i w(t) - u(t)]/\delta - Au(t) - \phi(t)\| \leq \epsilon/4\tilde{M}T \quad \text{for } 0 \leq t < T.$$

Assuming the scheme $\{\tilde{S}_i\}$ is stable, we have by (10), with $w=w(t)$, a δ_2 such that, for $0 < \delta < \delta_2$,

$$(13) \quad \|[(\tilde{S}_i \bar{w}(t))_1 - \bar{u}(t)]/\delta - Au(t) - \phi(t)\| \leq \epsilon/4\tilde{M}T \quad \text{for } 0 \leq t < T.$$

Note further that for w sufficiently smooth there is a δ_3 such that for $0 < \delta < \delta_3$

$$(14) \quad \|\bar{u} - u\| \leq \epsilon/4\tilde{M}T \quad \text{for } 0 \leq t \leq T.$$

It follows from (11), (12), and (13) that for $0 < \delta < \delta_1, \delta_2, \delta_3$

$$(15) \quad \|(\tilde{S}_i \bar{w}(t))_1 - \tilde{S}_i w(t)\| \leq 3\epsilon\delta/4\tilde{M}T \quad \text{for } 0 \leq t \leq T.$$

Let $\delta_j \rightarrow 0+$, and n_j be such that $n_j \delta_j \rightarrow t \in [0, T]$. Set

$$(16) \quad \begin{aligned} \psi_j &= [(\tilde{S}_{\delta_j})^{n_j} \bar{w}_0]_1 - \tilde{S}_{n_j \delta_j} w_0 \\ &= \sum_{k=0}^{n_j-1} [\tilde{S}_{\delta_j}^k (\tilde{S}_{\delta_j} - S_{\delta_j}) S_{(n_j-1-k)\delta_j} w_0]_1. \end{aligned}$$

Then, using the stability condition and (15), with $t = (n_j - 1 - k)\delta_j$, we have, for $\delta_j < \delta_0 = \min \{\delta_1, \delta_2, \delta_3\}$,

$$(17) \quad \|\psi_j\| \leq \sum_0^{n_j-1} \tilde{M}(3\epsilon\delta_j/4\tilde{M}T) = \frac{3}{4} (n_j \delta_j / T) \epsilon \leq \frac{3}{4} \epsilon.$$

Putting $s = |n_j \delta_j - t|$, we have

$$\pm (S_{n_j \delta_j} - S_t) w_0 = S_{t'} (S_s - I) w_0,$$

where the sign on the left denotes whether $n_j \delta_j$ is greater than or less than t , and where t' is the smaller of $n_j \delta_j$ and t . Therefore,

$$(18) \quad \|\tilde{S}_{n_j \delta_j} w_0 - \tilde{S}_t w_0\| \leq \|S_{t'}\| \cdot \|S_s w_0 - w_0\|,$$

which goes to zero as $s \rightarrow 0$ (i.e., as $j \rightarrow \infty$), according to (5d) and (5a). Thus, for any $\epsilon > 0$, there is a J_1 such that for $j > J_1$

$$(19) \quad \|\tilde{S}_{n_j \delta_j} w_0 - \tilde{S}_t w_0\| \leq \epsilon/4.$$

Letting J be greater than J_1 and also large enough so that $\delta_j < \delta_0$ for $j > J$, we have, then, from (15), (17), and (19),

$$(20) \quad \begin{aligned} \|(\tilde{S}_{\delta_j}^{n_j} \bar{w}_0)_1 - \tilde{S}_t w_0\| &= \|\psi_j + \tilde{S}_{n_j \delta_j} w_0 - \tilde{S}_t w_0\| \\ &\leq 3\epsilon/4 + \epsilon/4 = \epsilon. \end{aligned}$$

Thus, since ϵ was arbitrary, we have shown that $(\tilde{S}_{\delta_j})^{n_j}$ converges

strongly to S_i as $j \rightarrow \infty$, and, since the sequence $\{\delta_j\}$ was arbitrary, this proves convergence.

PART II

The concept of stability can be generalized to boundary value problems of the form:

$$(21) \quad \begin{aligned} Lu &= f, \\ \Gamma u &= \gamma, \end{aligned}$$

where L is a linear differential operator acting on functions $u = u(x)$ for x in a domain $\mathfrak{D}$ and Γ is a linear boundary operator mapping u into a function γ defined on $\partial\mathfrak{D}$.

If the problem is well-posed, there is a bounded linear operator S such that

$$(22) \quad S \begin{pmatrix} f \\ \gamma \end{pmatrix} = u$$

gives the solution u of (21). Clearly S is the inverse of

$$A = \begin{pmatrix} L \\ \Gamma \end{pmatrix}.$$

By a difference scheme we mean a parametrized sequence of meshes μ_δ becoming dense as $\delta \rightarrow 0$ and a family of operators $\tilde{S}_\delta$ intended⁶ to approximate S as $\delta \rightarrow 0$.

We say the scheme $\{\tilde{S}_\delta\}$ is *consistent* with (21) if for every sufficiently smooth function u

$$(23) \quad \|Au - \tilde{S}_\delta^{-1}\tilde{u}\| \rightarrow 0 \quad \text{as } \delta \rightarrow 0.$$

The scheme is called *stable* if the operators $\tilde{S}_\delta$ are uniformly bounded so there is a constant M such that

$$(24) \quad \|\tilde{S}_\delta \tilde{w}\| \leq M \|\tilde{w}\| \leq M'(\|f\| + \|\gamma\|).$$

Finally, the scheme is *convergent* if $\tilde{S}_\delta$ converges strongly to S as $\delta \rightarrow 0$ so that for a dense set of w

$$(25) \quad \|\tilde{S}_\delta \tilde{w} - Sw\| \rightarrow 0 \quad \text{as } \delta \rightarrow 0.$$

H. Keller observed⁷ that it is extremely simple to show that stability implies convergence in this situation also. Ignoring the distinction

⁶ In practice, of course, it is $\tilde{S}_\delta^{-1}$ which is given.

⁷ Private communication. This will appear in the forthcoming book on numerical analysis by E. Isaacson and H. Keller.

between such functions as u and $\bar{u}$ (the representative function agreeing with u on μ_δ and approaching u in the norm) the proof runs as follows:

From (23), there is, for any $\epsilon > 0$, a δ_0 such that, for $\delta < \delta_0$, $\|w - S_\delta^{-1}u\| < \epsilon$ where we write $Au = w$, $Sw = u$. On multiplying $w' = (w - \tilde{S}_\delta^{-1}u)$ by $\tilde{S}_\delta$ one has, from (24) $\|\tilde{S}_\delta w - u\| = \|S_\delta w'\| \leq M'\|w'\| < M'\epsilon$ which gives (25) and the scheme is convergent.

As in Part I, the converse follows from the Principle of Uniform Boundedness.

REFERENCE

1. P. Lax and R. Richtmeyer, *Survey of the stability of linear finite difference equations*, Comm. Pure Appl. Math. vol. 9 (1956) pp. 267-293.

UNIVERSITY OF CALIFORNIA, LIVERMORE

DUALITY IN HOMOGENEOUS PROGRAMMING¹

E. EISENBERG

The problem of maximizing a concave function subject to linear constraints does not have a dual, as is the case in linear programming, in which primal optimizing variables do not appear. As a special case of our principal result it will follow that such a dual does indeed exist whenever the objective function is also homogeneous.

In the linear case we are given an $m \times n$ matrix A and vectors $a \in R^n$, $b \in R^m$.² The feasibility sets X and Y are defined by: $X = R_+^m \cap \{x \mid xA \leq a\}$, $Y = R_+^n \cap \{y \mid Ay \geq b\}$. Since $xA \leq a$ if and only if $xAy \leq ay$ for all $y \in R_+^n$ (and similarly for $Ay \geq b$), we may write:

$$(1) \quad \begin{aligned} X &= R_+^m \cap \{x \mid xAy \leq \psi(y) \quad \text{all } y \in R_+^n\} \\ Y &= R_+^n \cap \{y \mid xAy \geq \phi(x) \quad \text{all } x \in R_+^m\} \end{aligned}$$

Received by the editors April 26, 1960; and in revised form, July 27, 1960 and September 12, 1960.

¹ Work on this paper was supported, in part, by the Logistics Branch of the Office of Naval Research under Contract NONR 562(15) at Brown University.

² R^m denotes the set of all real m -tuples. If $u, v \in R^m$ then $u \leq v$ means that the inequality holds for each component. In particular, $R_+^m = R^m \cap \{x \mid x \geq 0\}$. If M is a $p \times q$ matrix and N is a $q \times t$ matrix then MN represents the usual matrix product. To simplify notation, the same symbol is used for both a column vector and its transpose; the meaning will, in any case, be clear from the context.