NOTE ON THE TENSOR PRODUCT OF BANACH ALGEBRAS¹

BERNARD R. GELBAUM

0. **Introduction.** In [2] the maximal ideal space of the tensor product of two commutative Banach algebras was studied. One of the results was: Let $A_3 = A_1 \otimes_\theta A_2$ be the "greatest cross-norm" [3; 4] tensor product of two commutative Banach algebras A_1 and A_2 . Let $\mathfrak{M}_1, \mathfrak{M}_2, \mathfrak{M}_3$ be the corresponding spaces of regular maximal ideals. Then $\mathfrak{M}_3$ and $\mathfrak{M}_1 \times \mathfrak{M}_2$ are "naturally" homeomorphic if the weak* topologies are used in all spaces. In the following, we extend the discussion to the case in which no commutativity is assumed.²

1. **Tensor products.** Let A_1 and A_2 be Banach algebras and let C be the complex number system. We consider [3] a subset F of $C^{A_1 \times A_2}$:

$$F = \left\{ f \mid f \in C^{A_1 \times A_2}, f(0, x_2) = f(x_1, 0) = 0, \right.$$

$$\left. \sum_{(x_1, x_2)} |f(x_1, x_2)| \|x_1\|_1 \|x_2\|_2 < \infty \right\},$$

where $x_i \in A_i$, $i = 1, 2$. Since each f in F is nonzero on a set that is countable or finite, to each f there corresponds a sequence (finite or infinite) $\{(x_{11}, x_{21}), (x_{12}, x_{22}), \dots\}$ consisting of just the pairs (x_1, x_2) where $f(x_1, x_2) \neq 0$. Addition of elements of F is defined by addition of functions. Multiplication is defined via a form of convolution: If $f, g \in F$, $f * g = h$ is defined by:

$$h(0, x_2) = h(x_1, 0) = 0, \quad h(x_1, x_2) = \sum_{h_1 b_1 = x_1; a_2 b_2 = x_2} f(a_1, a_2) g(b_1, b_2)$$

if $x_1, x_2 \neq 0$. We note:

Presented to the Society September 4, 1959; received by the editors December 26, 1959 and, in revised form, September 3, 1960 and May 4, 1961.

¹ This research was supported in part by the United States Air Force through Air Force Office of Scientific Research of the Air Research and Development Command, under contract No. AF 49 (638)-526, and in part by the National Science Foundation under grant NSF-G11048. The author thanks these organizations for their assistance. Reproduction of the paper in whole or in part is permitted for any purposes of the United States Government.

² The author is much indebted to the referee of an earlier draft of this research. At his suggestion, arguments that were too compressed for intelligible reading have been elaborated.

$$\begin{aligned}
& \sum_{(x_1, x_2)} |h(x_1, x_2)| \|x_1\|_1 \|x_2\|_2 \\
& \leq \sum_{(a_1, a_2), (b_1, b_2)} |f(a_1, a_2)| |g(b_1, b_2)| \|a_1\|_1 \|b_1\|_1 \|a_2\|_2 \|b_2\|_2 \\
& = \left(\sum_{(a_1, a_2)} |f(a_1, a_2)| \|a_1\|_1 \|a_2\|_2 \right) \left(\sum_{(b_1, b_2)} |g(b_1, b_2)| \|b_1\|_1 \|b_2\|_2 \right) < \infty.
\end{aligned}$$

Thus $h \in F$ and the value of $h(x_1, x_2)$ is given by an absolutely convergent series. Relative to the operations defined, F is an algebra over C .

If $\|f\|'_3$ is taken as $\sum_{(x_1, x_2)} |f(x_1, x_2)| \|x_1\|_1 \|x_2\|_2$, then an analog of the proof that l_1 is a Banach space shows that F is a Banach space. The relation $\|f * g\|'_3 \leq \|f\|'_3 \|g\|'_3$ follows from the computation of the preceding paragraph. Hence F is a Banach algebra.

In F let I be the closed ideal generated by all functions f of one of the following forms (x_1, x'_1, x_2, x'_2 are arbitrary, α is a complex number):

- (i) $f(x_1, x_2) = f(x_1, x'_2) = -f(x_1, x_2 + x'_2)$, $f = 0$ otherwise;
- (ii) $f(x_1, x_2) = f(x'_1, x_2) = -f(x_1 + x'_1, x_2)$, $f = 0$ otherwise;
- (iii) $f(x_1, x_2) = -\alpha f(\alpha x_1, x_2)$, $f = 0$ otherwise;
- (iv) $f(x_1, x_2) = -\alpha f(x_1, \alpha x_2)$, $f = 0$ otherwise.

We assert that when F/I is given its quotient space norm it is isometrically isomorphic to A_3 , the greatest cross-norm tensor product $A_1 \otimes_\theta A_2$ of A_1 and A_2 . The argument rests on the following statements each of which is simply proved.

(a) In F , the set F_0 of functions f that are nonzero on sets that are at most finite is dense in F .

(b) F_0 is a normed algebra containing the generators of I . Let I_0 be the closed ideal generated in F_0 by the generators of I . Then I_0 is a linear space in F_0 and I_0 is dense in I .

(c) For f in F_0 , the quotient norm of f/I_0 is precisely the greatest cross-norm of the equivalence class represented by f .

(d) The mapping $\phi_0: F_0/I_0 \rightarrow A_3$ defined by: $\phi_0(f/I_0) = \text{equivalence class represented by } f$ is an isometry onto a dense subset of A_3 . The extension ϕ of ϕ_0 to F/I is an isometry of F/I onto A_3 .

In dealing with A_3 (where the norm will be denoted by $\|\cdot\|_3$) we shall often work with representatives of elements in F/I . For any $f \in F$, the element g given by: $g(x_1, x_2) = 1$ if $(x_1, x_2) = (f(x'_1, x_2)x'_1, x_2)$ and $g = 0$ otherwise, satisfies the equation $f/I = g/I$. Thus, ultimately an element of A_3 will be symbolized by a formal sum $\sum_{n=1}^{\infty} (x_{1n} \otimes x_{2n})$ where $\sum_{n=1}^{\infty} \|x_{1n}\|_1 \|x_{2n}\|_2 < \infty$. Very often we shall omit reference to representatives. For example, we shall say simply "the element

$\sum_{n=1}^{\infty} (x_{1n} \otimes x_{2n})$ " when we mean "the element represented by $\sum_{n=1}^{\infty} (x_{1n} \otimes x_{2n})$."

2. $\mathfrak{M}_1 \times \mathfrak{M}_2 \rightarrow \mathfrak{M}_3$. Let (M_1, M_2) be in $\mathfrak{M}_1 \times \mathfrak{M}_2$. Define the homomorphism $E_3: A_3 \rightarrow (A_1/M_1) \otimes_{\theta} (A_2/M_2)$ by the formula: $E_3(\sum_{n=1}^{\infty} (x_{1n} \otimes x_{2n})) = \sum_{n=1}^{\infty} (E_1(x_{1n}) \otimes E_2(x_{2n}))$, where E_1 and E_2 are the canonical epimorphisms $E_i: A_i \rightarrow A_i/M_i$, $i=1, 2$. Let $\|\cdot\|_I$, $\|\cdot\|_{II}$ denote the quotient norms in A_1/M_1 and A_2/M_2 . Since these algebras are nontrivial so is $(A_1/M_1) \otimes_{\theta} (A_2/M_2)$ nontrivial. Thus $E_3^{-1}(0) = M_3$ is a proper ideal in A_3 . If u_1 and u_2 are identities modulo M_1 and M_2 then $u = u_1 \otimes u_2$ is an identity modulo M_3 . Thus M_3 is regular. Since the norm $\|\cdot\|_{III}$ in $E_3(A_3)$ satisfies

$$\begin{aligned} \left\| E_3 \left(\sum_{n=1}^{\infty} (x_{1n} \otimes x_{2n}) \right) \right\|_{III} &\leq \sum_{n=1}^{\infty} \|E_1(x_{1n})\|_I \|E_2(x_{2n})\|_{II} \\ &\leq \sum_{n=1}^{\infty} \|x_{1n}\|_1 \|x_{2n}\|_2, \end{aligned}$$

we see that E_3 is norm-decreasing and hence that E_3 is continuous. Hence M_3 is closed.

If M_3 is not maximal, let $M_3 \subset N_3$, a regular maximal ideal. Let $F_3: A_3 \rightarrow A_3/N_3$ be the canonical epimorphism and define G_1 and G_2 by the formulas

$$\begin{aligned} G_1(x_1) &= E_3(ux_1), \\ G_2(x_2) &= E_3(x_2u). \end{aligned}$$

Then $G_i(A_i) \subset A_3/M_3$, $i=1, 2$. We prove now

LEMMA 1. *A regular ideal I_3 in A_3 is a right and left A_1 - and A_2 -ideal.*

PROOF. Let u be an identity modulo I_3 in A_3 , $x_1 \in A_1$, $v \in I_3$. Then $ux_1v - x_1v \in I_3$ and thus $ux_1v/I_3 - x_1v/I_3 = 0$. But $ux_1v/I_3 = (ux_1/I_3)(v/I_3) = 0$. Thus $x_1v \in I_3$. Similarly $vx_1 \in I_3$, and $x_2v, vx_2 \in I_3$ for $x_2 \in A_2$.

From Lemma 1 we see that $G_2(x_2y_2) = E_3(x_2y_2u) = E_3(x_2(m_3 + uy_2u)) = E_3(x_2uy_2u) = G_2(x_2)G_2(y_2)$ (where $m_3 \in M_3$). Note also: $\|G_2(x_1)\|_{III} = \|E_3(ux_1)\|_{III} \leq \|ux_1\|_3 \leq k\|x_1\|_1$. It is clear now that G_2 (and similarly G_1) is a continuous homomorphism.

If $(x_1, x_2) \in M_1 \times M_2$ then $ux_1 = (u_1x_1 \otimes u_2) = w$ is such that $E_3(w) = 0$, i.e., $G_1(x_1) = 0$. Thus $M_1 \subset G_1^{-1}(0)$. Similarly, $M_2 \subset G_2^{-1}(0)$. Hence either $M_i = G_i^{-1}(0)$ or $G_i(A_i) = 0$, $i=1, 2$. Since $G_i(u_i) \neq 0$, we see

³ A_3 may be regarded as a left and right A_1 - and A_2 -module, e.g., $x_1(y_1 \otimes y_2) = (x_1y_1 \otimes y_2)$, where $x_1, y_1 \in A_1$, $y_2 \in A_2$.

$G_i^{-1}(0) = M_i$. If H_i are engendered by F_3 as G_i are engendered by E_3 we find [2] $M_i = H_i^{-1}(0)$, $i = 1, 2$. Following the argument in [2], we obtain $H_i = \alpha_i G_i = \beta_i E_i$ where α_i, β_i are isometric C -automorphisms of A_i/M_i . Finally $E_3 = (\beta_1 \otimes \beta_2)^{-1} F_3$, whence $E_3(A_3)$ is simple and M_3 is maximal.

We remark that E_3 is an epimorphism. Indeed, if

$$\sum_{n=1}^{\infty} (E_1(x_{1n}) \otimes E_2(x_{2n})) = w \in (A_1/M_1) \otimes_{\theta} (A_2/M_2),$$

we can choose $(y_{1n}, y_{2n}) \in A_1 \times A_2$ so that $E_i(y_{in}) = E_i(x_{in})$, $i = 1, 2$, $n = 1, 2, \dots$, and such that $\|y_{1n}\|_1 < \|E_1(x_{1n})\|_I + 2^{-n}$, $\|y_{2n}\|_2 < \|E_2(x_{2n})\|_{II} + 2^{-n}$. (We may and do assume $\|E_1(x_{1n})\|_I = \|E_2(x_{2n})\|_{II}$. For, in each nontrivial term we may use the equivalent representative $\alpha E_1(x_{1n}) \otimes \alpha^{-1} E_2(x_{2n})$ where $\alpha = +[\|E_1(x_{1n})\|_I (\|E_2(x_{2n})\|_{II})^{-1}]^{1/2}$.) Then if $z = \sum_{n=1}^{\infty} (y_{1n} \otimes y_{2n})$ we see $E_3(z) = w$.

We note (for later use) that $G_1(A_1)$ and $G_2(A_2)$ are commuting subalgebras of A_3/M_3 . For

$$\begin{aligned} G_1(x_1)G_2(x_2) &= E_3(ux_1)E_3(x_2u) = E_3(u_1x_1 \otimes u_2)E_3(u_1 \otimes x_2u_2) \\ &= (E_1(u_1x_1) \otimes E_2(u_2)) \cdot (E_1(u_1) \otimes E_2(x_2u_2)) = E_1(x_1) \otimes E_2(x_2) \\ &= (E_1(u_1) \otimes E_2(x_2u_2)) \cdot (E_1(u_1x_1) \otimes E_2(u_2)) = E_3(x_2u)E_3(ux_1) \\ &= G_2(x_2)G_1(x_1). \end{aligned}$$

Hence [1] there is a homomorphism $T: G_1(A_1) \otimes_{\theta} G_2(A_2) \rightarrow A_3/M_3$ given by $T(\sum_{n=1}^{\infty} (G_1(x_{1n}) \otimes G_2(x_{2n}))) = \sum_{n=1}^{\infty} G_1(x_{1n})G_2(x_{2n})$. Since $T(G_1(u_1) \otimes G_2(u_2)) = \text{the identity of } A_3/M_3$, and since $G_1(A_1) \otimes_{\theta} G_2(A_2)$ is simple we see $T^{-1}(0) = 0$, i.e., T is a monomorphism. From our definitions it now follows that T is an isomorphism.

Similarly, if γ_i are isometric isomorphisms $\gamma_i: G_i(A_i) \rightarrow B_i$, $i = 1, 2$, there is an isomorphism $T': B_1 \otimes_{\theta} B_2 \rightarrow A_3/M_3$ given by

$$T' \left(\sum_{n=1}^{\infty} (y_{1n} \otimes y_{2n}) \right) = T \left(\sum_{n=1}^{\infty} (\gamma_1^{-1}(y_{1n}) \otimes \gamma_2^{-1}(y_{2n})) \right).$$

Finally, the association between pairs (M_1, M_2) and ideals M_3 is 1-1. For if $(M_1, M_2) \neq (M'_1, M'_2)$, e.g., if $M_1 \neq M'_1$, choose $x_1 \in M_1 \setminus M'_1$ and $x_2 \in M_2 \setminus M'_2$ if $M_2 \neq M'_2$ (otherwise choose $x_2 \notin M_2$). Then $E_3(x_1 \otimes x_2) = 0$, and $E'_3(x_1 \otimes x_2) \neq 0$, i.e., (M_1, M_2) and (M'_1, M'_2) engender distinct epimorphisms E_3 and E'_3 .

3. $\mathfrak{M}_3 \rightarrow \mathfrak{M}_1 \times \mathfrak{M}_2$. Consider the algebras $A_{i\epsilon_i}$ obtained by adjoining (if necessary) identities e_i to A_i , $i = 1, 2$. It is to be noted that $A_4 = A_{1\epsilon_1} \otimes_{\theta} A_{2\epsilon_2}$ is in general not the same as $A_{3\epsilon_3}$, the result of

adjoining (if necessary) an identity e_3 to A_3 . A typical element z of A_4 may be represented by $\alpha(e_1 \otimes e_2) + e_1 \otimes x_2 + x_1 \otimes e_2 + \sum_{n=1}^{\infty} (x_{1n} \otimes x_{2n})$ where $x_1, x_{1n} \in A_1$, $x_2, x_{2n} \in A_2$, $\alpha \in C$, $\sum_{n=1}^{\infty} \|x_{1n}\|_1 \|x_{2n}\|_2 < \infty$. There is an obvious isometric isomorphic copy of A_3 in A_4 . We denote the copy of A_3 by A_3 .

LEMMA 2. A_3 is an ideal in A_4 .

PROOF. Let z be as above, and let $w = \sum_{n=1}^{\infty} (w_{1n} \otimes w_{2n}) \in A_3$. Then $zw = \alpha w + wx_2 + x_1w + vw$ where $v = \sum_{n=1}^{\infty} (x_{1n} \otimes x_{2n})$. Since all four summands of zw are in A_3 the result follows.

A similar argument employing Lemma 1 shows the following.

LEMMA 3. A closed ideal I_3 of A_3 is a closed ideal in A_4 .

LEMMA 4. Let M_3 be a regular maximal ideal of A_3 , u an identity (in A_3) modulo M_3 : $uA_3 - A_3$, $A_3u - A_3 \subset M_3$. In A_4 let $M_4^0 = \{z \mid uz, zu \in M_3\}$, $A_4^0 = [x_1 \otimes e_2 + e_1 \otimes x_2] \oplus A_3$. Then M_4^0 is a (regular) ideal in A_4 , u is an identity (in A_4) modulo M_4^0 and $M_4^0 \not\subset A_4^0$.

PROOF. For $z \in M_4^0$, $w \in A_4$, we see in succession: uz , uzw , $uzwu$, $u(zwu) - zwu$, $zwu \in M_3$. Hence M_4^0 is a right ideal and by a similar proof is a left ideal.

Writing $e_4 = e_1 \otimes e_2$, we see $u(u - e_4) = u^2 - u \in M_3$, whence $u - e_4 \in M_4^0 \setminus A_4^0$.

Since (Lemma 2) A_3 is an ideal in A_4 , we find that if $z \in A_4$ then uz , $zu \in A_3$ and $(uz - z)u = u(zu) - zu$, $u(uz - z) = u(uz) - uz \in M_3$, whence $uz - z \in M_4^0$.

We now consider several cases:

Case I. Assume A_1 and A_2 have identities. Define mappings $F_i: A_i \rightarrow A_3/M_3$, $i=1, 2$ by the (symbolically given) formulas $F_1(A_1) = (A_1 \otimes e_2)/M_3$, $F_2(A_2) = (e_1 \otimes A_2)/M_3$. Then $F_i^{-1}(0)$ are regular ideals N_i of A_i , $i=1, 2$.

We find the pair associated with M_3 by treating several subcases.

Case 1. N_1 and N_2 are both maximal. Then as in §2, let them engender an $N_3 \in \mathfrak{M}_3$ and let G_3 be the canonical epimorphism $G_3: A_3 \rightarrow (A_1/N_1) \otimes_{\theta} (A_2/N_2)$. We shall show $N_3 \subset M_3$ (whence we shall have shown $N_3 = M_3$ since both are maximal ideals). Thus let $z = \sum_{n=1}^{\infty} (x_{1n} \otimes x_{2n}) \in A_3$ and assume $G_3(z) = 0$. Then

$$\sum_{n=1}^{\infty} (F_1(x_{1n}) \otimes F_2(x_{2n})) = 0.$$

$F_1(A_1)$ and $F_2(A_2)$ are commuting subalgebras of A_3/M_3 and thus $T: (A_1/N_1) \otimes_{\theta} (A_2/N_2) \rightarrow A_3/M_3$, given by $T(\sum_{n=1}^{\infty} (F_1(x_{1n}) \otimes F_2(x_{2n})))$

$= \sum_{n=1}^{\infty} F_1(x_{1n})F_2(x_{2n})$, is a homomorphism. (Since $T(F_1(u_1) \otimes F_2(u_2))$ = the identity of A_3/M_3 , and since $F_1(A_1) \otimes_{\theta} F_2(A_2)$ is simple, T is a monomorphism.) Hence

$$F_3(z) = \sum_{n=1}^{\infty} (F_1(x_{1n})F_2(x_{2n})) = T\left(\sum_{n=1}^{\infty} (F_1(x_{1n}) \otimes F_2(x_{2n}))\right) = T(0) = 0.$$

Thus $N_3 \subset M_3$.

Case 2. N_1 is not maximal; N_2 is maximal. Let $N_1 \subset M_1 \in \mathfrak{M}_1$. Then M_1 and N_2 engender an $M'_3 \in \mathfrak{M}_3$. We shall show $M'_3 \subset M_3$ (whence, again, we shall have shown $M'_3 = M_3$). This time let G'_3 be the canonical epimorphism $G'_3: A_3 \rightarrow (A_1/M_1) \otimes_{\theta} (A_2/N_2)$, and let G'_1, G'_2 arise from G'_3 according to the procedure in §2. Then $\alpha_2 G'_2 = F_2$, where α_2 is an isometric C -automorphism of $F_2(A_2)$. As indicated in §2, there is an isomorphism $T: G'_1(A_1) \otimes_{\theta} G'_2(A_2) \rightarrow A_3/M'_3$. If $L: A_1/N_1 \rightarrow (A_1/N_1)/(F_1(M_1))$ is the canonical epimorphism, then by virtue of the "second isomorphism theorem" $LF_1 = \alpha_1 G'_1$, where α_1 is an isometric C -automorphism of $G'_1(A_1)$. If $G_3(z) = 0$, then $\sum_{n=1}^{\infty} (E_1(x_{1n}) \otimes F_2(x_{2n})) = 0$, ($E_1: A_1 \rightarrow A_1/M_1$). Via C -automorphisms we can conclude $\sum_{n=1}^{\infty} (G'_1(x_{1n}) \otimes G'_2(x_{2n})) = 0$ and then that

$$\begin{aligned} (\alpha_1 \otimes \alpha_2) \left(\sum_{n=1}^{\infty} (G'_1(x_{1n}) \otimes G'_2(x_{2n})) \right) &= \sum_{n=1}^{\infty} (\alpha_1 G'_1(x_{1n}) \otimes \alpha_2 G'_2(x_{2n})) \\ &= \sum_{n=1}^{\infty} (LF_1(x_{1n}) \otimes F_2(x_{2n})) = 0. \end{aligned}$$

Remembering that α_1 and α_2 are automorphisms and applying T we find $\sum_{n=1}^{\infty} LF_1(x_{1n})F_2(x_{2n}) \equiv L_3(\sum_{n=1}^{\infty} F_1(x_{1n})F_2(x_{2n})) = L_3F_3(z) = 0$ (where $L_3: A_3/M_3 \rightarrow A_3/M'_3$ is defined by the initial identity). Since $L_3(F_1(u_1)F_2(u_2)) \neq 0$ and since A_3/M_3 is simple we see $F_3(z) = 0$, and so $M'_3 \subset M_3$.

Case 3. Both N_1 and N_2 are not maximal. Argue *mutatis mutandis* as in Case 2.

Case II. Not both A_1 and A_2 have identities. Let M_4 be a regular maximal ideal in A_4 such that (Lemma 4) $A_4^0 \not\supset M_4 \supset M_3$. Let M_4 engender M_{ie_i} , regular maximal ideals of A_{ie_i} , $i = 1, 2$ (Case I). If M_{1e_1} , say, is A_1 then clearly $A_3/M_4 = 0$, whence $u/M_4 = 0$, a contradiction. Hence $M_{ie_i} \not\supset A_i$, and the regular maximal ideals we seek are $M_i \equiv M_{ie_i} \cap A_i$, $i = 1, 2$.

In every case we get a pair (M_1, M_2) in $\mathfrak{M}_1 \times \mathfrak{M}_2$ and this pair in turn engenders M_3 . Since the association $\mathfrak{M}_1 \times \mathfrak{M}_2 \rightarrow \mathfrak{M}_3$ is 1-1, (M_1, M_2) is uniquely determined by M_3 : $(M_1, M_2) = \bar{i}(M_3)$. We have shown $\bar{i}$ is 1-1 and that $\bar{i}(\mathfrak{M}_3) = \mathfrak{M}_1 \times \mathfrak{M}_2$.

THEOREM. *When hk -topologies are used throughout, $\bar{i}$ is continuous but not generally bicontinuous.*

PROOF. We shall show that $\bar{i}^{-1}$ is closed. Let F_1 be closed in $\mathfrak{M}_1$. Then $F_3 = F_1 \times \mathfrak{M}_2$ is closed in $\mathfrak{M}_1 \times \mathfrak{M}_2$. Let $K_3 = \bar{i}^{-1}(F_3)$. We show that K_3 is closed. To this end let $M_3^0 \supset k(K_3)$ and assume $M_3^0 \notin K_3$. Thus if $\bar{i}(M_3^0) = (M_1^0, M_2^0)$, then $M_1^0 \notin F_1$. Hence $M_1^0 \not\supset k(F_1)$ and there is an x in $k(F_1)$ such that $x/M_1^0 \neq 0$. But if y is an arbitrary element of A_2 , then $z = x \otimes y$ in A_3 is actually in $k(K_3)$. For if $M_3 = \bar{i}^{-1}(M_1, M_2)$ is in K_3 , then M_1 is in F_1 and hence $x/M_1 = 0$, whence $z/M_3 = (x/M_1) \otimes (y/M_2) = 0$. Therefore z is in M_3 , and since M_3 is arbitrary in K_3 , z is in $k(K_3)$. Thus any such z is in M_3^0 . But if we choose y so that $y/M_2^0 \neq 0$, then $z/M_3^0 \neq 0$. This contradiction shows that K_3 is closed. Similarly, if F_2 is closed in $\mathfrak{M}_2$ then $\bar{i}^{-1}(\mathfrak{M}_1 \times F_2)$ is closed.

Since any closed set in $\mathfrak{M}_1 \times \mathfrak{M}_2$ is of the form

$$\bigcap_r [(F_1^r \times \mathfrak{M}_2) \cup (\mathfrak{M}_1 \times F_2^r)]$$

where F_i^r is closed in $\mathfrak{M}_i$, $i = 1, 2$, it follows that $\bar{i}^{-1}$ is closed and thus that $\bar{i}$ is continuous.

We are indebted to John Lindberg for suggesting the following example showing that $\bar{i}^{-1}$ need not be continuous.

EXAMPLE. Let A_1 be the commutative Banach algebra of functions analytic in the interior of the unit disc D_z and continuous on the entire unit disc D_z . The maximal ideal space of A_1 may be identified with D_z although the hk -topology constitutes a genuine weakening of the usual topology of D_z . In fact, a hk -closed set F in D_z has only countably many points in the (usual) interior of D_z unless F is the whole of D_z . Let A_2 be $C(D_w)$, i.e., the algebra of all continuous functions on D_w . Then the maximal ideal space of A_2 in the hk -topology is homeomorphic with D_w in its usual topology.

In A_3 let v be the element $z \otimes 1 - 1 \otimes w$ (z and w here are the usual complex variables). Then in $\mathfrak{M}_1 \times \mathfrak{M}_2$ the set $K = \{(z, w) \mid z = w\}$ is the hull of v , and so $\bar{i}^{-1}(K)$ is closed in the hk -topology of $\mathfrak{M}_3$. Thus $U = \mathfrak{M}_3 \setminus \bar{i}^{-1}(K)$ is open. We shall show that $\bar{i}(U)$ is not open in $\mathfrak{M}_1 \times \mathfrak{M}_2$.

If $\bar{i}(U)$ were open, there would be open sets U_i in $\mathfrak{M}_i$ such that $U_1 \times U_2 \subset \bar{i}(U)$, and U_2 could be taken as an ordinary open circle in the interior of D_w . Since $F_1 = \mathfrak{M}_1 \setminus U_1$ is hk -closed in D_z , F_1 has at most countably many points in the ordinary interior of D_z . Let U_2^* be the circle in D_z that consists of the same set of complex numbers as the set comprising U_2 . Then $U_2^* \setminus F_1 \neq \emptyset$ since F_1 meets the ordinary interior of D_z in at most countably many points. But then $U_1 \times U_2 \supset (U_2^* \setminus F_1) \times U_2$. Clearly, any point z in $U_2^* \setminus F_1$ corresponds to a point

w in U_2 such that $z=w$, i.e., $(U_2^* \setminus F_1) \times U_2$ meets K and thus $U_1 \times U_2$ cannot lie in the complement $\bar{i}(U)$ of K .

REMARK. In the last paragraph of p. 302 and the first two lines of p. 303 [2] the proof given is incorrect since the u_i chosen there vary with the M_i . The following amended proof should be substituted.

Let M_{03} be in $\mathfrak{M}_3$ and if $\bar{i}(M_{03}) = (M_{01}, M_{02})$, let $N(M_{0i})$ in $\mathfrak{M}_i$ be of the form:

$$N(M_{0i}) = \{M_i \mid |a_{ji}^+(M_i) - a_{ji}^+(M_{0i})| < r_i, j = 1, 2, \dots, J_i\}, \\ i = 1, 2.$$

To prove the continuity of $\bar{i}$ it suffices to find an $N(M_{03})$ such that $\bar{i}(N(M_{03})) \subset N(M_{01}) \times N(M_{02})$. To this end let u_{0i} be identities modulo M_{0i} , $i = 1, 2$ and let

$$N(M_{03}) = \{M_3 \mid \begin{aligned} &|(a_{j1}^k \otimes u_{02})^+(M_3) - (a_{j1}^k \otimes u_{02})^+(M_{03})| < s_k, \\ &k = 1, 2, j = 1, 2, \dots, J_1; \\ &|(u_{01} \otimes a_{j2}^m)^+(M_3) - (u_{01} \otimes a_{j2}^m)^+(M_{03})| < v_m, \\ &m = 1, 2, j = 1, 2, \dots, J_2; \\ &|(u_{01} \otimes u_{02})^+(M_3) - 1| < w \}. \end{aligned}$$

We treat two cases: (1) For all (j, i) $a_{ji}^+(M_{0i}) \neq 0$. Then for appropriate choices of the s_k , v_m and w , we can assure that

$$\left| \frac{(a_{j1}^2 \otimes u_{02})^+(M_{03})}{(a_{j1} \otimes u_{02})^+(M_{03})} - \frac{(a_{j1}^2 \otimes u_{02})^+(M_3)}{(a_{j1} \otimes u_{02})^+(M_3)} \right| \equiv |a_{j1}^+(M_{01}) - a_{j1}^+(M_1)| < r_1$$

and similarly that $|a_{j2}^+(M_{02}) - a_{j2}^+(M_2)| < r_2$, where $\bar{i}(M_3) = (M_1, M_2)$. In this case then, $\bar{i}(N(M_{03})) \subset N(M_{01}) \times N(M_{02})$. (2) Some $a_{ji}^+(M_{0i}) = 0$. For the (j, i) combinations for which $a_{ji}^+(M_{0i}) \neq 0$, s_k , v_m and w are chosen as in (1). For the other (j, i) combinations, since $|u_{0i}^+(M_i)|$ is bounded, we see that for suitable w and M_3 in $N(M_{03})$ $|u_{0i}^+(M_i)|$ is bounded away from 0 and then for suitable s_k and v_m $|a_{ji}^+(M_i)|$ is small. The result follows upon a final revised choice of s_k , v_m and w .

BIBLIOGRAPHY

1. N. Bourbaki, *Algèbre*, Chapter III, Paris, 1948.
2. B. Gelbaum, *Tensor products of Banach algebras*, Canad. J. Math. vol. 11 (1959) pp. 297-310.
3. A. Grothendieck, *Produits tensoriels topologiques et espaces nucléaires*, Memoirs Amer. Math. Soc. no. 16, 1955.
4. R. Schatten, *A theory of cross-spaces*, Princeton, New Jersey, Princeton University Press, 1950.

UNIVERSITY OF MINNESOTA