MATHEMATICAL PEARLS

The purpose of this department is to publish very short papers of an unusually elegant and polished character, for which there is normally no other outlet.

A RECURRENCE FORMULA FOR $\zeta(2n)$

L. CARLITZ¹

H. T. Kuo [2] has proved the formula

$$(1) \quad C_{2n} = \frac{\pi(2\pi)^{2n-1}}{4((n-1)!)^2(2n-1)} + \frac{1}{(n-1)!} \sum_{k=0}^{[n/2]} \frac{C_{2k}(2\pi)^{2n-2k}}{(n-2k)!(2n-2k)} \\ + \frac{1}{\pi} \sum_{k=0}^{[n/2]} \sum_{j=0}^{[n/2]} (-1)^{k-j} \frac{C_{2k}C_{2j}(2\pi)^{2n-2k-2j+1}}{(n-2k)!(n-2j)!(2n-2k-2j+1)}$$

where $n \geq 1$ and $C_0 = -1/2$, $C_{2k} = \zeta(2k)$ for $k = 1, 2, 3, \dots$.

Since

$$\zeta(2k) = \sum_{n=1}^{\infty} \frac{1}{n^{2k}} = (-1)^{k-1} \frac{(2\pi)^{2k}}{2(2k)!} B_{2k},$$

where B_{2k} is a Bernoulli number in the even suffix notation, (1) implies a rather complicated recursion formula for the Bernoulli numbers. It may be of interest to point out that a somewhat simpler and slightly more general formula of this kind can be obtained very rapidly by making use of the identity ([1; 3, p. 75])

$$(2) \quad B_m(x)B_n(x) = \sum_r \left\{ \binom{m}{2r} n + \binom{n}{2r} m \right\} \frac{B_{2r}B_{m+n-2r}(x)}{m+n-2r} \\ + (-1)^{m-1} \frac{m!n!}{(m+n)!} B_{m+n} \quad (m+n \geq 2),$$

where

$$B_n(x) = \sum_{k=0}^n \binom{k}{n} B_k x^{n-k}.$$

Integrating (2) from 0 to 1 and making use of

$$\int_0^1 B_{2k}(x) dx = \frac{B_{2k+1}(1) - B_{2k+1}}{2k+1} = 0 \quad (k \geq 1),$$

we get

Received by the editors March 6, 1961.

¹ Supported in part by National Science Foundation, grant G-9425.

$$\begin{aligned}
 (3) \quad & \sum_{j=0}^m \sum_{k=0}^n \binom{m}{j} \binom{n}{k} \frac{B_j B_k}{m+n-j-k+1} \\
 & = (-1)^{m-1} \frac{m!n!}{(m+n)!} B_{m+n} \quad (m+n \text{ even, } mn > 0).
 \end{aligned}$$

REFERENCES

1. L. Carlitz, *Note on the integral of the product of several Bernoulli polynomials*, J. London Math. Soc. vol. 34 (1959) pp. 361-363.
2. Huan-Ting Kuo, *A recurrence formula for $\zeta(2n)$* , Bull. Amer. Math. Soc. vol. 55 (1949) pp. 573-574.
3. N. Nielsen, *Traité élémentaire des nombres de Bernoulli*, Paris, Gauthier-Villars, 1923.

DUKE UNIVERSITY

MINKOWSKI'S THEOREM ON NONHOMOGENEOUS APPROXIMATION

IVAN NIVEN¹

For θ irrational, let γ be any real number such that $\gamma = \theta a + b$ has no solution in integers a and b . We give a short proof of Minkowski's classic result that there are infinitely many pairs of integers satisfying $F < 1/4$, where $F = F(\theta, \gamma, x, y) = |x| \cdot |\theta x + y + \gamma|$. First we prove that given any real numbers α and β there exists an integer u such that $|u - \beta| < 1$ and such that at least one of the following holds:

$$(A) \quad |u - \alpha| \cdot |u - \beta| \leq 1/4; \quad |u - \alpha| \cdot |u - \beta| \leq |\beta - \alpha|/2.$$

If β is an integer, set $u = \beta$. Otherwise define the integer n by $n < \beta < n+1$. If $n \leq \alpha \leq n+1$ then $|n - \alpha| \cdot |n+1 - \alpha| \leq 1/4$ and similarly for β , and so

$$|n - \alpha| \cdot |n - \beta| \cdot |n+1 - \alpha| \cdot |n+1 - \beta| \leq 1/16.$$

Hence $u = n$ or $u = n+1$ gives inequality (A₁). The cases $n > \alpha$ and $\alpha > n+1$ are symmetric, and we treat $n > \alpha$. We note that

$$\begin{aligned}
 & 2(n - \alpha)^{1/2}(n + 1 - \beta)^{1/2}(\beta - n)^{1/2}(n + 1 - \alpha)^{1/2} \\
 & \leq (n - \alpha)(n + 1 - \beta) + (\beta - n)(n + 1 - \alpha) = \beta - \alpha
 \end{aligned}$$

Received by the editors April 14, 1961.

¹ Supported in part by the Office of Naval Research.