$$\begin{aligned}
 (3) \quad & \sum_{j=0}^m \sum_{k=0}^n \binom{m}{j} \binom{n}{k} \frac{B_j B_k}{m+n-j-k+1} \\
 &= (-1)^{m-1} \frac{m!n!}{(m+n)!} B_{m+n} \quad (m+n \text{ even, } mn > 0).
 \end{aligned}$$

REFERENCES

1. L. Carlitz, *Note on the integral of the product of several Bernoulli polynomials*, J. London Math. Soc. vol. 34 (1959) pp. 361-363.
2. Huan-Ting Kuo, *A recurrence formula for $\zeta(2n)$* , Bull. Amer. Math. Soc. vol. 55 (1949) pp. 573-574.
3. N. Nielsen, *Traité élémentaire des nombres de Bernoulli*, Paris, Gauthier-Villars, 1923.

DUKE UNIVERSITY

MINKOWSKI'S THEOREM ON NONHOMOGENEOUS APPROXIMATION

IVAN NIVEN¹

For θ irrational, let γ be any real number such that $\gamma = \theta a + b$ has no solution in integers a and b . We give a short proof of Minkowski's classic result that there are infinitely many pairs of integers satisfying $F < 1/4$, where $F = F(\theta, \gamma, x, y) = |x| \cdot |\theta x + y + \gamma|$. First we prove that given any real numbers α and β there exists an integer u such that $|u - \beta| < 1$ and such that at least one of the following holds:

$$(A) \quad |u - \alpha| \cdot |u - \beta| \leq 1/4; \quad |u - \alpha| \cdot |u - \beta| \leq |\beta - \alpha|/2.$$

If β is an integer, set $u = \beta$. Otherwise define the integer n by $n < \beta < n+1$. If $n \leq \alpha \leq n+1$ then $|n - \alpha| \cdot |n+1 - \alpha| \leq 1/4$ and similarly for β , and so

$$|n - \alpha| \cdot |n - \beta| \cdot |n+1 - \alpha| \cdot |n+1 - \beta| \leq 1/16.$$

Hence $u = n$ or $u = n+1$ gives inequality (A₁). The cases $n > \alpha$ and $\alpha > n+1$ are symmetric, and we treat $n > \alpha$. We note that

$$\begin{aligned}
 & 2(n - \alpha)^{1/2}(n + 1 - \beta)^{1/2}(\beta - n)^{1/2}(n + 1 - \alpha)^{1/2} \\
 & \leq (n - \alpha)(n + 1 - \beta) + (\beta - n)(n + 1 - \alpha) = \beta - \alpha
 \end{aligned}$$

Received by the editors April 14, 1961.

¹ Supported in part by the Office of Naval Research.

and so (A_2) must hold for $u=n$ or $u=n+1$.

Now by the pigeon-hole method [1, p. 1 or 2, p. 42] it is known that there exist infinitely many pairs of integers h, k such that $|k| \cdot |k\theta - h| < 1$. For each such pair choose integers r, s such that $|rh - sk + \gamma k| \leq 1/2$. Apply (A) with

$$\alpha = r/k$$

and

$$\beta = (r\theta - s + \gamma)/(k\theta - h),$$

and define $x=r-uk, y=-s+uh$. Then we get $|\theta x + y + \gamma| < |k\theta - h|$ and $F < 1/4$ from (A_1) , $F \leq 1/4$ from (A_2) . Since $k\theta - h$ can be made arbitrarily small, and since $\theta x + y + \gamma \neq 0$, we get infinitely many pairs x, y satisfying $F \leq 1/4$. But at most one pair can give $F = 1/4$, because

$$\theta x_1 + y_1 + \gamma = \pm (4x_1)^{-1} \quad \text{and} \quad \theta x_2 + y_2 + \gamma = \pm (4x_2)^{-1}$$

would imply the rationality of $\theta(x_1 - x_2) + y_1 - y_2$ and hence of θ . This proof can be readily extended to Minkowski's theorem on the product of two linear forms, as will be shown elsewhere. A proof that $1/4$ is the best possible constant is given in [1, p. 49].

REFERENCES

1. J. W. S. Cassels, *An introduction to Diophantine approximation*, Cambridge Tract No. 45, 1957.
2. Ivan Niven, *Irrational numbers*, Carus Mathematical Monographs, No. 11, New York, Wiley, 1956.

UNIVERSITY OF OREGON