ON QUASI-REFLEXIVE BANACH SPACES

Y. CUTTLE

1. Introduction. In the first part of this paper we study some properties of Schauder bases in quasi-reflexive Banach spaces. It is shown that if a quasi-reflexive space of order 1 has a Schauder basis, this basis must be either shrinking or boundedly complete. This result should be compared with the well-known theorem which states that if a reflexive Banach space has a Schauder basis, then this basis must be both shrinking and boundedly complete. If a quasi-reflexive space has a shrinking (boundedly complete) Schauder basis, then its n th conjugate space has a boundedly complete (shrinking) basis if n is odd or a shrinking (boundedly complete) basis if n is even. Furthermore if the first conjugate space of a quasi-reflexive space has a boundedly complete basis, then the space itself has a shrinking basis.

The last two sections give conditions under which a quasi-reflexive space is in fact reflexive: if every continuous linear functional attains its supremum on the unit sphere, or if the space is either smooth or rotund, then it is reflexive.

2. Definitions and notation. Let B be a Banach space, B^* and B^{**} its first and second conjugate spaces. The canonical isomorphism of B into B^{**} will be denoted by π . If A is a subspace of B we will denote by A^+ the annihilator of A in B^* . Following [1] we define B to be *quasi-reflexive of order n* if $B^{**}/\pi B$ is (finite) n -dimensional.

Let w denote the set of positive integers. A sequence $(x_i; i \in w) \subset B$ will be called a *basis* if for each $x \in B$ there exists a unique sequence of real numbers $(a_i; i \in w)$ such that $\lim_{n \in w} \|\sum_{i \leq n} a_i x_i - x\| = 0$; we then write $x = \sum_{i \in w} a_i x_i$. A basis $(x_i; i \in w)$ for B will be called *boundedly complete* if for each sequence $(a_i; i \in w)$ of real numbers such that the sequence $(\|\sum_{i \leq n} a_i x_i\|; n \in w)$ is bounded there exists an $x \in B$ such that $x = \sum_{i \in w} a_i x_i$; it will be called *shrinking* if $\lim_{n \in w} \|x^*\|_n = 0$ for each $x^* \in B^*$ where $\|x^*\|_n = \sup \{x^*(x) : x = \sum_{i > n} a_i x_i \text{ and } \|x\| \leq 1\}$, and *monotone* if for each $x = \sum_{i \in w} a_i x_i$ in B $\|\sum_{i \leq m} a_i x_i\|$ is a non-decreasing function of m .

A Banach space is *smooth* if for each $x \in B$ with $\|x\| = 1$ there exists a unique $x^* \in B^*$ such that $\|x^*\| = 1$ and $x^*(x) = 1$. If for each $x^* \in B^*$ with $\|x^*\| = 1$ there exists at most one $x \in B$ such that $\|x\| = 1$ and $x^*(x) = 1$, then the space is called *rotund*. It is easy to see that this terminology is equivalent to that of [2]. Finally a Banach space is

Received by the editors July 22, 1960 and, in revised form, November 21, 1960.

called *subreflexive* [6] if the set of bounded linear functionals which attain their supremum on the unit sphere of B is dense in B^* .

3. Basis in quasi-reflexive spaces. If B is a quasi-reflexive space of order n then, by definition, we write $B^{**} = \pi B \oplus F$ where F is a finite n -dimensional subspace. It follows that if $(x_i; i \in w)$ is a basis for B then the sequence $(y_1^{**}, \dots, y_n^{**}, \pi x_i; i \in w)$, where $y_1^{**}, \dots, y_n^{**}$ is a basis for F , is a basis for B^{**} which we will call a basis corresponding to $(x_i; i \in w)$.

3.1. THEOREM. *Let B be a quasi-reflexive space. If B has a basis, then this basis is shrinking (boundedly complete) if and only if the corresponding basis in B^{**} is shrinking (boundedly complete).*

PROOF. If $(x_i; i \in w)$ is a shrinking basis for B and $(y_1^{**}, \dots, y_n^{**}, \pi x_i; i \in w)$ a corresponding basis for B^{**} , then for $x^{***} \in B^{***}$ we have

$$\lim_m \|x^{***}\|_m = \lim_{m; m > n} \|x^{***}\|_m \leq \lim_m \sup_{\|x\| \leq 1} \left\{ x^{***}(\pi x) : x = \sum_{i \in m} a_i x_i \right\}.$$

By Theorem 15 [3], $B^{***} = \pi_1 B^* \oplus (\pi B)^+$, where π_1 is the canonical isomorphism of B^* into B^{***} , so if $x^{***} = \pi_1 x^* + y^{***}$, where $x^* \in B^*$ and $y^{***} \in (\pi B)^+$ we obtain

$$\lim_m \|x^{***}\|_m \leq \lim_m \sup_{\|x\| \leq 1} \left\{ x^*(x) : x = \sum_{i \in m} a_i x_i \right\} = \lim_m \|x^*\|_m = 0.$$

The corresponding basis for B^{**} is therefore shrinking.

If $(x_i; i \in w)$ is a boundedly complete basis for B , let $(a_i; i \in w)$ be a sequence of real numbers such that the sequence

$$\left(\left\| \sum_{i \leq \min(m, n)} a_i y_i^{**} + \sum_{n < i \leq m} a_i \pi x_{i-n} \right\| ; m \in w \right)$$

is bounded. Since B is quasi-reflexive there exists a continuous projection of B^{**} onto πB , the sequence $\{\|\sum_{n < i \leq m} a_i \pi x_{i-n}\|; m \in w\} = \{\|\sum_{i \leq m} a_{n+i} x_i\|; m \in w\}$ is bounded and there exists $x \in B$ such that $x = \sum_{i \in w} a_{n+i} x_i$. Let $y^{**} = \sum_{i \leq n} a_i y_i^{**} + \pi x$, then $y^{**} = \sum_{i \leq n} a_i y_i^{**} + \sum_{n < i} a_i \pi x_{i-n}$ and the corresponding basis in B^{**} is therefore boundedly complete. The converse in each case is clear.

3.2. COROLLARY. *Let B be a quasi-reflexive space with a monotone shrinking basis; then $B^{(n)}$ is subreflexive for all $n \geq 0$ where $B^{(0)} = B$ and $B^{(n)} = (B^{(n-1)})^*$.*

PROOF. This is a direct consequence of [6] and the above theorem.

3.3. THEOREM. *Let B be a quasi-reflexive space. If B has a boundedly complete basis then B^* has a shrinking basis.*

PROOF. Let $(x_i; i \in w)$ be a boundedly complete basis for B . Let $(x_i^*; i \in w)$ be the sequence in B^* biorthogonal to $(x_i; i \in w)$ and A the closed subspace of B^* spanned by $(x_i^*; i \in w)$. Then by Lemma 2 [2, p. 70], $B^{**} = \pi B \oplus A^+$ and since B is quasi-reflexive A^+ must be finite n -dimensional; let $y_1^{**}, \dots, y_n^{**}$ be a basis for A^+ and pick $y_1^*, \dots, y_n^*$ in B^* such that $y_i^{**}(y_j^*) = \delta_{ij}$. Let R be the subspace spanned by $y_1^*, \dots, y_n^*$, then $B^* = R \oplus A$ and $(y_1^*, \dots, y_n^*, x_i^*; i \in w)$ is a basis for B^* , since $(x_i^*; i \in w)$ is a basis for A .

Let $(x_i^{**}; i \in w)$ be the sequence in B^{**} biorthogonal to $(y_1^*, \dots, y_n^*, x_i^*, i \in w)$. We will show that $(x_i^{**}; i \in w)$ spans B^{**} ; by Lemma 1 [2, p. 70] this will prove that the basis for B^* is shrinking. Let $(y_1^{**}, \dots, y_n^{**}, \pi x_i; i \in w)$ be the corresponding basis for B^{**} , then one verifies easily that $x_i^{**} = y_i^{**}$, $i \leq n$ and $x_{n+i}^{**} = \pi x_i - \sum_{k \leq n} y_k^*(x_i) x_k^{**}$, $i \in w$. If $x^{**} \in B^{**}$, $x^{**} = \pi x + y^{**}$ with $x \in B$ and $y^{**} = \sum_{i \leq n} \beta_i x_i^{**} \in A^+$ and one can write $x^{**} = \sum_{i \in w} x_i^*(x) \pi x_i + \sum_{i \leq n} \beta_i x_i^{**} = \sum_{i \in w} x_i^*(x) x_{n+i}^{**} + \sum_{i \leq n} (\beta_i + y_i^*(x)) x_i^{**}$. Therefore the sequence $(x_i^{**}; i \in w)$ spans B^{**} and the basis for B^* is shrinking.

3.4. COROLLARY. *Let B be a quasi-reflexive space. If B has a basis which is shrinking (boundedly complete) then there exist quasi-reflexive spaces $B^{(j)}$, $j \in w$ such that $B^0 = B$, $B^{(j+1)} = (B^{(j)})^*$, $B^{(2k)}$ has a shrinking (boundedly complete) basis and $B^{(2k+1)}$ has a boundedly complete (shrinking) basis.*

PROOF. The proof follows from Lemma 3.4 [1], Corollary 1 [2, p. 71] and Theorems 3.1 and 3.3 above.

Let B^* have a basis. Whether or not this implies the existence of a basis for B appears to be an open question. In case B is quasi-reflexive the following theorem provides a partial answer.

3.5. THEOREM. *Let B be a quasi-reflexive Banach space. If B^* has a boundedly complete basis then B has a shrinking basis.*

PROOF. Let $(x_i^*; i \in w)$ be a boundedly complete basis. By Lemma 3.4 [1] B^* is quasi-reflexive and so, as in the proof of the preceding theorem $B^{***} = \pi_1 B^* \oplus A^+$ where A is the subspace spanned by $(x_i^*; i \in w)$ the sequence biorthogonal to $(x_i^*; i \in w)$ and A^+ is finite n -dimensional. Also $B^{**} = R \oplus A$ where R is finite n -dimensional and A is a total subspace of B^{**} . By Theorems 14 and 16 of [3] there exists an isomorphism T from B^* onto A^* such that $(Tx^*)(y^{**}) = y^{**}(x^*)$ for all $x^* \in B^*$ and $y^{**} \in A$. By Theorem 3.6 of [1] there exists an

isomorphism S of A onto B . It is easy to verify that the sequence $(Sx_i^{**}; i \in w)$ is a basis for B . To see that this basis is shrinking consider the sequence $(z_i^*; i \in w)$ in B^* biorthogonal to $(Sx_i^{**}; i \in w)$. The sequence $(Tx_i^*; i \in w)$ in A^* is biorthogonal to $(x_i^{**}; i \in w)$, therefore $Sy^{**} = S \sum_{i \in w} (Tx_i^*) (y^{**}) x_i^{**} = \sum_{i \in w} y^{**}(x_i^*) Sx_i^{**}$ for all $y^{**} \in A$; it follows that $z_i^*(Sy^{**}) = y^{**}(x_i^*)$ for all $i \in w$ and all $y^{**} \in A$; since A is total we have $T^{-1}S^*z_i^* = x_i^*$ and so $(z_i^*; i \in w)$ is the isomorphic image of the basis $(x_i^*; i \in w)$. We conclude that $(Sx_i^{**}; i \in w)$ is a shrinking basis for B .

If B is a reflexive space and B has a basis, then it is well known (see, for example, [2]) that this basis must be both shrinking and boundedly complete. The following theorem holds for quasi-reflexive spaces.

3.6. THEOREM. *Let B be a quasi-reflexive space of order 1, then any basis for B is either shrinking or boundedly complete but not both.*

PROOF. Let $(x_i; i \in w)$ be a basis for the space B which is quasi-reflexive of order 1. If $(x_i; i \in w)$ is not shrinking, then, by Lemma 1 [2, p. 70], A the closed subspace spanned by $(x_i^*; i \in w)$, the sequence in B^* biorthogonal to $(x_i; i \in w)$, is a proper subspace of B^* and consequently A^+ is nonvoid. Since B has order 1 we can write $B^{**} = \pi B \oplus \{ay^{**}\}$, $y^{**} \notin \pi B$, a any real number. Let $z_0^{**} \in A^+$; then $z_0^{**} = \pi x_0 + a_0 y^{**}$ for some $x_0 \in B$ and $a_0 \neq 0$, and for any $x^* \in A$ we have $x^*(x_0) = -a_0 y^{**}(x^*)$. If $z^{**} = \pi x + ay^{**} \in A^+$ we must have $x^*(x) = (a/a_0)x^*(x_0)$ for all $x^* \in A$; A being total this implies that $x = (a/a_0)x_0$ and so $z^{**} = (a/a_0)(\pi x_0 + a_0 y^{**}) = (a/a_0)z_0^{**}$. We conclude that $A^+ = \{az_0^{**}\}$ is 1-dimensional. It is easy to see that one can write $B^{**} = \pi B \oplus A^+$. By Lemma 2 [2, p. 70] this implies that $(x_i; i \in w)$ is boundedly complete. Finally if $(x_i; i \in w)$ were both shrinking and boundedly complete then B would be reflexive by Theorem 3 [2, p. 71], contradicting the hypothesis.

4. The supremum of functionals in quasi-reflexive spaces. James [4] has shown that if all bounded linear functionals on a separable Banach space attain their supremum on the unit sphere then the space is reflexive. Whether this result is true for a general Banach space is not known. However the result can be extended to non-separable quasi-reflexive Banach space.

4.1. THEOREM. *Let B be a quasi-reflexive Banach space. If all bounded linear functionals on B attain their supremum on the unit sphere of B then B is reflexive.*

PROOF. Assume the hypothesis of the theorem. If B is separable then

James' result can be applied. If B is not separable then by Theorem 4.6 of [1] there exists a nonseparable reflexive subspace Z such that B/Z is separable.

The mapping T of Z^+ onto $(B/Z)^*$ defined by $Tx^*(x+Z) = x^*(x)$ for all $x^* \in Z^+$ and $x \in B$, is an isometric isomorphism. If $x^* \in Z^+$ there exists $x_0 \in B$ such that $\|x_0\| = 1$ and $x^*(x_0) = \|x^*\|$, and therefore $Tx^*(x_0+Z) = x^*(x_0) = \|x^*\| = \|Tx^*\|$. Every functional on B/Z attains its supremum on the unit sphere of B/Z consequently B/Z , as well as Z is reflexive. It follows that B is reflexive.

5. Rotundity and smoothness in quasi-reflexive spaces.

5.1. THEOREM. *If B is a quasi-reflexive Banach space, there exists an equivalent norm for B such that if B is either rotund or smooth in this norm, then B is reflexive.*

PROOF. By Theorem 3.5 [1] there exists an equivalent norm for B and quasi-reflexive Banach spaces $B^{(-5)}$, $B^{(-4)}$ and $B^{(-1)}$ such that $(B^{(-5)})^{****} = B^{(-1)}$, $(B^{(-4)})^{****} = B$ and $(B^{(-1)})^* = B$. By Theorem 20 [3], if B is rotund in this norm we must have $B^{(-4)}$ reflexive. It follows that B is reflexive. If B is smooth, then by [5, p. 12] $B^{(-1)}$ is rotund and so $B^{(-5)}$ is reflexive; B is therefore reflexive.

BIBLIOGRAPHY

1. Paul Civin and Bertram Yood, *Quasi reflexive spaces*, Proc. Amer. Math. Soc. vol. 8 (1957) pp. 906-911.
2. Mahlon M. Day, *Normed linear spaces*, Berlin, Springer-Verlag, 1958.
3. J. Dixmier, *Sur un théorème de Banach*, Duke Math. J. vol. 15 (1948) pp. 1057-1071.
4. Robert C. James, *Reflexivity and the supremum of linear functionals*, Ann. of Math. vol. 66 (1957) p. 159.
5. V. L. Klee, *Convex bodies and periodic homeomorphisms in Hilbert space*, Trans. Amer. Math. Soc. vol. 74 (1953) pp. 10-43.
6. R. R. Phelps, *Subreflexive normed linear spaces*, Arch. Math. vol. 8 (1957) pp. 444-450.

CANADIAN MATHEMATICAL CONGRESS, KINGSTON AND
UNIVERSITY OF SASKATCHEWAN