

ON RUNS OF RESIDUES¹

D. H. LEHMER AND EMMA LEHMER

According to a theorem of Alfred Brauer [1] all sufficiently large primes have runs of l consecutive integers that are k th power residues, where k and l are arbitrarily given integers. In this paper we consider the question of the first appearance of such runs.

Let p be a sufficiently large prime and let

$$r = r(k, l, p)$$

be the least positive integer such that

$$(1) \quad r, \quad r + 1, \quad r + 2, \dots, \quad r + l - 1$$

are all congruent modulo p to k th powers of integers > 0 . It is natural to ask, when k and l are given, how large is this minimum r and are there primes p for which r is arbitrarily large? If we let

$$\Lambda(k, l) = \limsup_{p \rightarrow \infty} r(k, l, p)$$

then is Λ infinite or finite, and if finite what is its value?

It is easy to see that

$$\Lambda(2, 2) = 9$$

so that every prime $p > 5$ has a pair of consecutive quadratic residues which appears not later than the pair (9, 10). In fact if 10 is not a quadratic residue of p then either 2 or 5 is, and so we have either (1, 2) or (4, 5) as a pair of consecutive residues.

By an elaboration of this reasoning M. Dunton has shown that

$$\Lambda(3, 2) = 77,$$

and more recently W. H. Mills has shown that

$$\Lambda(4, 2) = 1224.$$

Both of these proofs are as yet unpublished.

In contrast to these results we prove in this paper that

$$(2) \quad \Lambda(2, 3) = \infty,$$

and

Presented to the Society, November 19, 1960 under the title *On the distribution of consecutive quadratic and cubic residues*; received by the editors December 5, 1960.

¹ This paper is the result of unsupported research.

$$(3) \quad \Lambda(k, 4) = \infty, \quad k \leq 1048909.$$

In other words, by proper choice of p the appearance of a run of 3 quadratic residues or of 4 higher residues can be postponed as long as desired.

PROOF OF (2). Let N be a positive integer. Then it suffices to prove that there is a prime p for which

$$(4) \quad r(2, 3, p) > N.$$

Let

$$q_1, q_2, \dots, q_t$$

be all the primes $\leq N$.

By the quadratic reciprocity law, those primes which have a particular prime q_i as a quadratic residue belong to a set of arithmetic progressions of common difference $4q_i$. Those primes which have q_i for a nonresidue likewise belong to another set of arithmetic progressions of difference $4q_i$. If we combine the progressions of the first kind for every prime $q_i \equiv 1 \pmod{3}$ with those of the second kind for every prime $q_i \equiv 2 \pmod{3}$ and use Dirichlet's theorem on primes in arithmetic progressions we see that there exists a prime p such that

$$\left(\frac{q}{p}\right) \equiv q \pmod{3} \quad (q \neq 3, q \leq N).$$

Using the multiplicative property of Legendre's symbol we see that

$$(5) \quad \left(\frac{m}{p}\right) \equiv m \pmod{3} \quad (m \not\equiv 0 \pmod{3}, m \leq N).$$

But among any three consecutive numbers $\leq N$ there is one congruent to $-1 \pmod{3}$ and hence, by (5), is a nonresidue of this prime p . Hence the first run of three consecutive quadratic residues lies beyond N . This proves (4) and (2).

PROOF OF (3). The following theorem enables one to prove that for $l \geq 4$, $\Lambda(k, l) = \infty$ for all k up to high limits. It is clear that for such a program one may confine k to odd prime values and take $l=4$.

THEOREM A. *Let k and $p^* = kn+1$ be odd primes. Suppose further that 2 is not a k th power residue of p^* , and p^* is small enough so that it has no run of 4 consecutive k th power residues. Then $\Lambda(k, 4) = \infty$.*

For the proof we need the following lemma which is a special case of a theorem of Kummer [2].

LEMMA. Let k be an odd prime and q_1, q_2, \dots, q_t be any set of distinct primes different from k . Let $\gamma_1, \gamma_2, \dots, \gamma_t$ be a set of k th roots of unity. Then there exist infinitely many primes $p \equiv 1 \pmod{k}$ with corresponding k th power character χ modulo p such that

$$\chi(q_i) = \gamma_i \quad (i = 1(1)t).$$

To prove the theorem let N be an arbitrarily large integer and let q_1, q_2, \dots, q_t be the primes $\leq N$ with the exception of the prime p^* . Choosing a nonprincipal character, let γ_i be the k th power character of q_i modulo p^* . By the lemma there exist infinitely many primes $p \equiv 1 \pmod{k}$ such that the q 's have the same characters modulo p as modulo p^* . By the multiplicative property of characters this will be true of all the integers $m \leq N$ that are not divisible by p^* . Hence p has no run of 4 consecutive residues $\leq N$ unless one of these residues is a multiple of p^* . But two units on either side of this multiple of p^* we find numbers congruent to $\pm 2 \pmod{p^*}$ which are nonresidues of p^* and hence of p . Hence there is also no run of 4 residues which includes a multiple of $p^* \leq N$. This proves the theorem.

The fact that $\Lambda(3, 4) = \infty$ follows from the theorem by setting $k=3$ and $p^*=7$. Similarly by taking $k=5$ and $p^*=11$ we have $\Lambda(5, 4) = \infty$.

There is good reason to believe that $\Lambda(k, 4) = \infty$ for all k . To prove this it would suffice to prove for each prime k the existence of a prime $p^* = kn+1$ satisfying the hypothesis of the theorem. If n is not too large, then $p^* = kn+1$ will not have 4 consecutive k th power residues. In fact n is precisely the number of residues altogether. Trivially, if $n=2$ we have $\Lambda(k, 4) = \infty$ as with $k=3, 5, 11, 23$, etc. With a little more effort we can prove

THEOREM B. If $n \leq 12$ then $\Lambda(k, 4) = \infty$.

PROOF. We may suppose that $k > 5$. Let $p^* = kn+1$ be a prime not satisfying the hypothesis of Theorem A. This failure is not due to the fact that 2 is a residue of p^* . In fact if 2 were a residue, p^* would divide $2^n - 1$ by Euler's criterion. Since n is even and ≤ 12 this restricts p^* to the values

$$3, 5, 7, 11, 13, 17, 31.$$

In each case the corresponding value of k is ≤ 5 . Hence 2 must be a nonresidue along with -2 and $(p \pm 1)/2$. Hence we may suppose that p^* has a run of 4 residues

$$2 < a, a+1, a+2, a+3 < (p-1)/2$$

as well as the negatives of these modulo p^* . Besides these 8 residues there are the two residues congruent to $\pm(a+2)/a \not\equiv \pm 1$. These two are isolated since

$$\frac{a+2}{a} - 1 = \frac{2}{a} \quad \text{and} \quad \frac{a+2}{a} + 1 = \frac{2}{a}(a+1)$$

are obvious nonresidues. The reciprocals $\pm a/(a+2)$ are also isolated residues and they are new because

$$\frac{a+2}{a} \equiv -\frac{a}{a+2} \pmod{p^*}$$

implies

$$a(a+2) \equiv -2 \pmod{p^*}$$

in which a product of two residues is congruent to a nonresidue. Including the residues ± 1 we have accounted for at least 14 distinct k th power residues of p^* . Hence $14 \leq n \leq 12$, a contradiction. Therefore p^* must satisfy the hypothesis of Theorem A and so $\Lambda(k, 4) = \infty$.

A more elaborate argument involving the factors of $3^n - 1$ and the Fibonacci numbers yields a theorem in which the 12 in Theorem B is replaced by 36.

Let $p_0 = kn_0 + 1$ be the least prime congruent to 1 modulo k . Primes k for which $n_0(k) \geq 38$ are relatively rare, only about 3% of all the primes < 50000 by actual count. The least such prime is $k = 1637$ with $n_0 = 38$, and the largest value for n_0 for primes less than 50000 is $n_0 = 80$ for $k = 47303$. The values of $k < 50000$ were calculated on the SWAC and were tested on the 7090 by John Selfridge for pairs of consecutive k th power residues. It was discovered that in this range the only pairs are the trivial pairs $(\omega, \omega + 1)$ and $(\omega^2 \equiv p - \omega - 1, \omega^2 + 1 \equiv p - \omega)$, which appear whenever n_0 is a multiple of six. Since such pairs cannot obviously combine to make a quadruplet they were eliminated from the next run, made entirely on the 7090 by John Selfridge, for $k \leq 1048909$ in which no nontrivial pairs occurred. The largest value of $n_0 = 156$ occurred for $k = 707467$. These numerical results for which we are very grateful enable us to state the following theorem, using Theorem A.

THEOREM C. *If $k \leq 1048909$, then $\Lambda(k, 4) = \infty$.*

More generally one can ask about the first appearance of l consecutive numbers each with specified k th power character modulo $p = kn + 1$, excluding of course the case already considered in which all

the numbers are k th power residues. This seemingly more difficult problem is unexpectedly simple. Regardless of l the first appearance of such a set of consecutive numbers may be delayed indefinitely by proper choice of p . In fact if we set all the γ 's in the lemma at 1 we can find primes p having all the primes $\leq N$ and hence all the numbers $\leq N$ as k th power residues. Hence if the specified characters contain as much as a single nonresidue the first appearance can be made to occur beyond N .

In a future paper, written jointly with W. H. Mills, we determine the finite numbers $\Lambda(5, 2)$, $\Lambda(6, 2)$ and $\Lambda(3, 3)$.

REFERENCES

1. Alfred Brauer, *Über Sequenzen von Potenzresten*, S.-B. Deutsch. Akad. Wiss. Berlin 1928, 9–16.
2. D. Hilbert, *Die Theorie der algebraischen Zahlkörper*, Jber. Deutsch. Math. Verein. 4 (1897), 426.

UNIVERSITY OF CALIFORNIA, BERKELEY