ON A PROBLEM IN THE THEORY OF PARTITIONS¹

E. G. STRAUS

In this note we give an affirmative answer to a problem posed by Sherman K. Stein (Bull. Amer. Math. Soc. 66 (1960), 510). That is, we show that for every number m , $1 \leq m \leq p(n)$ where $p(n)$ is the number of partitions of the positive integer n , there exists a set $A(m, n)$ of positive integers such that n has m partitions into elements of $A(m, n)$. Our result generalizes to the case of partitions of vectors with nonnegative integral components.

DEFINITION 1. Let $S = \{s_1, s_2, \dots, s_k\}$ be a set of positive integers; then $p(S, n)$ denotes the number of partitions of n into elements of S . In particular we define $p(S, 0) = 1$ and $p(S, n) = 0$ for $n < 0$.

LEMMA 1.

$$p(S, n) = \sum_{i=1}^k p(S - S_i, n - s_i)$$

where $S_i = \{s_1, s_2, \dots, s_{i-1}\}$, $S_1 = \emptyset$. (Note: this lemma is independent of the ordering of S .)

PROOF. The i th term in the sum counts the number of partitions of n into elements of S where the first term occurring is s_i .

COROLLARY 1. If $1 \in S$ then $p(S, n)$ is a nondecreasing function of n .

COROLLARY 2.

$$p(S, n) \leq \sum_{l=0}^{n-1} p(S, l).$$

PROOF. By Lemma 1

$$\begin{aligned} p(S, n) &= \sum_{i=1}^k p(S - S_i, n - s_i) \\ &\leq \sum_{i=1}^k p(S, n - s_i) \\ &\leq \sum_{l=0}^{n-1} p(S, l). \end{aligned}$$

Received by the editors January 9, 1961.

¹ This work has been supported in part by a grant from the National Science Foundation.

COROLLARY 3. If $1 \in S$ then $p(S, n+1) \leq p(S, n) + p(S, n-1)$.

PROOF. We use simultaneous induction on n and k , the number of elements of S . If $n=0$ then

$$p(S, 1) \leq 1 = p(S, 0) = p(S, 0) + p(S, -1)$$

for all S . If $k=1$ then $S = \{1\}$ and

$$p(\{1\}, n+1) = 1 = p(\{1\}, n) \leq p(\{1\}, n) + p(\{1\}, n-1)$$

for all nonnegative n .

Now assume we had the minimal n and a minimal S for which the result is false then there is an $s \in S$, $s \neq 1$ and by Lemma 1

$$\begin{aligned} p(S, n+1) &= p(S, n+1-s) + p(S - \{s\}, n+1) \\ &\leq p(S, n-s) + p(S, n-1-s) + p(S - \{s\}, n) \\ &\quad + p(S - \{s\}, n-1) \\ &\leq p(S, n) + p(S, n-1). \end{aligned}$$

LEMMA 2. If $1 = a_1 \leq a_2 \leq \dots \leq a_k \leq \dots$ and $a_{k+1} \leq 2a_k$ where the a_k are integers, then for every integer n with $1 \leq n \leq a_1 + a_2 + \dots + a_k$ there exist indices $1 \leq i_1 < i_2 < \dots < i_j \leq k$ so that $n = a_{i_1} + \dots + a_{i_j}$.

PROOF. By induction on k . If $k=1$ then $n=1=a_1$. Assume the lemma proved for $k-1$. If $n \leq a_1 + \dots + a_{k-1}$ then the lemma is true by the induction hypothesis. If $n > a_1 + \dots + a_{k-1}$ then

$$n > a_k \left(\frac{1}{2} + \frac{1}{4} + \dots + \frac{1}{2^{k-1}} \right) = a_k \left(1 - \frac{1}{2^{k-1}} \right) \geq a_k - 1$$

so that $n \geq a_k$. Thus $0 \leq n - a_k \leq a_1 + \dots + a_{k-1}$ and the lemma holds, by the induction hypothesis.

LEMMA 3. If $1 \in S$ then the sequence $p(S, 1), p(S, 2), \dots, p(S, k), \dots$ satisfies the hypothesis of Lemma 2.

PROOF. We have $p(S, 1) = 1$ and $p(S, k+1) \geq p(S, k)$ by Corollary 1; finally $p(S, k+1) \leq 2p(S, k)$ by Corollary 3 of Lemma 1.

LEMMA 4. If $S \subset \{1, 2, \dots, [n/2]\}$ and $T \subset \{[n/2]+1, \dots, n\}$ then $p(S \cup T, n) = p(S, n) + \sum_{t \in T} p(S, n-t)$.

PROOF. Since $t+t' > n$ for any $t, t' \in T$ a partition of n into elements of $S \cup T$ is either a partition into elements of S alone or t plus a partition of $n-t$ into elements of S .

COROLLARY. If $1 \in S$ and $S \subset \{1, 2, \dots, [n/2]\}$ then for any integer m with

$$\begin{aligned}
 p(S, n) &\leq m \\
 &\leq p(S, n) + \sum_{l=0}^{[(n-1)/2]} p(S, l)
 \end{aligned}$$

there exists a set $T \subset \{[n/2] + 1, \dots, n\}$ so that $p(S \cup T, n) = m$.

LEMMA 5.

$$p\left(S \cup \left\{\left[\frac{n}{2}\right]\right\}, n\right) \leq 1 + p(S, n) + \sum_{l=0}^{[(n-1)/2]} p(S, l).$$

PROOF. If $[n/2] \in S$ the result is obvious. If $[n/2] \notin S$ then

$$\begin{aligned}
 &p\left(S \cup \left\{\left[\frac{n}{2}\right]\right\}, n\right) \\
 &= p(S, n) + p\left(S, n - \left[\frac{n}{2}\right]\right) + p\left(S, n - 2\left[\frac{n}{2}\right]\right) \\
 &\leq 1 + p(S, n) + p\left(S, \left[\frac{n-1}{2}\right] + 1\right)
 \end{aligned}$$

so that the lemma reduces to

$$p(S, m+1) \leq \sum_{l=0}^m p(S, l), \quad m = \left[\frac{n-1}{2}\right];$$

which is Corollary 2 of Lemma 1.

LEMMA 6. If $1 \in S$ and $S \subset \{1, 2, \dots, m-1\}$ where $m < [n/2]$ then

$$\begin{aligned}
 p(S \cup \{m\}, n) &\leq 1 + p\left(S \cup \left\{m+1, \dots, \left[\frac{n}{2}\right]\right\}, n\right) \\
 &\quad + \sum_{l=0}^{[(n-1)/2]} p\left(S \cup \left\{m+1, \dots, \left[\frac{n}{2}\right]\right\}, l\right) \\
 &= 1 + p(S \cup \{m+1, \dots, n\}, n).
 \end{aligned}$$

PROOF. The hypothesis implies $m \geq 2$. By repeated application of Lemma 1 we get

$$p(S \cup \{m\}, n) = \sum_{l=0}^{[n/m]} p(S, n - lm)$$

and by Corollary 3 of Lemma 1 together with Lemma 1

$$\begin{aligned}
& \sum_{l=0}^{\lfloor n/m \rfloor} p(S, n - lm) \\
& \leq p\left(S, n - \left\lfloor \frac{n}{m} \right\rfloor m\right) + p(S, n) + \sum_{l=1}^{\lfloor n/m \rfloor - 1} p(S, n - lm) \\
& \leq p\left(S, n - \left\lfloor \frac{n}{m} \right\rfloor m\right) + p(S, n) \\
& \quad + \sum_{l=1}^{\lfloor n/m \rfloor - 1} (p(S, n - lm - 1) + p(S, n - lm - 2)) \\
& \leq 1 + p(S, n) + \sum_{l=0}^{n-m-1} p(S, l) \\
& \leq 1 + p(S, n) + \sum_{k=1}^{n-m} p(S \cup \{m+1, \dots, m+k\}, n - m - k) \\
& = 1 + p(S \cup \{m+1, \dots, n\}, n).
\end{aligned}$$

THEOREM 1. *For every integer m , $1 \leq m \leq p(n)$, there exists a set $S \subset \{1, 2, \dots, n\}$ such that $p(S, n) = m$.*

PROOF. We construct a sequence $S_1, S_2, \dots, S_n$ of subsets of $\{1, 2, \dots, \lfloor n/2 \rfloor\}$ as follows:

(1) $S_1 = \{1\}$, $S_2 = \{1, \lfloor n/2 \rfloor\}$.

(2) Let r be the maximal element $\leq \lfloor n/2 \rfloor$ which is not contained in S_i then $S_{i+1} = S_i \cup \{r\} - \{r+1, \dots, \lfloor n/2 \rfloor\}$.

It is clear that this sequence terminates only with $S_N = \{1, 2, \dots, \lfloor n/2 \rfloor\}$. According to the corollary to Lemma 4 the numbers $p(S_i \cup T, n)$, where $T \subset \{\lfloor n/2 \rfloor + 1, \dots, n\}$, fill the interval I_i of numbers $p(S_i, n) \leq m \leq p(S_i \cup \{\lfloor n/2 \rfloor + 1, \dots, n\}, n)$. By Lemmas 5 and 6 the union $J_i = I_1 \cup I_2 \cup \dots \cup I_i$ is itself an interval so that J_N contains all m with $1 \leq m \leq p(n)$.

One might ask whether our theorem has an analogue if $p(n)$ is replaced by $p(S, n)$, that is whether for any m with $1 \leq m \leq p(S, n)$ there is a subset S' of S so that $p(S', n) = m$. The answer is obviously no, since $p(\{1, 2\}, n) = \lfloor n/2 \rfloor + 1$ while $p(\{1\}, n) = 1$ and $p(\{2\}, n) \leq 1$.² It is, however, possible to generalize our result to the partition of vectors.

DEFINITION. Let $n = (n_1, n_2, \dots, n_k)$ where the n_i are integers; then $p_k(n)$ is the number of partitions of n into k -vectors whose components are nonnegative integers not all of which are zero.

THEOREM 2. *Let m be an integer, $1 \leq m \leq p_k(n)$ then there exists a set*

² The answer is yes if we let $S = \{m, m+1, m+2, \dots\}$ be the set of all integers $\geq m$, m a positive integer, without appreciable change in the proof.

S of nonnegative nonzero k-vectors such that the number $p_k(S, n)$ of partitions of n into elements of S is m .

The proof involves only simple modifications of the steps which led to Theorem 1. We outline them here. Lemma 1 remains valid. In Corollary 1 the condition $1 \in S$ is to be replaced by the condition that S contains all the unit-vectors. By "nondecreasing" we now refer to the partial ordering given by $a < b$ if $b - a$ has nonnegative components. In Corollary 2 the summation is understood to be over all vectors which precede n in the above sense.

In Corollary 3 the vectors $n+1$ and $n-1$ must be understood as the result of adding and subtracting from n the same (arbitrary) unit vector. We need a slight modification of this result:

If S contains all unit vectors and $n+1 = n' + 1'$ where 1 and $1'$ are two different unit vectors then

$$p_k(S, n+1) \leq p_k(S, n) + p_k(S, n').$$

Instead of the subsequent lemmas we get

LEMMA 3'. *If S contains the unit vectors then the set $\{p_k(S, l)\}$ where l ranges over all nonnegative vectors with $l < n$ can be arranged as a non-decreasing sequence which satisfies the hypothesis of Lemma 2.*

LEMMA 4'. *If $S \cap T = \emptyset$ and $2t > n$ for every $t \in T$, then*

$$p_k(S \cup T, n) = p_k(S, n) + \sum_{t \in T} p_k(S, n - t).$$

COROLLARY. *Let S contain the unit vectors and $S \cap T = \emptyset$ where T is the set of all vectors t which satisfy $2t > n$. Then for every integer m with $p_k(S, n) \leq m \leq p_k(S \cup T, n)$ there exists a $T' \subset T$ so that $p_k(S \cup T', n) = m$.*

LEMMA 5'. *Let $n^{(i)} = (n_1, n_2, \dots, [n_i/2], \dots, n_k)$ and let T be as in the preceding corollary. Then*

$$p_k(S \cup \{n^{(i)}\}, n) \leq 1 + p_k(S \cup T, n).$$

LEMMA 6'. *Let T be as above and let S contain the unit vectors with $S \cap T = \emptyset$. Let $m \in S \cup T$ be a vector all whose successors are contained in $S \cup T$. Let S_m denote the set of successors of m . Then*

$$p_k(S \cup \{m\} - S_m, n) \leq 1 + p_k(S \cup T, n).$$

None of the proofs involves any new ideas and we therefore omit them. Theorem 2 now follows by exactly the same argument that led to Theorem 1.