A NOTE ON HOMOGENEOUS COMPLEX CONTACT MANIFOLDS

WILLIAM M. BOOTHBY¹

1. Introduction. A complex contact manifold is by definition a complex analytic manifold of odd dimension² $2n+1$ over which is defined to within an analytic scalar multiple an analytic pfaffian form ω of class $2n+1$, i.e. such that $\omega \wedge d\omega^n \neq 0$ at any point. Such a manifold is given [4] by a covering by coordinate neighborhoods $\{U_i\}$ on each of which is defined a form ω_i of the type described with $\omega_i = f_{ij}\omega_j$ whenever $U_i \cap U_j \neq \emptyset$, f_{ij} being nonvanishing, analytic scalar functions. A natural question arises as to whether the contact form may be chosen globally, that is the ω_i so chosen that each $f_{ij} \equiv 1$. It was shown by S. Kobayashi [4] that this can be done if and only if the first Chern class, $c_1(M)$, vanishes. Here and in what follows we will suppose M to be compact. Under the additional assumption that M is simply connected and homogeneous both with respect to the complex and the contact structure the author [2] characterized and enumerated those manifolds for which $c_1(M) \neq 0$. In what follows it is shown that these additional assumptions already imply $c_1(M) \neq 0$, so that combining this with the results of [2] we obtain:

THEOREM I. *Let M be a compact, simply connected, homogeneous complex contact manifold containing more than one point. Then $c_1(M) \neq 0$, hence ω can not be globally defined; M is Kählerian; and there is exactly one such manifold corresponding to each of the classes of simple Lie groups A_n , B_n , C_n , and D_n and the five exceptional simple groups. No other manifolds satisfying these hypotheses exist.*

When we make no homogeneity assumption, the classical example of such a manifold is the bundle of complex "co-directions" over a complex analytic manifold of dimension $n+1$: the fibres are complex projective spaces of dimension n , and with local coordinates $z^1, \dots, z^{n+1}$ in the base space, $p_1, \dots, p_{n+1}$ (homogeneous) coordinates in the fibre we have $\omega = p_1 dz^1 + \dots + p_{n+1} dz^{n+1}$. In this connection the question arises as to whether the spaces of Theorem I are already included in this example. This question is answered by the following:

Received by the editors March 13, 1961.

¹ The author was supported by a grant from the National Science Foundation.

² Throughout dimension refers to complex dimension and analytic means complex analytic.

THEOREM II. *With the exception of the manifolds corresponding to A_n , none of the manifolds of Theorem I can be homeomorphic to a bundle of complex "co-directions" over a complex manifold, i.e. to an example of the type described above.*

2. Proof of Theorem I. Following H. C. Wang [5], we will refer to a compact, simply connected, homogeneous complex manifold as a C -space. The notation used in this section is that of [5] and [2] and for brevity will not be redefined. Suppose G/L to be a C -space with a globally defined contact form ω which is G -invariant. Without loss of generality we suppose G to be complex semi-simple. By proceeding as in the real case [3, §4] we will arrive at a contradiction which shows the impossibility of a globally defined contact form. This together with the results of [2] gives Theorem I. Let $K = \{g \in G \mid \text{ad}(g)^* \omega^* = \omega^*\}$ where $\text{ad}(g)^*$ denotes the transformation on forms on the Lie algebra $\hat{G}$ induced by the inner automorphism $\text{ad}(g): G \rightarrow G$ and $\omega^* = p^* \omega$, $p: G \rightarrow G/L$ being the natural map. Clearly K is a closed subgroup and contains L . As in the real case we see $\hat{K} = \{X \in \hat{G} \mid d\omega^*(X, \hat{G}) = 0\}$ and therefore $\hat{K}$ is a complex Lie algebra; hence K is a closed, complex subgroup and G/K a homogeneous complex manifold. There is a natural analytic map of G/L onto G/K so we know G/K to be compact and connected. If K_0 is the identity component of K , then the same statements apply to K_0 and G/K_0 provided L is connected, which is, in fact, the case since G is connected and G/L is simply connected. Then from the homotopy sequence of $G/L \rightarrow G/K_0$ it follows that G/K_0 is simply connected and hence is a C -space. It contains more than a single point for, as in the real case, $\omega \wedge d\omega^n \neq 0$ implies $\dim \hat{K} = 1 + \dim \hat{L}$ and, if G/K_0 is a single point, then $K_0 = G$ and K_0/L is an abelian group of complex dimension one. Since it is compact and simply connected, it too contains but one point which is contrary to assumption.

Now let $Z \in \hat{G}$ correspond to ω^* relative to the Killing form (X, Y) on $\hat{G}$. Then $\omega^*(X) = (Z, X)$ and $d\omega^*(X, Y) = \omega([X, Y])/2$. Thus $d\omega^*(X, Y) = 0$ if and only if $([Z, X], Y) = (Z, [X, Y]) = 0$. Since the Killing form is nondegenerate, the first of these vanishes for all $Y \in \hat{G}$ if and only if $[X, Z] = 0$, therefore $\hat{K} = \hat{C}(Z)$, the centralizer of Z , i.e. $\hat{C}(Z) = \{X \mid [X, Z] = 0\}$. The desired contradiction is then obtained from the following:

LEMMA. *Let G be a connected, complex, semi-simple Lie group and K a closed, connected, complex subgroup whose Lie algebra $\hat{K}$ is the centralizer of some $Z \in \hat{G}$. Then, if G/K is a C -space, it contains only one point.*

PROOF. According to Wang [5], if G/K is a C -space, then for the

Lie algebras we have the following situation. There exists a Cartan subalgebra $\hat{H} \subset \hat{G}$ with rational basis $h_1, \dots, h_b, h_{b+1}, \dots, h_a, h_{a+1}, \dots, h_l$, with $b = 2u$, where $\hat{K}$ has as basis $k_1, \dots, k_u, h_{b+1}, \dots, h_a, h_{a+1}, \dots, h_l; E_{\pm\alpha_1}, \dots, E_{\pm\alpha_s}, E_{\sigma_1}, \dots, E_{\sigma_r}$, where $k_1 = ih_1 + ch_2$, $k_2 = ih_3 + ch_4, \dots, k_u = ih_{b-1} + ch_b$, c a real number, and $\alpha_1, \dots, \alpha_s, \sigma_1, \dots, \sigma_r$ are all positive roots and $\pm\alpha_1, \dots, \pm\alpha_s$ are all those roots which vanish on each of the vectors $h_1, \dots, h_a$. The maximal semi-simple subalgebra $\hat{Q}$ of $\hat{K}$ is spanned by $h_{a+1}, \dots, h_l$ and $E_{\pm\alpha_1}, \dots, E_{\pm\alpha_s}$. Since $Z \in \hat{K}$, we may write with suitable $h \in \hat{H} \cap \hat{K}$, $Z = h + \sum_{i=1}^s a_i E_{\alpha_i} + \sum_{i=1}^s b_i E_{-\alpha_i} + \sum_{j=1}^r d_j E_{\sigma_j}$. We shall prove $Z = 0$, whence $\hat{G} = \hat{K}$ and G/K is a single point. If $X \in \hat{K}$ is either in $\hat{H}$ or a root vector E_r , then since $[X, Z] = 0$, each summand of $[X, h] + \sum_{i=1}^s a_i [X, E_{\alpha_i}] + \sum_{i=1}^s b_i [X, E_{-\alpha_i}] + \sum_{j=1}^r d_j [X, E_{\sigma_j}]$ vanishes since each lies in a different summand of a direct sum decomposition of $\hat{G}$. Thus $[E_\sigma, h] = \sigma(h)E_\sigma$ vanishes for each positive root σ , hence $h = 0$. Multiplying by $E_{+\alpha}$ or $E_{-\alpha}$ we obtain $a_i = 0 = b_i$ for each i . Finally let σ_{j_0} be one of the roots $\sigma_1, \dots, \sigma_r$ and $k, 1 \leq k \leq a$, be such that $\sigma_{j_0}(h_k) \neq 0$. There must be at least one such or σ_{j_0} would be among the $\pm\alpha_i$. If $b+1 \leq k \leq a$, then, since $h_k \in \hat{K}$ and thus $[h_k, Z] = 0$, we at once get $d_{j_0}[h_k, E_{\sigma_{j_0}}] = \sigma_{j_0}(h_k)d_{j_0}E_{\sigma_{j_0}} = 0$ and thus $d_{j_0} = 0$. But, if $1 \leq k \leq b$, then either $[ih_k + ch_{k+1}, Z]d_{j_0} = 0$ or $[ih_{k-1} + ch_k, Z]d_{j_0} = 0$ according to the parity of k . In either case $d_{j_0} = 0$ unless the term in brackets vanishes. But this cannot happen since, say in the first case, vanishing of the term in brackets would imply $i\sigma_{j_0}(h_k) + c\sigma_{j_0}(h_{k+1}) = 0$ and thus since σ_{j_0} is real on h_k and h_{k+1} , $\sigma_{j_0}(h_k) = 0 = \sigma_{j_0}(h_{k+1})$, contrary to our choice of h_k . A corresponding argument applies to the other case and to each of the $\sigma_j, j = 1, \dots, r$. Thus each $d_j = 0$ and so $Z = 0$ as was claimed.

3. Proof of Theorem II. We now consider the possibility that the homogeneous complex contact manifold $M = G/L$ might be a bundle of complex co-directions over a complex analytic manifold B of dimension $n+1$. In this case the fibre F would be a complex projective space of dimension n . By Theorem I M is a Kähler manifold, F is also Kählerian, and by Blanchard [1, Prop. II.2, p. 184] we see that B is Kählerian. Concerning the topology of M, F , and B we use the fact that $H^1(F) = 0$, which implies that the transgression $H^1(F) \rightarrow H^2(B)$ vanishes, together with the fact that $\pi_1(B) = 0$, which follows from the fact that F and M are both simply connected spaces and, of course, implies that $\pi_1(B)$ acts trivially on the cohomology groups of F . This allows us to use Blanchard, op. cit., Theorem II 1.2, p. 178, according to which the real cohomology of M is isomorphic

to that of $B \times F$. In particular for the Poincaré polynomials of M , F , and B we have

$$P_M(t) = P_F(t)P_B(t) = (1 + t^2 + \cdots + t^n)(1 + at^2 + \cdots + t^{n+1})$$

where $a \geq 1$ since B is Kählerian. It follows that the second Betti number of M , $p_2(M) \geq 2$. Now let $M = G/L$, G complex semi-simple as before. It is known that $\pi_2(G) = 0$. The homotopy sequence of the bundle $G \rightarrow G/L$ yields the exact sequence:

$$0 \rightarrow \pi_2(G/L) \rightarrow \pi_1(L) \rightarrow \pi_1(G) \rightarrow 0,$$

which, in particular, says that the rank of $\pi_2(G/L)$ is no greater than the rank of $\pi_1(L)$. Since G/L is simply connected, $\pi_2(G/L) \approx H^2(G/L)$; thus $p_2(M) \leq \text{rank } \pi_1(L)$. Now examining the individual cases in Theorem I and replacing G and L by their maximal compact subgroups G' , L' , cf. [2], shows that $L' = A \times T^r$ where A is semi-simple and thus $\text{rank } \pi_1(L) = r$. Since $2 \leq p_2(M) \leq r$ we must have $r \geq 2$, but this occurs only in the case corresponding to A_n where $G' = SU(n+1)$ and $L' = SU(n-1) \times T^2$.

4. Remarks. I. Since every *homogeneous* complex contact manifold, at least if it is simply connected, is a Kähler manifold, one might ask if this is true in general. However, from Blanchard, op. cit., we see that if B is a non-Kählerian, compact, complex analytic manifold then no bundle over it can be Kählerian. Thus the bundle of co-directions furnishes an example of a non-Kählerian compact complex contact manifold. It will be simply connected if B is chosen simply connected.

II. In the one exceptional case of Theorem II we have $SU(n+1) \supset SU(n) \times T^1 \supset SU(n-1) \times T^2$ so that denoting $SU(n+1)/SU(n-1) \times T^2$ by M , $SU(n+1)/SU(n) \times T^1$ by B and $SU(n) \times T^1/SU(n-1) \times T^2$ by F we see that M is a bundle over B whose fibre F is a projective space $P_{n-1}(C)$ of dimension $n-1$ and whose base B is a projective space $P_n(C)$ of dimension n . And in fact the bundle is homeomorphic to the bundle of tangent co-directions to n dimensional complex projective space.

A proof of this last statement may be sketched as follows. We regard $P_n(C)$ as the space of directions through the origin in C_{n+1} , $n+1$ dimensional Hermitian space, and the complex n -plane orthogonal to a given direction as the tangent space to the corresponding point of $P_n(C)$. Then $SU(n+1)$ is transitive on the pairs $(x, d) = (\text{point } x, \text{tangent direction at } x)$ of $P_n(C)$. The subgroups fixing respectively x and (x, d) are $SU(n) \times T^1$ and $SU(n-1) \times T^2$. Then

the result follows easily if we note that the space of tangent co-directions to $P_n(C)$ is homeomorphic to the space of tangent directions as we may see by choosing a Hermitian metric on $P_n(C)$.

BIBLIOGRAPHY

1. A. Blanchard, *Sur les variétés analytiques complexes*, Ann. Sci. École Norm. Sup. (3) **73** (1956), 157–202.
2. W. M. Boothby, *Homogeneous complex contact manifolds*, Proc. Sympos. Pure Math. Vol. 3, pp. 144–154, Amer. Math. Soc., Providence, R. I., 1961.
3. W. M. Boothby and H. C. Wang, *On contact manifolds*, Ann. of Math. **68** (1958), 721–734.
4. S. Kobayashi, *Remarks on complex contact manifolds*, Proc. Amer. Math. Soc. **10** (1959), 164–167.
5. H. C. Wang, *Closed manifolds with homogeneous complex structures*, Amer. J. Math. **76** (1953), 1–32.

NORTHWESTERN UNIVERSITY