9. W. Rudin, *Measure algebras on abelian groups*, Bull. Amer. Math. Soc. **65** (1959), 227-247.

10. G. E. Šilov, *On decomposition of a commutative normed ring in a direct sum of ideals*, Mat. Sb. **32** (1953), 353-364.

11. A. Weil, *L'intégration dans les groupes topologiques et ses applications*, Actualités Sci. Ind. no. 869, Hermann, Paris, 1940.

12. J. Wermer, *Ideals in a class of commutative Banach algebras*, Duke Math. J. **20** (1953), 273-278.

OHIO STATE UNIVERSITY AND
UNIVERSITY OF NOTRE DAME

A MULTI-POINT GENERALIZATION OF NEWTON'S DIVIDED DIFFERENCE FORMULA

HERBERT E. SALZER

Newton's divided difference formula for the polynomial of degree $n-1$ determined by the values of the function $f(x)$ and its divided differences up to the $(n-1)$ th order originating from any single point, has the following multi-point generalization:

THEOREM I. *The unique interpolation polynomial $P(x)$, of degree $\sum_{i=1}^n r_i - 1$, which is determined by $f(x_i) \equiv f_i \equiv \mathcal{D}^0 f_i$ and the first $r_i - 1$ divided differences, $\mathcal{D}^m f_i \equiv [x_i x_{i_1} \cdots x_{i_m}]$, $m = 1, \cdots, r_i - 1$, originating from the n points x_i , $i = 1, \cdots, n$, is expressible as*

$$(1) \quad P(x) = \sum_{i=1}^n \Pi_i(x) \psi_i(x), \text{ where}$$

$$(2) \quad \Pi_i(x) = \prod_{j=1, j \neq i}^n (x - x_j)(x - x_{j_1}) \cdots (x - x_{j_{r_j-1}}),$$

$$(3) \quad \psi_i(x) = \sum_{m=0}^{r_i-1} [x_i x_{i_1} \cdots x_{i_m}] \psi_{i,m}(x), \text{ and}$$

$$(4) \quad \psi_{i,m}(x) = \sum_{s=0}^{r_i-m-1} [x_{i_m} x_{i_{m+1}} \cdots x_{i_{m+s}}]_{1/\Pi_i} (x - x_i)(x - x_{i_1}) \cdots (x - x_{i_{m+s-1}}).$$

In (3) and (4), $x_{i_0} \equiv x_i$; in (3), $[x_i] \equiv f_i$; in (4), $[x_{i_m}]_{1/\Pi_i} \equiv 1/\Pi_i(x_{i_m})$, the s th order divided difference $[x_{i_m} x_{i_{m+1}} \cdots x_{i_{m+s}}]_{1/\Pi_i}$ is for the function $1/\Pi_i(x)$, and $\psi_{i,0}(x)$ begins $[x_i]_{1/\Pi_i} + \cdots$.

Presented to the Society, April 22, 1961; received by the editors February 23, 1961.

The quantities in braces in (7) are

$$[x_{i_m} \cdots x_{i_{m+s}}]_{\Pi_i}, \cdots, [x_{i_{m+s-1}} x_{i_{m+s}}]_{\Pi_i}.$$

We now employ the Popoviciu-Steffensen theorem for the m th divided difference of the product $f(x)\phi(x)$ in terms of the divided differences of $f(x)$ and $\phi(x)$ [3; 4]:¹

$$(8) \quad [x_0 x_1 \cdots x_m]_{f\phi} = \sum_{i=0}^m [x_0 \cdots x_i]_f [x_i \cdots x_m]_{\phi}.$$

Applying (8) to the product $1 \equiv \{1/\Pi_i(x)\}\Pi_i(x)$ for the points $(x_{i_m}, x_{i_{m+1}}, \cdots, x_{i_{m+s}})$, we have

$$(9) \quad \begin{aligned} 0 &= (1/\Pi_i(x_{i_m})) [x_{i_m} \cdots x_{i_{m+s}}]_{\Pi_i} + \cdots \\ &+ [x_{i_m} \cdots x_{i_{m+s-1}}]_{1/\Pi_i} [x_{i_{m+s-1}} x_{i_{m+s}}]_{\Pi_i} \\ &+ [x_{i_m} \cdots x_{i_{m+s}}]_{1/\Pi_i} \Pi_i(x_{i_{m+s}}). \end{aligned}$$

Comparison of (9) with (7) shows $a_{i_m}^{(s)} = [x_{i_m} \cdots x_{i_{m+s}}]_{1/\Pi_i}$.

The uniqueness of $P(x)$ is apparent, since specification by $f_i, \mathfrak{D}f_i, \cdots, \mathfrak{D}^{r_i-1}f_i, i=1, \cdots, n$, is equivalent to specification by certain $\sum_{i=1}^n r_i$ functional values $f_i, f_{i_1}, \cdots, f_{i_{r_i-1}}$ at $x_i, x_{i_1}, \cdots, x_{i_{r_i-1}}$, for which the Lagrange interpolation polynomial of degree $\sum_{i=1}^n r_i - 1$ (a rearrangement of $P(x)$) is unique.

REFERENCES

1. H. E. Salzer, *Hermite's general osculatory interpolation formula and a finite difference analogue*, J. Soc. Indust. Appl. Math. 8 (1960), 18-27.
2. L. M. Milne-Thomson, *The calculus of finite differences*, Macmillan, London, 1933, p. 7.
3. T. Popoviciu, *Sur quelques propriétés des fonctions d'une ou de deux variables réelles*, Mathematica 8 (1934), 1-85.
4. J. F. Steffensen, *Note on divided differences*, Danske Vid. Selsk. Math.-Fys. Medd. 17, no. 3 (1939), 12 pp.

GENERAL DYNAMICS/ASTRONAUTICS

¹ Popoviciu published a statement of (8) without proof in his thesis [3], apparently unknown to Steffensen whose proof appeared 5 or 6 years later [4].