UPPER SEMICONTINUOUS DECOMPOSITIONS OF THE n -SPHERE

KYUNG WHAN KWUN

We consider conditions under which an upper semicontinuous decomposition has the decomposition space which is a topological n -sphere. A special emphasis is placed on the case in which the decomposition has only a countable number of nondegenerate elements. If M is an n -manifold with boundary, $\text{Bd } M$ and $\text{Int } M$ will be used to denote the usual boundary and the interior of M . The set theoretical boundary of a subset A in a space will be denoted by $b(A)$ and the closure of A by $\text{Cl } A$.

1. *I-sets in S^n .* A subset F of the n -sphere S^n is called an *I-set* if $S^n - F$ is homeomorphic to $S^n - A$ for some countable subset A of S^n .

LEMMA 1. *Let A be a countable set in n -space E^n , $n > 1$. Then $E^n - A$ is connected.*

LEMMA 1'. *If X is a connected n -manifold with boundary, $n > 1$, $X - A$ is connected for any countable set A .*

THE PROOF OF LEMMAS 1 AND 1'. If A is dense in E^n , Lemma 1 follows from Proposition A, p. 44, of [5] and the general case can be proved using this special case. For the proof of Lemma 1', use Lemma 1.

LEMMA 2. *Each component of an I-set F in S^n , $n > 1$, is closed.*

PROOF. Suppose a component C of F is not closed. Let $x \in \text{Cl } C - C$. Let h be a homeomorphism of $S^n - F$ onto $S^n - A$ for some countable set A . Since A is countable, there exist n -cells $D_1, D_2, \dots$ such that $\text{Int } D_i \supset D_{i+1}$, $\text{Bd } D_i \cdot A = \emptyset$ for each i and the intersection of the D_i is the point $h(x)$. Let U_i be the complementary domain of $S_i = h^{-1}(\text{Bd } D_i)$ containing x and U_i' the other complementary domain. Then $(U_i - F) + (U_i' - F)$ is a disjoint sum equal to a disjoint sum $h^{-1}(\text{Int } D_i - A) + h^{-1}(S^n - A - D_i)$. Since the four sets are open in $S^n - S_i - F$ and the last two sets are connected by Lemma 1', it follows that $U_i - F = h^{-1}(\text{Int } D_i - A)$. Each U_i contains $\text{Cl } U_{i+1}$ and the intersection U of the U_i is a compact set contained in $F + x$ as the intersection of D_i is $h(x)$. Since C is connected, U_i is open and S_i is disjoint from F , each U_i contains C . Since C is closed in F , it follows that $\text{Cl } C = C + x$. Let $y \in b(C) - x$. (It is impossible that $b(C) = x$.)

Received by the editors March 16, 1961.

Let $y_1, y_2, \dots$ be a sequence of points in $S^n - F$ that converges to y . Let $h(y_{i_1}), h(y_{i_2}), \dots$ be a converging subsequence of $h(y_i)$. Then $y' = \lim h(y_{i_j})$ is on A . Let E_i be n -cells such that $\text{Int } E_i \supset E_{i+1}$, $\text{Bd } E_i \cdot A = \emptyset$ for each i and the intersection of E_i is y' . Let V_i be the complementary domain of $S'_i = h^{-1}(\text{Bd } E_i)$ containing y . Then $V_i - F = h^{-1}(\text{Int } E_i - A)$ as V_i contains infinitely many y_j . Then each Cl V_i must contain Cl C for the same reason as before, but this is impossible as the intersection of E_i is disjoint from $h(x)$.

Let F be any subset whatsoever of S^n . By $G(F)$ we denote the collection of the components of F and the points of $S^n - F$.

THEOREM 1. *Let G be a decomposition of S^n , $n > 1$, having only a countable number of nondegenerate elements. In order that G be an upper semicontinuous decomposition of S^n having an n -sphere as decomposition space, it is necessary and sufficient that $G = G(F)$ for some I -set F in S^n .*

PROOF. *The necessity.* Let g be an arbitrary element of G and X and f respectively denote the decomposition space and the quotient map of G . Let $D_1, D_2, \dots$ be n -cells in X such that $\text{Bd } D_i$ do not meet the image under f of any nondegenerate element of G and the intersection of D_i is $f(g)$. Then g is the intersection of a decreasing sequence of compact connected sets U_i , where U_i are the closures of the complementary domains of $f^{-1}(\text{Bd } D_i)$ containing g . Hence g is connected. Then $G = G(F)$ for the sum F of the nondegenerate elements of G .

The sufficiency. We first show that $G(F)$ is upper semicontinuous. Let $x \in S^n - F$. It is easy to see that x has an arbitrarily small neighborhood which is the sum of elements of $G(F)$. (Recall the proof of Lemma 2.) Let C be a component of F and h be a homeomorphism of $S^n - F$ onto $S^n - A$ for some countable subset A of S^n . Let $x_1, x_2, \dots$ be a sequence of points in $S^n - F$ that converges to a point $x \in C$.¹ Then by an argument similar to one in the proof of Lemma 2, we see that C is contained in the intersection of a decreasing sequence of compact connected sets Cl V_i , where V_i is a complementary domain of an $(n-1)$ -sphere disjoint from F , and the intersection of Cl V_i is contained in F . That the intersection in fact is C follows from the fact that C is a component of F and the intersection is connected. Then a V_i with a sufficiently large i is a very close neighborhood of C which is the sum of elements of $G(F)$.

We now define a map f of S^n onto itself as follows. For each point $x \in S^n - F$, we let $f(x) = h(x)$. For each component C of F , let $y_1, y_2, \dots$ be a sequence of points in $S^n - F$ that converges to a point $y \in C$. Let $h(y_{i_1}), h(y_{i_2}), \dots$ be a subsequence of $h(y_i)$ that con-

¹ The existence of x is assured by the closedness of C . (See Lemma 1.)

verges to a point y' . Then $y' \in A$ as y_i do not have a limit point in $S^n - F$. Let $f(C) = y'$. The map f is then well-defined. For if y' depended on y , etc. and in fact had two y' 's, by an argument similar to the proof of Lemma 2, we would find C contained in two disjoint neighborhoods. We can also prove that f is onto and sends distinct elements of $G(F)$ into distinct points. To prove the continuity of f , let D be an n -cell in S^n such that $\text{Bd } D$ is disjoint from A . Then $f^{-1}(\text{Int } D)$ is a complementary domain of $f^{-1}(\text{Bd } D) = h^{-1}(\text{Bd } D)$ which is certainly open. Since each open set in S^n is the sum of such $\text{Int } D$'s, f is continuous.

COROLLARY 1. *An I -set F in S^n , $n > 1$, has only a countable number of components.*

COROLLARY 2. *Let A and B be two countable subsets of S^n , $n > 1$. Then there exists a homeomorphism h of S^n onto itself such that $h(A) = B$ if and only if there is a homeomorphism h' of $S^n - A$ onto $S^n - B$. Moreover, h can be taken as an extension of h' .*

REMARK. Counter-examples show that Lemmas 1, 1' and 2, Theorem 1 and Corollaries 1 and 2 are false for $n = 1$ and that in the statement of Corollary 2 the words "countable subsets" cannot be replaced by "arcs."

THEOREM 2. *Let G be an upper semicontinuous decomposition of S^3 having a countable number of nondegenerate elements. If the decomposition space X is a topological 3-sphere, then each element of G has an open 3-cell as complement.*

PROOF. Let f be the quotient map of G . We denote by F the sum of the nondegenerate elements of G . Let g be an arbitrary element of G . Since $f(F)$ is countable, there exist 3-cells $D_1, D_2, \dots$ in X such that $\text{Int } D_i \supset D_{i+1}$, $\text{Bd } D_i \cdot f(F) = \emptyset$ for each i and the intersection of D_i is $f(g)$. Let $U_i = f^{-1}(\text{Int } D_i)$. Then U_i is the complementary domain of $S_i = f^{-1}(\text{Bd } D_i)$ containing g . By the approximation [2] of Bing and the lemma below there exists a polyhedral 2-sphere S'_i separating $\text{Cl } U_{i+1}$ from $S^3 - U_{i-1}$ for each $i > 1$. Hence there exist polyhedral 3-cells $E_1, E_2, \dots$ such that $\text{Int } E_i \supset E_{i+1}$ for each i and the intersection of E_i is f . A repeated application of Theorem 1 of [6] proves the theorem.

LEMMA 3. *Let S be an $(n-1)$ -sphere in S^n separating two connected closed subsets A and B in $S^n - S$. Then there exists a positive number ϵ such that if h is an ϵ -homeomorphism of S onto $h(S)$ in $S^n - A - B$ then $h(S)$ separates A from B .*

PROOF. We first show that there is a positive number ϵ such that if h is an ϵ -homeomorphism mentioned in the lemma, then there is a homotopy connecting the inclusion map of S into S^n and the map h . Since a neighborhood of S can be uniformly continuously imbedded in E^n , we may assume that S is in E^n and construct a homotopy involving a very close neighborhood of S . Let $D_1, D_2, \dots, D_k$ be a finite number of convex n -cells such that $\sum \text{Int } D_i \supset S$. Let T be a triangulation of S so fine that each simplex Δ of T lies in an $\text{Int } D_{i(\Delta)}$. Let ϵ be a positive number such that each ϵ -neighborhood of each simplex Δ of T lies in $\text{Int } D_{i(\Delta)}$. Let h be an ϵ -homeomorphism of S onto $h(S)$. Notice that $h(\Delta)$ lies in $\text{Int } D_{i(\Delta)}$. Then it is easy to construct the desired homotopy within $\sum \text{Int } D_i$. Now we come back to S^n and suppose that ϵ was chosen such that not only h has the property mentioned at the beginning of the proof but also the homotopy takes place outside the sum of an open connected neighborhood U of A and an open connected neighborhood V of B . Before proceeding any further, we refer the reader to [7, Definition 8.2 and Theorem 8.3, p. 266]. Let $M = S^n - U - V$. A cycle Γ irreducibly carried by S and a cycle γ irreducibly carried by the sum of a point of A and a point of B are linked. Since Γ is homologous to $h(\Gamma)$ in M , $h(\Gamma)$ and γ are also linked. Since A and B are connected and disjoint from $h(S)$, $h(S)$ must separate A from B .

REMARKS. A result for S^2 that is stronger than Theorem 2 is well known. While it may be true that an analogous theorem holds in S^n , we have no argument to prove it.

Suppose F is a subset of S^3 having a countable number of components of which each has an open 3-cell complement. In general, F is not an I -set. However, we can prove

THEOREM 3. *Let F be a subset of S^3 having only a countable number of components. Suppose F is a G set and $G(F)$ is upper semicontinuous. Then F is an I -set if and only if each component of F has the open 3-cell complement.*

PROOF. The "if" part follows from Theorem 1 and [1, Theorem 1]. The "only if" part follows from Theorems 1 and 2.

THEOREM 4. *Let F be the sum of a countable number of disjoint tame arcs in S^3 . Then F is an I -set if and only if $G(F)$ is upper semicontinuous.*

PROOF. Theorem 1 and [1, Theorem 3].

2. **Similar n -cells in S^n .** Let S^{n-1} denote the "equator" of S^n . A topological $(n-1)$ -sphere S in S^n is called *tame* if there exists a homeo-

morphism of S^n onto itself that sends S onto S^{n-1} . A result [3] of Brown implies that S is tame if and only if S has a product neighborhood, i.e., the image of a homeomorphism of $S^{n-1} \times [0, 1]$ into S^n that sends $S^{n-1} \times 1/2$ onto S . If S is tame, its product neighborhood can be taken in any given neighborhood of S . An n -cell D in S^n is called *tame* if $\text{Bd } D$ is tame. Let $R_i, i = 1, 2$, be the subsets of E^n consisting of the points x with $\|x\| \leq i$. Two n -cells in S^n are called *concentric* if there is a homeomorphism of one onto R_2 that sends the other onto R_1 . If D_1 and D_2 are concentric, we also say that D_1 is concentric to D_2 and vice versa. Two n -cells D and D' with $D \subset D'$ are concentric if and only if $D' - \text{Int } D$ is homeomorphic to $S^{n-1} \times [0, 1]$. A finite number of disjoint n -cells $D_1, \dots, D_k$ in S^n are called *similar* if there is an n -cell to which each D_i is concentric. The following problem should be important.

THE CONCENTRICITY PROBLEM. Let D' be a tame n -cell in S^n containing another tame n -cell D in its interior. Are D and D' concentric?

This problem is equivalent to the following:

THE SIMILARITY PROBLEM. Let $D_1, \dots, D_k$ be disjoint tame n -cells in S^n . Are they similar?

These problems have the affirmative answers for $n \leq 3$. They are open for $n \geq 4$. We will require in certain theorems or lemmas that certain n -cells be similar or concentric. Such requirements may be deleted if and when either of the above problems is solved in the affirmative.

Let D be an n -cell in S^n . An n -cell D' in S^n is called an ϵ -pseudo-approximation of D if $\text{Int } D'$ is an ϵ -neighborhood of D .

LEMMA 4. *Let D and E be two tame n -cells in S^n such that $D \subset \text{Int } E$. Then given a positive number ϵ there exists an ϵ -pseudo-approximation D' of D such that D' is concentric to E .*

LEMMA 5. *Let $x_1, x_2, \dots, x_k$ be distinct points in the interior of an n -cell D . Let $y_1, y_2, \dots, y_k$ be distinct points in $\text{Int } D$. Then there exists a homeomorphism of D onto itself such that it leaves $\text{Bd } D$ point-wise fixed and sends each x_i to y_i .*

The proofs of the above two lemmas are omitted.

THEOREM 5. *Let $D_1, \dots, D_k$ be disjoint n -cells contained in, and concentric to, an n -cell D . Let $D'_1, \dots, D'_k$ be disjoint n -cells contained in, and concentric to, D . Then there exists a homeomorphism h of D onto itself that moves no point of $\text{Bd } D$ and takes D_i onto D'_i .*

PROOF. Let $x_i \in \text{Int } D_i$ and $y_i \in \text{Int } D'_i$. By Lemma 5, there exists a homeomorphism h_i of D_i onto itself that leaves each point of $\text{Bd } D_i$

fixed and takes x_i to y_i . Let h_2 be a homeomorphism of D onto itself that leaves $\text{Bd } D$ pointwise fixed and pulls $h_1(D_i)$ toward y_i such that $h_2h_1(D_i) \subset \text{Int } D'_i$. Now $h_2h_1(D_i)$ is concentric to D'_i for each i . Therefore there exists a homeomorphism h_3 that moves no point that is far from $\sum D'_i$ and takes $h_2h_1(D_i)$ onto D'_i . We finally let $h = h_3h_2h_1$.

COROLLARY. *Let $D_1, D_2, \dots, D_k$ be disjoint n -cells contained in, and concentric to, an n -cell D . Let $E_1, E_2, \dots, E_k$ be disjoint n -cells contained in, and concentric to, an n -cell E . Then for any given homeomorphism h of $\text{Bd } D$ onto $\text{Bd } E$, there exists an extension homeomorphism H of h that takes D onto E and each D_i onto E_i .*

THEOREM 6. *Let G be an upper semicontinuous decomposition of S^n , $n > 1$, having a countable number of nondegenerate elements whose sum F is closed. Then the decomposition space X of G is an n -sphere if and only if each element of G has an open n -cell complement.*

PROOF. Suppose X is an n -sphere. We denote the quotient map of G by f . For given $g \in G$, there exist n -cells $D_1, D_2, \dots$ in X such that each D_i is tame, $\text{Int } D_i \supset D_{i+1}$, $\text{Bd } D_i \cdot f(F) = \emptyset$ and the intersection of D_i is the point $f(g)$. Since each $(n-1)$ -sphere $f^{-1}(\text{Bd } D_i)$ is apart from F by a positive distance, it is tame. A repeated application of Lemma 4 shows that $S^n - g$ is an open n -cell.

The converse follows, since F is a G_δ set, from [1, Theorem 1]. (Actually, Bing proves his theorem for $n=3$ but the proof works for $n>3$ as well. Also, Theorem 6 can be slightly strengthened. In fact, we only need to assume that $F+A$ is closed for some countable set A .)

THEOREM 7. *Let G be a monotone upper semicontinuous decomposition of S^n and F be the sum of the nondegenerate elements of G . Suppose for each positive number ϵ , there exist a finite number of disjoint n -cells $D_1, D_2, \dots, D_k$ such that $\text{Cl } F \subset \sum \text{Int } D_i$ and each D_i lies in an ϵ -neighborhood of an element of G it contains. Then the decomposition space X is a topological n -sphere.*

PROOF. Let $D_1, \dots, D_k$ be disjoint tame n -cells such that $\text{Cl } F \subset \sum \text{Int } D_i$ and each D_i lies in a 1-neighborhood of an element of G it contains. Let ϵ_1 be a positive number smaller than $1/2$ such that no $2\epsilon_1$ -neighborhood of any $\text{Bd } D_i$ meets $\text{Cl } F$. Let $D'_1, \dots, D'_{k_1}$ be disjoint n -cells such that $\text{Cl } F \subset \sum \text{Int } D'_i$ and each D'_i lies in an ϵ_1 -neighborhood of an element of G it contains. Then $\sum D'_i \subset \sum \text{Int } D_i$. In general, there exist disjoint n -cells $D^{(p)}_1, \dots, D^{(p)}_{k_p}$ such that $\text{Cl } F \subset \sum \text{Int } D^{(p)}_i$, each $D^{(p)}_i$ lies in a $1/2^p$ -neighborhood of an element of G it contains and $\sum D^{(p)}_i \subset \sum \text{Int } D^{(p-1)}_i$.

If g is an element of G contained in $\text{Cl } F$, there exists a sequence $D_{i_0}, D_{i_1}, \dots, D_{i_p}^{(p)}, \dots$ such that each $D_{i_p}^{(p)}$ contains g . Let $g_p \subset D_{i_p}^{(p)}$ be an element of G of which $D_{i_p}^{(p)}$ is in a $1/2^p$ -neighborhood. Let $g' \in G$ be a limit set of g_p 's. Then an ϵ -neighborhood U of g' meets infinitely many g_p . Therefore, by the upper semicontinuity of G , U contains infinitely many g_p . Hence, infinitely many $D_{i_p}^{(p)}$ lie in $(\epsilon + 1/2^p)$ -neighborhoods of g' . This means that the intersection of $D_{i_p}^{(p)}$ is g' and $g' = g$. On the other hand, if $x \notin \text{Cl } F$, there exists p such that $x \notin \sum D_{i_p}^{(p)}$.

We now simplify the situation a little. By shrinking $D^{(p)}$'s slightly inward, we may assume they are tame. Then using pseudo-approximations of Lemma 4, we may assume that D_i are similar and $D_i(p)$ is concentric to $D_j^{(p-1)}$ if the latter contains the former. Using Corollary to Theorem 5 and the method of Harrold [4], we see that X is an n -sphere.

REFERENCES

1. R. H. Bing, *Upper semicontinuous decompositions of E^3* , Ann. of Math. **65** (1957), 363-374.
2. ———, *Approximating surfaces with polyhedral ones*, Ann. of Math. **65** (1957), 456-483.
3. M. Brown, *A proof of the generalized Schoenflies theorem*, Bull. Amer. Math. Soc. **66** (1960), 74-76.
4. O. G. Harrold, Jr., *A sufficient condition that a monotone image of the three-sphere be a topological three-sphere*, Proc. Amer. Math. Soc. **9** (1958), 846-850.
5. W. Hurewicz and H. Wallman, *Dimension theory*, Princeton, 1941.
6. E. E. Moise, *Affine structures in 3-manifolds. II. Positional properties of 2-spheres*, Ann. of Math. **55** (1952), 172-176.
7. R. L. Wilder, *Topology of manifolds*, Amer. Math. Soc. Colloq. Publ. Vol. 32, Amer. Math. Soc., New York, 1949.

THE SEOUL NATIONAL UNIVERSITY, SEOUL, KOREA