ON TOTALLY MINIMAL SETS

HSIN CHU

The following theorem: "Let (X, T, π) be a transformation group, where T is the additive group of all real numbers with the usual topology and X is a compact Hausdorff space containing more than one point and let X be an almost periodic minimal orbit-closure under T . Then X is not totally minimal under T ," which we may find in [3], is due to Professor E. E. Floyd. In this note, we show his theorem is still true if we weaken the condition that T is the real group to be any locally compact, non-totally-disconnected, abelian group and change the condition that X contains more than one point to be there is $a \in X$ such that aG_0 contains more than one point, where G_0 is the connected component of the identity of T . Thus we find a large class of transformation groups which is not totally minimal. The Pontrjagin duality of locally compact abelian groups is used in the proof. For the notations occurring here consult [3].

THEOREM. *Let (X, T, π) be a transformation group, where T is a locally compact, non-totally-disconnected, abelian group and X is a compact Hausdorff space and there is $a \in X$ such that aG_0 contains more than one point where G_0 is the connected component of identity in T , and let X be an almost periodic minimal orbit-closure under T . Then X is not totally minimal under T .*

PROOF. By a known result (see [3, p. 39]), X has a compact group structure such that a is the identity and $\pi_a: T \rightarrow X$ is a continuous homomorphism such that $\text{Cl}((T)\pi_a) = \text{Cl}(aT) = X$. It follows that $\text{Cl}(aG_0)$ is a closed connected subgroup of X , which is not trivial. It is known there exists a continuous homomorphism f_a from $\text{Cl}(aG_0)$ onto the circle group K and we can extend f_a to a continuous homomorphism $f: X \rightarrow K$ such that $f|_{\text{Cl}(aG_0)} = f_a$. Define $g: T \rightarrow K$ by $g = \pi_a f$. Consider $g|_{G_0}: G_0 \rightarrow K$. Since $(G_0)\pi_a = aG_0$ is a nontrivial connected subgroup of X , which is dense in $\text{Cl}(aG_0)$, $(G_0)g = (G_0\pi_a)f$ is a nontrivial connected subgroup K . However, the only nontrivial connected subgroup of the circle group K is K itself. It follows that $(G_0)g = K$. Hence $g: T \rightarrow K$ is an onto mapping. Let $g^{-1}(0) = L$ then L is a closed syndetic subgroup of T (see [3, 2.03]) and

$$0 \rightarrow L \xrightarrow{i} T \xrightarrow{g} K \rightarrow 0$$

is exact with i and g are relatively open mappings (with respect to their images), where i is the inclusion mapping. Thus, we have the

Received by the editors April 3, 1961.

following commutative diagram.

$$\begin{array}{ccccccc}
 & & & X & & & \\
 & & r \nearrow & \uparrow \pi_a & \nwarrow f & & \\
 0 & \longrightarrow & L & \xrightarrow{i} & T & \xrightarrow{g} & K \longrightarrow 0
 \end{array}$$

where $r = i\pi_a$. Then the following diagram

$$\begin{array}{ccccccc}
 & & & X^* & & & \\
 & & r^* \nearrow & \uparrow \pi_a^* & \nwarrow f^* & & \\
 0 & \longleftarrow & L^* & \xleftarrow{i^*} & T^* & \xleftarrow{g^*} & K^* \longleftarrow 0
 \end{array}$$

is also commutative, where L^*, T^*, K^*, X^* are the character groups, with the usual compact-open topology, of L, T, K, X respectively. We know from [4], that $0 \rightarrow L^* \xrightarrow{i^*} T^* \xrightarrow{g^*} K^* \leftarrow 0$ is exact, where i^* and g^* are relatively open mappings (with respect to their images) and $K^* \cong Z$, where Z is the integer group with the discrete topology and f^* is isomorphic into such that f^* is relatively open with respect to its image and X^* is a nontrivial discrete group. From [1], we know $\pi_a^*: X^* \rightarrow T^*$ is continuously isomorphic into. Since g^*i^* is trivial, r^* cannot be isomorphic into. By a result in [1], $r: L \rightarrow X$ is not dense, i.e. $\text{Cl}((L)r) = \text{Cl}(aL) \neq X$. This shows (X, L, π) is not a minimal set. This theorem is proved.

Using a similar argument, we have the following result:

COROLLARY. *Let (X, T, π) be a transformation group, where T is a locally compact abelian group and X is a compact, totally disconnected, Hausdorff space containing more than one point. If X is an almost periodic minimal orbit-closure under T . Then X is not totally minimal under T .*

REFERENCES

1. H. Anzai and S. Kakutani, *Bohr compactifications of a locally compact abelian group* I, Proc. Imperial Acad. Tokyo 19 (1943), 476-480.
2. ———, *Bohr compactifications of a locally compact abelian group*. II, Proc. Imperial Acad. Tokyo 19 (1943), 533-539.
3. W. Gottschalk and G. Hedlund, *Topological dynamics*, Amer. Math. Soc. Colloq. Publ. Vol. 36, Amer. Math. Soc., Providence, R. I., 1955.
4. A. Weil, *L'intégration dans les groupes topologiques et ses applications*, Actualités Sci. Ind. Nos. 869-1145, Paris, Hermann, 1953.

NORTHWESTERN UNIVERSITY