$$(5) \quad f(x, y) \leq (2\pi)^{-2} D \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) dx dy.$$

If $[A]$ is the matrix $2\pi[a]^{-1}$ the analogue of (3) is

$$D\phi(0, 0) = \sum_{m=-\infty}^{\infty} \sum_{n=-\infty}^{\infty} f(A_{11}m + A_{12}n + x, A_{21}m + A_{22}n + y),$$

and (5) follows in the same way as (1).

REFERENCES

1. S. Bernstein, *Sur la borne supérieure du module de la dérivée d'une fonction de degré fini*, Dokl. Akad. Nauk SSSR 51 (1946), 567-568; Collected Works, Vol. 2, Izdatelstvo Akademii Nauk SSSR, Moscow, 1954, pp. 396-398.
2. R. P. Boas, Jr., *Entire functions*, Academic Press, New York, 1954.
3. S. Bochner, *Vorlesungen über Fouriersche Integrale*, Akademische Verlagsgesellschaft, Leipzig, 1932.
4. L. H. Loomis, *An introduction to abstract harmonic analysis*, Van Nostrand, New York, 1953.
5. M. Plancherel and G. Pólya, *Fonctions entières et intégrales de Fourier multiples* Comment. Math. Helv. 9 (1937), 224-248; 10 (1938), 110-163.

NORTHWESTERN UNIVERSITY

REPRESENTATIONS OF BANACH SPACES¹

G. S. YOUNG

Banach and Mazur² proved that every separable Banach space B can be represented as the space $C(M)$ of continuous real functions on a compact metric space M . Since M is the continuous image of the Cantor set K , $C(M)$ can be imbedded in $C(K)$, and since functions in $C(K)$ can be extended preserving norm to functions over I^1 , they conclude that B can be represented as a subspace of $C(I^1)$.

If B is not separable, it can be represented as $C(H)$, where H is compact Hausdorff. A compact Hausdorff space is the continuous image of some totally disconnected compact Hausdorff space T —for example, give the space the discrete topology, and let T be its Stone-Čech compactification. It follows that B is isomorphic to a subspace of $C(T)$. If T could be given a linear order inducing the same topology, we could fill in the missing intervals and obtain a compact con-

Received by the editors May 20, 1961.

¹ The work on this note was supported by the National Science Foundation.

² S. Banach, *Théorie des opérations linéaires*, Warsaw, 1932.

nected ordered space S , and so represent B as a subspace of $C(S)$. I cannot have been the only one who observed this, and failed in ordering T .

Mardešić and Papić³ have proved the unexpected and elegant result that *if ΠX_α is a product of two or more nondegenerate continua, and there is a continuous map of a compact ordered continuum onto ΠX_α , then each X_α is metric*. From this it follows, e.g., that *if L is the "long interval," obtained by inserting open intervals between consecutive points of the set of ordinals not greater than the first uncountable ordinal, then $L \times I^1$ is not the continuous image of any compact ordered space, T .*

For suppose that there is a map $f: T \rightarrow L \times I^1$, onto. Let a_1, a_2 be two points of T with no point between them. Let $f(a_j) = (l_j, t_j)$, $j = 1, 2$, and suppose $l_1 \leq l_2$ in L . The set L contains an "interval" $l_1 l_2$, possibly a point, and in I^1 there is an interval $t_1 t_2$, also possibly a point. The set $X = l_1 l_2 \times t_1 \cup l_2 \times t_1 t_2$ is a continuum ordered by separation. Between a_1 and a_2 insert a copy X_{12} of $X - f(a_1) - f(a_2)$ so that the natural map of $a_1 \cup X_{12} \cup a_2$ onto X is a homeomorphism. Having done this for each pair a_1, a_2 , we obtain a compact connected ordered space T^* . We define $f^*: T^* \rightarrow L \times I^1$ to agree with f on T and to be the natural map on each set X_{12} . It is easily seen that f^* is continuous. But the Mardešić-Papić result then implies that L is metric, a contradiction. There are obvious generalizations. A corollary is that there exist compact totally disconnected Hausdorff spaces that cannot be ordered.

The algebra of $C(L \times I^1)$ must be quite different from that of $C(I^1 \times I^1)$. It would be interesting to see how this property manifests itself algebraically.

THE TULANE UNIVERSITY OF LOUISIANA

³ S. Mardešić and P. Papić, Glasnik Mat. Fiz. Astr. 15 (1960). I am indebted to Professor Mardešić for informing me of their results.