Since $\int_0^\infty |F| < \infty$, the first equation in (4) has a solution, say θ_1 , such that $\theta_1 \rightarrow 0$ as $x \rightarrow \infty$. Put $r_1 = \exp - \int_0^x F \cos^2(f + \theta_1)$. Then r_1 and θ_1 constitute a solution of (4). Set $y_1 = r_1 \sin(f + \theta_1)$. One can construct y_2 similarly.

REFERENCES

1. N. Levinson, *The asymptotic nature of solutions of linear systems of differential equations*, Duke Math. J. 15 (1948), 111-126.
2. A. Wintner, *Asymptotic integrations of the adiabatic oscillator*, Amer. J. Math. 69 (1947), 251-272.

REMINGTON RAND UNIVAC, WASHINGTON, D. C. AND
HOWARD UNIVERSITY

REMARK ON $\pi(x) = o(x)$

S. E. MAMANGAKIS¹

In Hardy and Wright [1], Landau [2], and Prachar [3] there are proofs of $\pi(x) = o(x)$. The purpose of this paper is to provide a simpler proof.

THEOREM. $\pi(x) = o(x)$.

PROOF. Let M_r denote the product of the first r primes. Given any positive real number $x \geq 2$, it is clear that there exist unique positive integers k and r , with $1 \leq k \leq p_{r+1} - 1$, such that

$$(1) \quad kM_r \leq x < (k+1)M_r.$$

For any x satisfying (1) we must also have

$$(2) \quad \pi(x) \leq (k+1)\phi(M_r) + r$$

(where $\phi(M_r)$ is the totient of M_r). From (1) and (2) we obtain

$$(3) \quad 0 \leq \frac{\pi(x)}{x} \leq \frac{(k+1)\phi(M_r)}{kM_r} + \frac{r}{kM_r}.$$

Since $kM_r \geq kr! \geq r!$ and $(k+1)/k \leq 2$, we replace (3) by

$$(4) \quad 0 \leq \frac{\pi(x)}{x} \leq 2 \prod_{i=1}^r (1 - p_i^{-1}) + \frac{1}{(r-1)!}.$$

Received by the editors August 14, 1961 and, in revised form, March 19, 1962.

¹ This work was supported in part by contract Nonr285(46) with the Office of Naval Research.

Since there are infinitely many primes (a fact already used implicitly in defining k and r), $r \rightarrow \infty$ as $x \rightarrow \infty$. We may therefore write

$$0 \leq \limsup_{x \rightarrow \infty} \frac{\pi(x)}{x} \leq 2 \prod_{i=1}^{\infty} (1 - p_i^{-1}) + \lim_{r \rightarrow \infty} \frac{1}{(r-1)!} = 0,$$

where the divergence of $\prod_{i=1}^{\infty} (1 - p_i^{-1})$ to zero follows from the divergence of $\sum_{i=1}^{\infty} p_i^{-1}$.

REFERENCES

1. G. H. Hardy and E. M. Wright, *An introduction to the theory of numbers*, 4th ed., Clarendon Press, Oxford, 1960, pp. 349-350.
2. E. Landau, *Handbuch der Lehre von der Verteilung der Primzahlen*, Erster Band, B. G. Teubner, Leipzig, 1909, pp. 67-71.
3. K. Prachar, *Primzahlverteilung*, Springer-Verlag, Berlin, 1957, pp. 13-14.

COURANT INSTITUTE OF MATHEMATICAL SCIENCES,
NEW YORK UNIVERSITY