A CHARACTERIZATION OF CERTAIN QUASI-OPEN MAPPINGS

ROBERT H. KASRIEL¹

1. Introduction. In this paper we shall deal with mappings (continuous functions) defined on regions (open and connected) subsets of a plane P with the range of the functions also contained in P . In reference [1], M. K. Fort defined such a mapping f to be *minimal* provided that for each closed 2-cell N contained in the domain of f , $f(N) \subset g(N)$ for every mapping g whose domain contains N and which is such that $f|_{\text{Fr } N} = g|_{\text{Fr } N}$. The following result was obtained in [1].

If f is light, then f is open if and only if f is minimal.

In this paper, we prove the following similar theorem for certain quasi-open mappings.

1.1. THEOREM. *Let $f: X \rightarrow P$ be a compact mapping defined on a simply connected region X in a plane P with $f(X) \subset P$. Then f is quasi-open if and only if f is minimal and for no $x \in f(X)$ is it true that $f^{-1}(x)$ separates X .*

The sufficiency of the condition in 1.1 is established in §2 and the necessity in §3.

Recall that a mapping $f: X \rightarrow P$ is compact provided that for each compact set $K \subset f(X)$, $f^{-1}(K)$ is compact. f is quasi-open provided that for any $y \in f(X)$ and any open set U in X containing a compact component of $f^{-1}(y)$, $y \in \text{int } f(U)$. Here and throughout the paper such terms as *interior*, *closed*, *closure* (cl), *boundary* (Fr) will always be relative to the containing plane.

With $f: X \rightarrow P$ a mapping into a plane P , if R is an *elementary region* (a bounded plane region whose boundary consists of a finite number of simple closed curves) whose closure is in the domain of f and $x \in P - f(\text{Fr } R)$, then there is defined the topological index or winding number $\mu(x, f, R)$. Use will be made of the following well-known properties of the index.

1.2. $\mu(x, f, R)$ is constant on every component of $P - f(\text{Fr } R)$.

1.3. Suppose $x \in P - f(\text{Fr } R)$. If $g: X \rightarrow P$ is such that $g(\bar{x}) = f(\bar{x})$ for $\bar{x} \in \text{Fr}(R)$, then $\mu(x, f, R) = \mu(x, g, R)$.

1.4. Let R_j ; $j = 1, 2, \dots, k$ be a sequence of pair-wise disjoint ele-

Received by the editors August 11, 1961.

¹ This research supported in part by the National Science Foundation Grant G-7343 at the Georgia Institute of Technology.

mentary regions in R such that

$$(\text{cl } R) \cdot f^{-1}(x) \subset \bigcup_j R_j.$$

Then $\mu(x, f, R) = \sum_j \mu(x, f, R_j)$. (See Theorem 3 on p. 126 in [7].)

1.5. If $\mu(x, f, R) \neq 0$, then $x \in f(R)$.

1.6. If N is a closed 2-cell and f is a homeomorphism on N , then $\mu(x, f, \text{int } N) = \pm 1$ for each $x \in f(\text{int } N)$.

In terms of this index, Fort [1] proved the following.

1.7. With $f: X \rightarrow P$ as before, f is minimal if and only if for each closed 2-cell N , $f(N) = f(\text{Fr } N) + \{p \mid \mu(p, f, \text{int } N) \neq 0\}$.

2. Compact plane mappings. In this section we make the following assumption.

2.1. $f: X \rightarrow P$ is a compact mapping from a simply connected region in a plane P into P .

From a result of Whyburn (§10 in [4]), there exists a factorization $f = lm$ in which m is monotone and compact and l is light and compact. Throughout the middle space $m(X)$ will be designated as M .

If the further assumption is made that for no $x \in f(X)$ it is true that $f^{-1}(x)$ separates X , it can be shown that no component of $f^{-1}(x)$ separates X . But then for no $z \in M$, is it true that $m^{-1}(z)$ separates X . 2.3 below then follows from the following result of Whyburn. (See Note 1, p. 313 in [5].)

If m is a monotone mapping from a plane X onto a space Y , then Y is a plane if and only if m is compact and $m^{-1}(y)$ does not separate X for each $y \in Y$.

2.3. Let f be as in 2.1 and let $f = lm$ be a factorization as before. In addition assume that for $x \in f(X)$, $f^{-1}(x)$ does not separate X . Then the middle space M is a topological plane.

2.4. Let $f: X \rightarrow P$, l, m, M be as in 2.3. Then if N is a closed 2-cell such that $N \subset M$, $m^{-1}(\text{int } N)$ is a simply connected region.

PROOF. $M - \text{int } N$ is connected. Therefore, since m is compact and monotone, $m^{-1}(M - \text{int } N) = X - m^{-1}(\text{int } N)$ is connected. (See 8.3 in [4].) Since m is compact, $m^{-1}(N)$ and thus $\text{cl } (m^{-1}(\text{int } N))$ is compact. Thus $m^{-1}(\text{int } N)$ is a plane region whose closure is compact and whose complement in the plane is connected. Hence $m^{-1}(\text{int } N)$ is simply connected.

2.5. Let $f: X \rightarrow P$ be a compact mapping. Let $x \in f(X)$ and let U be an open set such that $f^{-1}(x) \subset U$. Then there exists an open set $V \subset P$ such that $x \in V$ and $f^{-1}(V) \subset U$.

This follows easily from the compactness of f .

2.6. Let $f: X \rightarrow P$ be a mapping as in 2.3. Furthermore assume that f

is minimal. Then f is quasi-open.

PROOF. Let $f=lm$ be a factorization as in 2.3. By 2.3 the middle space M may be taken to be a complex plane. Let $x \in f(X)$ and suppose K is a component of $f^{-1}(x)$. Let U be an open set in X such that $K \subset U$. From the definition of quasi-open, the proof will be complete if we show that x is an interior point of $f(U)$. Toward this end, let $p = m(K)$. Note that $p \in l^{-1}(x)$. Since $l^{-1}(x)$ is totally disconnected it can be shown that there exists arbitrarily small 2-cells N such that $(\text{Fr } N) \cdot l^{-1}(x) = \emptyset$ and $p \in \text{int } N$. By 2.5, N may be chosen so that $m^{-1}(N) \subset U$. Let $G \equiv m^{-1}(\text{int } N)$. By 2.4, G is simply connected. Since f is compact, $f^{-1}(x) \cdot \text{cl } G$ is compact and since $f^{-1}(x) \cdot \text{Fr } G = \emptyset$ it follows that $f^{-1}(x) \cdot G$ is compact. Then since G is simply connected, there exists a closed 2-cell T such that $f^{-1}(x) \cdot G \subset \text{int } T \subset T \subset G$. Further by 1.7 $\mu(x, f, \text{int } T) \neq 0$. But then from 1.2 and 1.5, $f(T) \supset Q_x$ where Q_x is the component of $P - f(\text{Fr } T)$ that contains x . Since Q_x is open and $x \in Q_x \subset f(T) \subset f(U)$ it follows that x is an interior point of $f(U)$.

3. Compact quasi-open mappings. In this section we make the following assumption.

3.1. $f: X \rightarrow P$ is a compact and quasi-open mapping defined on a simply connected region X in the plane P and $f(X) \subset P$.

Assuming 3.1, it is known (10.2 and 10.4 in [4]) that there is a monotone-light open factorization $f=lm$ for which m is a compact monotone mapping and l is compact, light, and open relative to $f(X)$. Further from the definition of quasi-open as given in the introduction, $f(X)$ is open in P and hence l is open relative to P . Furthermore, we have the following.

3.2. For no point $x \in f(X)$ is it true that $f^{-1}(x)$ separates X .

PROOF. Suppose $f^{-1}(x)$ separates X . Then there exists a separation

$$X - f^{-1}(x) = A + B.$$

Since $f^{-1}(x)$ is compact $A + f^{-1}(x)$ or $B + f^{-1}(x)$ is compact. Assume that $A + f^{-1}(x)$ is compact. Then $f(A) + x$ is compact. However A is open and is the union of components of point inverses. Hence from the quasi-openness of f , $f(A)$ is open. This is a contradiction since then x would be the boundary of an open set $f(A)$ in the plane.

On the basis of 3.2 and 2.3 with f as in 3.1, the middle space M can be taken to be a plane. Recall that $f(X)$ is open in P . Thus there is available the following result of Whyburn.

3.3. (See VIII, 1.1 and 1.11 in [3].) There exists an integer k and a completely scattered set $D \subset f(X)$, such that for each $x \in f(X) - D$, $l^{-1}(x)$ consists of k distinct points. Furthermore on the set $M - l^{-1}(D)$, l is a

local homeomorphism. [Note that for each point $x \in D$, x is an isolated point of D . Also D is closed.]

By making use of 3.3 and 2.4, the following can be obtained.

3.4. For each $z \in f(X) - D$, there exists arbitrarily small closed 2-cells N such that $z \in \text{int } N \subset N \subset f(X) - D$ and such that $f^{-1}(\text{int } N)$ is the union of a finite number of pairwise disjoint open two-cells Q_i for which $f(Q_i) = \text{int } N$ and $f^{-1}(z^*) \cdot Q_i$ consists of a single component of $f^{-1}(z^*)$ for each $z^* \in \text{int } N$.

3.5. $X - f^{-1}(D)$ is a connected open set.

PROOF. Notice that $X - f^{-1}(D) = m^{-1}(M - l^{-1}(D))$ is open and furthermore is connected since $M - l^{-1}(D)$ is connected and m is a compact monotone mapping.

3.6. DEFINITION OF $\mu(x)$. On the open connected set $S \equiv X - f^{-1}(D)$ define the function μ as follows. Let $x \in S$. Then $f^{-1}f(x)$ consists of exactly k compact components. (See 3.3.) Let K_x be that component that contains x . There exists an elementary region R such that

$$(1) \quad K_x \subset R \subset \text{cl } R \subset X - f^{-1}(D) \quad \text{and} \quad (f^{-1}f(x)) \cdot \text{cl } R = K_x.$$

It will be shown in 3.7 that $\mu(x, f, R)$ is independent of the particular choice of R provided that R satisfies (1). On that basis the following may be defined:

$$\mu(x) \equiv \mu(f(x), f, R).$$

(Because of the nature of point inverses of f , μ is a point function version of the function defined in II. 3.4 of [7].)

3.7. $\mu(x, f, R_1) = \mu(x, f, R_2)$ provided that R_1 and R_2 satisfy (1).

PROOF. Let Q be the component of $R_1 \cdot R_2$ that contains K_x . There exists an elementary region R such that

$$K_x \subset R \subset \text{cl } R \subset Q \subset R_1 \cdot R_2.$$

From 1.4, $\mu(f(x), f, R_1) = \mu(f(x), f, R) = \mu(f(x), f, R_2)$.

Use will be made of the following in investigating the function μ .

3.8. YOUNG'S MODIFICATION THEOREM. (See [6].) Let N be a closed 2-cell with interior G and boundary J . Suppose X is a continuum and $m: X \rightarrow N$ is a monotone mapping of X onto N such that $m^{-1}(G)$ is an open 2-cell. Then there is a mapping $h: X \rightarrow N$ of X onto N such that $h(x) = m(x)$ for $x \in m^{-1}(J)$ and $h^{-1}h(x) = x$ for $x \in m^{-1}(G)$.

3.9. $|\mu(x)| = 1$ for each $x \in X - f^{-1}(D)$.

PROOF. Let $x \in X - f^{-1}(D)$. Let R be an elementary region admissible for the computation of $\mu(x)$. (See 3.6.) Let $y = m(x) \in M$. Note that l is a local homeomorphism at y and recall that m is compact.

Hence by 2.5 there exists a closed 2-cell neighborhood N of y such that $m^{-1}(N) \subset R$ and such that l is a homeomorphism on N . Let $G = m^{-1}(\text{int } N)$. By 2.4 G is an open 2-cell. Thus 3.8 can be applied to $m|_{m^{-1}(N)}$ and using this a modification $h: X \rightarrow M$ of m can be defined such that $h(z) = m(z)$ for $z \in X - G$ and h is one-to-one on G . Let z^* be the unique point in G such that $h(z^*) = y$. Note also $h^{-1}(y) \cdot R = z^*$. Let W be a closed 2-cell such that $z^* \in \text{int } W \subset W \subset G$. Note that $lh|_W$ is a homeomorphism. Then by 1.6 and from the fact that $lh(z^*) = l(y) = f(x)$, it follows that $\mu(f(x), lh, \text{int } W) = \pm 1$. Further since $[(lh)^{-1}f(x)] \cdot R = z^*$, it follows from 1.4 that $\mu(f(x), lh, \text{int } W) = \mu(f(x), lh, R)$. Moreover by 1.3, since $f = lh$ on $\text{Fr}(R)$, $\mu(f(x), f, R) = \mu(f(x), lh, R)$. Combining this with the two previous equalities involving μ yields $\mu(f(x), f, R) = \pm 1$.

3.10. μ is continuous on $X - f^{-1}(D)$. (Hence by 3.5 $\mu(x)$ is constant on $X - f^{-1}(D)$.)

PROOF. Let $x \in X - f^{-1}(D)$ and K_x the component of $f^{-1}(x)$ that contains x . Choose R as in (1) of 3.6 so that $\mu(x) = \mu(f(x), f, R)$. Choose a closed 2-cell N such that $f(x) \in \text{int } N \subset \text{cl } N \subset f(X) - D - f(\text{Fr}(R))$ and is chosen as in 3.4. Let Q be the component of $f^{-1}(\text{int } N)$ that contains K_x . Since $f(Q_i) = \text{int } N$ for each component Q_i of $f^{-1}(\text{int } N)$ (see 3.4) and since $f^{-1}(x) \cdot R = K_x$, it follows that Q is the only component of $f^{-1}(\text{int } N)$ that intersects R .

Now let $y \in Q$. Let K_y be the component of $f^{-1}(f(y))$ that contains y . However from the way in which N was chosen (see 3.4) K_y is the only component of $f^{-1}f(y)$ that intersects R . Thus R is admissible for the computation of $\mu(y)$ and $\mu(y) = \mu(f(y), f, R)$. However by 1.2, $\mu(f(y), f, R) = \mu(f(x), f, R)$ and thus $\mu(x) = \mu(y)$.

3.11. Let $f: X \rightarrow P$ be as in 3.1. Then f is minimal.

PROOF. Let N be a closed 2-cell in X . From 1.5 and 1.7 we need show only that $\mu(x, f, \text{int } N) \neq 0$ for each $x \in f(N) - f(\text{Fr } N)$. Let $x \in f(N) - f(\text{Fr } N)$. Define D as in 3.3. Let U_x be the component of $P - f(\text{Fr } N)$ that contains x . There exists a point $y \in U_x \cap [f(\text{int } N) - D]$. This follows from the quasi-openness of f and the definition of D . By 1.2, $\mu(y, f, \text{int } N) = \mu(x, f, \text{int } N)$. Let $K_1, K_2, \dots, K_h$ be the components of $f^{-1}(y)$ that intersect and hence are contained in $\text{int } N$. There exist elementary regions $R_1, R_2, \dots, R_h$ such $K_i \subset R_i \subset \text{cl } R_i \subset \text{int } N$ and $R_i \cdot R_j = \emptyset$ if $i \neq j$. By 1.4, $\mu(y, f, \text{int } N) = \sum_i \mu(y, f, R_i)$. Let $x_i \in K_i$. Thus, from definition of μ , 3.9, and 3.10 $|\sum_i \mu(y, f, R_i)| = |\sum_i \mu(x_i)| = h$ and $\mu(x, f, \text{int } N) = \mu(y, f, \text{int } N) \neq 0$.

THEOREM 1.1. *The validity of the Theorem 1.1 stated in the introduction now follows from 2.6, 3.2 and 3.11.*

REFERENCES

1. M. K. Fort, *A characterization of plane light open mappings*, Proc. Amer. Math. Soc. **2** (1951), 175-177.
2. G. T. Whyburn, *Analytic topology*, Amer. Math. Soc. Colloq. Publ. Vol. 28, Amer. Math. Soc., New York, 1942.
3. ———, *Topological analysis*, Princeton Univ. Press, Princeton, N. J., 1958.
4. ———, *Open mappings on locally compact spaces*, Mem. Amer. Math. Soc. No. 1 (1950).
5. ———, *Compactness of certain mappings*, Amer. J. Math. **81** (1959), 306-314.
6. J. W. T. Youngs, *Homeomorphic approximations to monotone mappings*, Duke Math. J. **15** (1948), 87-94.
7. T. Rado and P. V. Reichelderfer, *Continuous transformations in analysis*, Springer-Verlag, Berlin, 1955.

THE GEORGIA INSTITUTE OF TECHNOLOGY