FUNCTION ALGEBRAS WITH CLOSED RESTRICTIONS¹

I. GLICKSBERG

Let $C(X)$ be the (supremum normed) algebra of all complex continuous functions on a compact Hausdorff space X , and A a closed subalgebra containing the constants and separating the points of X . For $F \subset X$ let $f|F$ be the restriction of the function f to F , $A|F = \{f|F : f \in A\}$. The purpose of this note is to prove the following result, which characterizes $C(X)$ among its closed separating subalgebras.

THEOREM. *Suppose that for every closed $F \subset X$, $A|F$ is closed in $C(F)$. Then $A = C(X)$.*

Our proof leans heavily on some facts obtained recently by Katznelson [1]. In fact little else, beyond the Riesz representation theorem, is needed (for some facts concerning Banach algebras the reader is referred to [2; 3]).

Let F be a closed subset of X . $kF = \{f : f \in A, f(F) = 0\}$, the kernel of F , is an ideal in A , and we can form the quotient algebra A/kF ; for an element $f+kF$ ($f \in A$) of A/kF , $\|f+kF\|$ will denote the usual quotient norm ($\inf \{\|f+g\| : g \in kF\}$) under which A/kF is a Banach algebra. For $x \in F$, $f+kF \rightarrow f(x)$ is a multiplicative linear functional on A/kF , so necessarily of norm ≤ 1 , whence $\|f|F\| = \sup |f(F)| \leq \|f+kF\|$, where $\|f|F\|$ is of course the norm inherited by the closed subalgebra $A|F$ of $C(F)$. Now clearly $f+kF \rightarrow f|F$ is an isomorphism of A/kF onto $A|F$, and so, by the open mapping theorem, topological: there is a constant c_F for which

$$(1) \quad \|f+kF\| \leq c_F \|f|F\|, \quad f \in A,$$

a relation which we shall need somewhat later.

To begin the proof proper we shall show that for closed disjoint subsets F and K of X there is an f in A which is 1 on F , 0 on K . Indeed let x and y be distinct elements of X . Since A separates X , there is an f in A with $f(x) = 0, f(y) = 1$, so we can find closed neighborhoods V_x of x and W_x of y for which $|f(V_x)| \leq \frac{1}{4}, |1-f(W_x)| \leq \frac{1}{4}$. As is well known there is a sequence $\{p_n\}$ of polynomials in one complex variable which converge uniformly on $\{z : |z| \leq \frac{1}{4}\} \cup \{z : |1-z| \leq \frac{1}{4}\}$ to the characteristic function of $\{z : |z| \leq \frac{1}{4}\}$; consequently

Received by the editors January 5, 1962.

¹ Work supported in part by the National Science Foundation through Grant G14779, and in part by the Air Force Office of Scientific Research.

$\{(p_n \circ f) \mid (V_x \cup W_x)\}$ is a sequence of elements of $A \mid (V_x \cup W_x)$ which converges uniformly on $V_x \cup W_x$, tending to 1 on V_x , 0 on W_x . Since $A \mid (V_x \cup W_x)$ is closed under uniform convergence by hypothesis, there is an e in A which is 1 on V_x , 0 on W_x .

Now let y be a fixed element of K . F can be covered by finitely many V_x , $x \in F$, say $V_{x_1}, \dots, V_{x_n}$. If $e_1, \dots, e_n$ are the corresponding elements of A , then (as in a familiar argument) clearly $e'_1 = e_1 + e_2 - e_1 e_2$ is 1 on $V_{x_1} \cup V_{x_2}$ and 0 on the neighborhood $W_{x_1} \cap W_{x_2}$ of y ; similarly $e'_2 = e'_1 + e_3 - e'_1 e_3$ is 1 on $V_{x_1} \cup V_{x_2} \cup V_{x_3}$ and 0 on $W_{x_1} \cap W_{x_2} \cap W_{x_3}$. Continuing we thus obtain an element f_y of A which is 1 on F , and vanishes on some neighborhood U_y of y .

But now finitely many U_y cover K , so that the product f of the corresponding f_y yields an element of A which is 1 on F and 0 on K , as desired.

Alternatively stated, we have found an idempotent in $A/k(K \cup F)$ which separates F and K . What we now require is a bound on the norms of such idempotents, and to obtain this we repeat an argument of Katznelson's [1, Lemmas 3, 4, 5]. Let us say A is *bounded on a subset V of X* if there is a constant C_V for which, whenever F is closed and $F \subset V$, any idempotent in A/kF has norm $< C_V$. (Our next lemmas are taken from Katznelson [1], and are repeated here for the convenience of the reader.)

LEMMA 1. *Let V_1 and V_2 be open in X , and A be bounded on each V_i . Then A is bounded on every closed subset F of $V_1 \cup V_2$.*

PROOF. Since $F \setminus V_2$ is a closed subset of V_1 we can find an open $W_1 \supset F \setminus V_2$, $W_1^- \subset V_1$; and since $F \setminus W_1 \subset V_2$ there is an open $W_2 \supset F \setminus W_1$, $W_2^- \subset V_2$. $F \setminus W_1$ and $F \setminus W_2$ are disjoint closed subsets of X so there is a ϕ in A for which $\phi(F \setminus W_1) = 0$, $\phi(F \setminus W_2) = 1$, as we have seen.

Let P be a closed subset of F and let $f + kP$ be an idempotent in A/kP , so f is zero or 1 at each point of P . Then $f + k(P \cap W_i^-)$ is an idempotent in $A/k(P \cap W_i^-)$ and so (by the definition of the quotient norm) we can choose $f_i \in f + k(P \cap W_i^-)$ with $\|f_i\| < C_{V_i}$, $P \cap W_i^-$ being a closed subset of V_i , $i = 1, 2$. But now $f = \phi f_1 + (1 - \phi) f_2$ on P , since $P \subset F \subset W_1 \cup W_2$, and thus

$$\|f + kP\| \leq \|\phi f_1 + (1 - \phi) f_2\| < \|\phi\| C_{V_1} + \|1 - \phi\| C_{V_2}$$

so the right side may be taken as C_F .

Still following Katznelson let us say A is *bounded at $x \in X$* if A is bounded on V , for some neighborhood V of x . Then as an obvious consequence of Lemma 1 and compactness we have

LEMMA 2. *If F is closed and A is bounded at each x in F there is an*

open $V \supset F$ on which A is bounded.

LEMMA 3. *There are at most finitely many x in X at which A is not bounded.*

PROOF. If there were infinitely many we could find a sequence $\{x_n\}$ of them with disjoint open neighborhoods $\{V_n\}$ for which A is not bounded on V_n . So each V_n would contain a closed F_n for which there is an idempotent $f_n + kF_n$ in A/kF_n of norm $\geq n$.

Now let $F = (\bigcup F_n)^-$; since $V_n \cap F_m = \emptyset$ for $n \neq m$, $F = F_n \cup (\bigcup_{m \neq n} F_m)^-$, $F_n \cap (\bigcup_{m \neq n} F_m)^- = \emptyset$ and thus there is a ϕ in A with $\phi(F_n) = 1$, $\phi(F \setminus F_n) = 0$. Consequently ϕf_n is zero or 1 at each x in F , and $\phi f_n + kF$ is an idempotent in A/kF ; but

$$n \leq \|f_n + kF_n\| = \|\phi f_n + kF_n\| \leq \|\phi f_n + kF\|$$

(since $kF \subset kF_n$) so the idempotents in A/kF are not bounded in norm. By (1) they must be bounded, and the lemma is proved.

Let $\{x_1, \dots, x_n\}$ be the finite set given by Lemma 3 and let F be any closed set in X disjoint from $\{x_1, \dots, x_n\}$. By Lemma 2, A is bounded on a neighborhood of F ; so there is a constant C_F for which, whenever K is closed, $K \subset F$, each idempotent in A/kK has norm $< C_F$.

Now suppose μ is a (complex regular Borel) measure on F orthogonal to the closed subalgebra $A|_F$ of $C(F)$, and let K be a closed subset of F , $\epsilon > 0$. Choose a closed $K_0 \subset F \setminus K$ for which the total variation of μ on $(F \setminus K) \setminus K_0$ is less than ϵ/C_F , and then choose $f \in A$ which is 1 on K , 0 on K_0 , and of norm $< C_F$. Then

$$0 = \int f d\mu = \int_K 1 d\mu + \int_{K_0} 0 d\mu + \int_{(F \setminus K) \setminus K_0} f d\mu,$$

so $|\mu(K)| = |\int_{(F \setminus K) \setminus K_0} f d\mu| \leq C_F \cdot \epsilon/C_F = \epsilon$. Thus since $\epsilon > 0$ is arbitrary, $\mu K = 0$ for every closed $K \subset F$, and μ itself vanishes by regularity. Consequently $A|_F = C(F)$.

Now this implies any measure μ orthogonal to A is carried by $\{x_1, \dots, x_n\}$. For if not $\mu F \neq 0$ for some closed F disjoint from $\{x_1, \dots, x_n\}$ by regularity, and if V is an open neighborhood of $\{x_1, \dots, x_n\}$ with $V^- \cap F = \emptyset$, we can choose $f \in A$ with $f(V^-) = 0$, $f(F) = 1$; then $f\mu$ (the measure corresponding to the functional $g \rightarrow \int g f d\mu$ on $C(X)$) is a measure carried by $X \setminus V$, and orthogonal to A and so orthogonal to $A|_{(X \setminus V)} = C(X \setminus V)$, hence the zero measure. But $f\mu(F) = \int f f d\mu = \mu F \neq 0$, the desired contradiction, and μ orthogonal to A must be carried by $\{x_1, \dots, x_n\}$.

Since A separates the points of X this is clearly impossible unless all orthogonal measures vanish, i.e., unless $A = C(X)$, which completes the proof of the theorem.

REFERENCES

1. Y. Katznelson, *A characterization of the algebra of all continuous functions on a compact Hausdorff space*, Bull. Amer. Math. Soc. **66** (1960), 313–315.
2. L. H. Loomis, *Abstract harmonic analysis*, Van Nostrand, New York, 1953.
3. C. Rickart, *General theory of Banach algebras*, Van Nostrand, New York, 1960.

INSTITUTE FOR ADVANCED STUDY