ON THE LOGARITHMIC MEANS OF THE SUCCESSIVELY DERIVED CONJUGATE SERIES OF A FOURIER SERIES

SULAXANA KUMARI

1. **Definition [6].** A series $\sum c_n$ is said to be summable $(R, \log \omega, k)$ or summable (R, k) , to sum s , if

$$R_k(\omega) = \frac{1}{(\log \omega)^k} \sum_{n < \omega} \left(\log \frac{\omega}{n} \right)^k c_n$$

tends to a limit s , as $\omega \rightarrow \infty$. $R_k(\omega)$ is called the $(R, \log \omega, k)$ mean of $\sum c_n$.

Let the Fourier series of a function $f(\theta) \in L$ over $(-\pi, \pi)$ and periodic outside this range with period 2π , be

$$(1.1) \quad f(\theta) \sim \frac{a_0}{2} + \sum_{n=1}^{\infty} (a_n \cos n\theta + b_n \sin n\theta) = \frac{A_0}{2} + \sum_{n=1}^{\infty} A_n(\theta).$$

The series conjugate to the above Fourier series is

$$(1.2) \quad \sum_{n=1}^{\infty} (b_n \cos n\theta - a_n \sin n\theta) = \sum_{n=1}^{\infty} B_n(\theta),$$

and the r th derived series of the above conjugate series is

$$(1.3) \quad \sum_{n=1}^{\infty} \frac{d^r}{d\theta^r} (b_n \cos n\theta - a_n \sin n\theta).$$

Suppose that $P(t)$ is a polynomial of $(r-1)$ th degree in t , such that

$$h(t) = \frac{1}{2t^r} [\{f(x+t) - P(t)\} - (-1)^r \{f(x-t) - P(-t)\}]$$

is integrable (L) over $(-\pi, \pi)$ and is defined by periodicity outside this range with period 2π .

In what follows we shall write

$$\begin{aligned} \psi(t) &= \frac{1}{2} \{f(x+t) - f(x-t)\}, \\ \Psi_\alpha(t) &= \frac{1}{\Gamma(\alpha)} \int_0^t (t-u)^{\alpha-1} \psi(u) du, \quad \alpha > 0, \\ \psi_\alpha(t) &= \frac{\Gamma(\alpha+1)}{t^\alpha} \Psi_\alpha(t), \end{aligned}$$

Received by the editors January 31, 1962.

$$\overline{\Psi}_\alpha(t) = \frac{1}{\Gamma(\alpha)} \int_t^\pi \left(\log \frac{u}{t} \right)^{\alpha-1} \frac{\psi(u)}{u} du, \quad \alpha > 0,$$

$$\overline{\Psi}_\alpha(t) = \Gamma(\alpha + 1) \left(\log \frac{1}{t} \right)^{-\alpha} \overline{\Psi}_\alpha(t),$$

and define $\theta_\alpha(t)$, $\overline{\Theta}_\alpha(t)$, $\overline{\theta}_\alpha(t)$, $\overline{H}_\alpha(t)$ and $\overline{h}_\alpha(t)$ in a similar way.

Also we shall denote by $R_0^\delta(\omega)$ and $R_r^\delta(\omega)$ the (R, δ) means of (1.2) and (1.3), respectively and by $B^\delta(\omega)$ the (R, δ) mean of the conjugate series of $h(t)$.

2. It is known that if $\theta_\alpha(t) \rightarrow 0$ as $t \rightarrow 0$, for $\alpha \geq 0$, then

$$(2.1) \quad C_\alpha(n) \sim \frac{2l}{\pi} \log n,$$

where $C_\alpha(n)$ denotes the (C, α) mean of the conjugate series (1.2) at $\theta = x$. This result for $\alpha = 0$ is the classical result of Lukàcs [2] and for $\alpha > 0$ it has been obtained by Obrechhoff [5].

Considering logarithmic means instead of Cesàro means, we find that if for $p > -1$, $\alpha \geq 0$,

$$\int_t^\pi \frac{|\overline{\theta}_\alpha(u)|}{u} du = o \left\{ \left(\log \frac{1}{t} \right)^{p+1} \right\}, \text{ as } t \rightarrow 0,$$

then

$$R_0^\alpha(\omega) = \frac{2l}{\pi(\alpha + 1)} \log \omega + o \{ (\log \omega)^{p+1} \}, \text{ as } \omega \rightarrow \infty.$$

(See Lemma 2 of the present paper.) For $p = 0$ this gives a result analogous to (2.1), which, although not explicitly mentioned, may also be obtained from a result of Misra [3].

For the first derived series of the conjugate series, Mohanty and Nanda [4] have recently proved the following theorem:

THEOREM A. If

$$\begin{aligned} \int_t^\pi \left| \{ f(x+u) + f(x-u) - 2f(x) \} / \left(4 \sin \frac{u}{2} \right) - l \right| u^{-1} du \\ = o \left(\log \frac{1}{t} \right), \text{ as } t \rightarrow 0, \end{aligned}$$

then $\lim_{n \rightarrow \infty} (\sigma_{2n} - \sigma_n) = l/\pi \log 2$, where σ_n denotes the $(R, \log n, 1)$ mean of the first derived conjugate series, at $\theta = x$.

The object of the present paper is to generalize this theorem to the case of the r th derived series of the conjugate series and to obtain a more refined result by replacing σ_{2n} and σ_n by $R_r^r(\lambda\omega)$ and $R_r^r(\omega)$ respectively and also by replacing the condition of the theorem by a set of weaker conditions. Thus we prove the following theorem which gives a more far-reaching result.

THEOREM 1. *If*

$$\int_t^\pi |h(u) - l| u^{-1} du = o\left\{\left(\log \frac{1}{t}\right)^r\right\}, \text{ as } t \rightarrow 0, \text{ for } r \geq 1,$$

and

$$\int_t^\pi |\tilde{h}_{r-1}(u) - s| u^{-1} du = o\left(\log \frac{1}{t}\right), \text{ as } t \rightarrow 0, \text{ for } r \geq 1,$$

then

$$[R_r^r(\lambda\omega) - R_r^r(\omega)] \sim \frac{2l}{\pi} \frac{r!}{r+1} \log \lambda, \quad \lambda > 1.$$

I am indebted to Professor B. N. Prasad for his kind encouragement during the preparation of this paper.

3. We require the following lemmas for the proof of our theorem:

LEMMA 1 [8]. *If*

$$M_p(t) = \int_0^1 \left(\log \frac{1}{u}\right)^p \sin ut \, du, \quad p > -1,$$

then

$$\frac{d}{dt} \{t M_p(t)\} = p M_{p-1}(t), \quad p > 0.$$

LEMMA 2. *If for* $p > -1$, $\alpha \geq 0$,

$$\int_t^\pi \frac{|\bar{\theta}_\alpha(u)|}{u} du = o\left\{\left(\log \frac{1}{t}\right)^{p+1}\right\}, \text{ as } t \rightarrow 0,$$

then

$$R_0^\alpha(\omega) = \frac{2l}{\pi(\alpha+1)} \log \omega + o\{(\log \omega)^{p+1}\}, \text{ as } \omega \rightarrow \infty.$$

PROOF OF LEMMA 2. Suppose that $h = [\alpha]$. We have by well-known arguments

$$\begin{aligned}
R(\omega) &= \sum_{n < \omega} B_n(x) = \frac{1}{\pi} \int_0^\pi \psi(t) \cot \frac{t}{2} (1 - \cos \omega t) dt + o(1) \\
&= \frac{1}{\pi} \int_0^\pi \theta(t) \cot \frac{t}{2} (1 - \cos \omega t) dt \\
&\quad + \frac{l}{\pi} \int_0^\pi \cot \frac{t}{2} (1 - \cos \omega t) dt + o(1) \\
&= \frac{1}{\pi} \int_0^\pi \theta(t) \left(\cot \frac{t}{2} - \frac{2}{t} \right) (1 - \cos \omega t) dt \\
&\quad + \frac{2}{\pi} \int_0^\pi \theta(t) \frac{1 - \cos \omega t}{t} dt + \frac{2l}{\pi} \log \omega + O(1) \\
&= \frac{2}{\pi} \int_0^\pi \theta(t) \frac{1 - \cos \omega t}{t} dt + \frac{2l}{\pi} \log \omega + O(1). \\
&= \frac{2\omega}{\pi} \int_0^\pi \overline{\Theta}_1(t) \sin \omega t dt + \frac{2l}{\pi} \log \omega + O(1).
\end{aligned}$$

Considering first the case when α is nonintegral, we have

$$\begin{aligned}
R_0^\alpha(\omega) &= \frac{1}{(\log \omega)^\alpha} \sum_{n < \omega} \left(\log \frac{\omega}{n} \right)^\alpha B_n = \frac{\alpha}{(\log \omega)^\alpha} \int_1^\omega \left(\log \frac{\omega}{x} \right)^{\alpha-1} \frac{R(x)}{x} dx \\
&= \frac{2l}{\pi} \frac{\alpha}{(\log \omega)^\alpha} \int_1^\omega \left(\log \frac{\omega}{x} \right)^{\alpha-1} \frac{\log x}{x} dx \\
&\quad + \frac{2\alpha}{\pi} \frac{1}{(\log \omega)^\alpha} \int_1^\omega \left(\log \frac{\omega}{x} \right)^{\alpha-1} dx \int_0^\pi \overline{\Theta}_1(t) \sin x t dt + O(1) \\
&= \frac{2l}{\pi} \frac{\log \omega}{\alpha + 1} + \frac{2\alpha}{\pi (\log \omega)^\alpha} \int_0^\pi \overline{\Theta}_1(t) dt \int_1^\omega \left(\log \frac{\omega}{x} \right)^{\alpha-1} \sin x t dt \\
&\quad + O(1) \\
&= \frac{2l}{\pi} \frac{\log \omega}{\alpha + 1} + \frac{2\alpha\omega}{\pi (\log \omega)^\alpha} \int_0^\pi \overline{\Theta}_1(t) M_{\alpha-1}(\omega t) dt \\
&\quad - \frac{2\alpha\omega}{\pi (\log \omega)^\alpha} \int_0^\pi \overline{\Theta}_1(t) \int_0^{1/\omega} \left(\log \frac{1}{v} \right)^{\alpha-1} \sin(\omega t v) dv + O(1) \\
&= \frac{2l}{\pi} \frac{\log \omega}{\alpha + 1} + \frac{2\alpha\omega}{\pi (\log \omega)^\alpha} \int_0^\pi \frac{\overline{\Theta}_1(t)}{t} t M_{\alpha-1}(\omega t) dt + O(1);
\end{aligned}$$

where

$$M_p(t) = \int_0^1 \left(\log \frac{1}{u} \right)^p \sin ut \, du, \quad p > -1.$$

Thus, integrating by parts,

$$\begin{aligned}
 R_0^\alpha(\omega) &= \frac{2l}{\pi} \frac{\log \omega}{\alpha + 1} + c \frac{\omega}{(\log \omega)^\alpha} \int_0^\pi \bar{\Theta}_{h+1}(t) M_{\alpha-h-1}(\omega t) dt + O(1) \\
 &= \frac{2l}{\pi} \frac{\log \omega}{\alpha + 1} + c_1 \frac{\omega}{(\log \omega)^\alpha} \int_0^\pi M_{\alpha-h-1}(\omega t) \\
 &\quad \cdot \left\{ \int_t^\pi \left(\log \frac{u}{t} \right)^{h-\alpha} \frac{\bar{\Theta}_\alpha(u)}{u} du \right\} dt + O(1) \\
 &= \frac{2l}{\pi} \frac{\log \omega}{\alpha + 1} + \frac{c_1 \omega}{(\log \omega)^\alpha} \int_0^\pi \frac{\bar{\Theta}_\alpha(u)}{u} \\
 (3.1) \quad &\quad \cdot \left\{ \int_0^u \left(\log \frac{u}{t} \right)^{h-\alpha} M_{\alpha-h-1}(\omega t) dt \right\} du + O(1) \\
 &= \frac{2l}{\pi} \frac{\log \omega}{\alpha + 1} + \frac{c_1}{(\log \omega)^\alpha} \\
 &\quad \cdot \left[\left\{ \int_0^{1/\omega} + \int_{1/\omega}^\pi \right\} \frac{\bar{\Theta}_\alpha(u)}{u} (1 - \cos \omega u) du \right] + O(1) \\
 &= \frac{2l}{\pi} \frac{\log \omega}{\alpha + 1} + \frac{c_1}{(\log \omega)^\alpha} [J_1 - J_2] + O(1), \text{ say.}
 \end{aligned}$$

Now

$$\begin{aligned}
 J_1 &= O \left[\int_0^{1/\omega} \frac{|\bar{\Theta}_\alpha(u)|}{u} \omega^2 u^2 du \right] \\
 &= O \left[\omega \int_0^{1/\omega} \frac{|\bar{\Theta}_\alpha(u)|}{u} \cdot u du \right] \\
 (3.2) \quad &= O \left[\omega \left\{ -u \int_u^\pi \frac{|\bar{\Theta}_\alpha(u)|}{u} du \right\}_{u=0}^{u=1/\omega} \right. \\
 &\quad \left. + \omega \left\{ \int_0^{1/\omega} \int_u^\pi \frac{|\bar{\Theta}_\alpha(u)|}{u} du \right\} \right] \\
 &= o[(\log \omega)^{\alpha+p+1}].
 \end{aligned}$$

Again

$$(3.3) \quad J_2 = O \left[\int_{1/\omega}^{\pi} \frac{|\bar{\Theta}_\alpha(u)|}{u} du \right] = o\{(\log \omega)^{\alpha+p+1}\},$$

and the result follows from (3.1), (3.2) and (3.3).

For the case when α is an integer, the proof becomes simpler.

LEMMA 3 [3]. If

$$\int_t^{\pi} |\bar{\Theta}_\alpha(u)| u^{-1} du = o\left\{\left(\log \frac{1}{t}\right)^{\alpha+1}\right\}, \text{ as } t \rightarrow 0, \text{ for } \alpha \geq 0,$$

then

$$\lim_{\omega \rightarrow \infty} [R_0^{\alpha+1}(\lambda\omega) - R_0^{\alpha+1}(\omega)] = \frac{2l}{\pi(\alpha+2)} \log \lambda, \quad \lambda > 1.$$

LEMMA 4 [1].

$$R_r^r(\omega) = (-1)^r \left[\alpha_0 B^r(\omega) + \alpha_1 \frac{B^{r-1}(\omega)}{\log \omega} + \cdots + \alpha_r \frac{B^0(\omega)}{(\log \omega)^r} \right],$$

where $\alpha_0 = (-1)^r r!$ and α_p 's for $p=1, 2, \dots, r$, are constants independent of ω .

LEMMA 5 [7]. Let $V(\omega)$ and $W(\omega)$ be two positive nondecreasing functions of ω . Let

$$U_r(\omega) = \{V(\omega)\}^{1-p/k} \{W(\omega)\}^{p/k}.$$

Then (i) $h^k(\omega) = o\{W(\omega)\}$ and (ii) $h(\omega) = o\{V(\omega)\}$ together imply that $h^p(\omega) = o\{U_p(\omega)\}$ for $0 < p < k$, where $h^p(\omega)$ denotes the (R, p) mean of $h(\omega)$. Also the result remains true if in any one of the conditions (i) and (ii), " o " is replaced by " O ".

4. Proof of Theorem 1. By Lemma 4 we have

$$R_r^r(\omega) = r! B^r(\omega) + (-1)^r \left[\alpha_1 \frac{B^{r-1}(\omega)}{\log \omega} + \cdots + \alpha_r \frac{B^0(\omega)}{(\log \omega)^r} \right].$$

Also by Lemma 2 and by the first condition of the theorem

$$B^0(\omega) = o\{(\log \omega)^r\}, \quad \text{for } r > 1,$$

and by the second condition

$$\begin{aligned} B^{r-1}(\omega) &= \frac{2l}{\pi r} \log \omega + o(\log \omega) \\ &= O(\log \omega). \end{aligned}$$

Now by Lemma 5 we have

$$\begin{aligned} B^1(\omega) &= o\{(\log \omega)^{-1}\}, \\ B^2(\omega) &= o\{(\log \omega)^{-2}\}, \\ &\dots \dots \dots, \\ B^{r-2}(\omega) &= o\{(\log \omega)^2\}. \end{aligned}$$

Thus, supposing that $r > 1$,

$$\begin{aligned} R_r^r(\lambda\omega) - R_r^r(\omega) &= r!\{B^r(\lambda\omega) - B^r(\omega)\} + (-1)^r \alpha_1 \left\{ \frac{B^{r-1}(\lambda\omega)}{\log(\lambda\omega)} - \frac{B^{r-1}(\omega)}{\log \omega} \right\} \\ &\quad + \dots + (-1)^r \alpha_r \left\{ \frac{B^0(\lambda\omega)}{\{\log(\lambda\omega)\}^r} - \frac{B^0(\omega)}{(\log \omega)^r} \right\} \\ &= r!\{B^r(\lambda\omega) - B^r(\omega)\} + (-1)^r \frac{\alpha_1 2l}{\pi r} \left\{ \frac{\log(\lambda\omega)}{\log(\lambda\omega)} - \frac{\log \omega}{\log \omega} \right\} \\ &\quad + o\left\{ \frac{\log(\lambda\omega)}{\log(\lambda\omega)} \right\} + o\left\{ \frac{\log \omega}{\log \omega} \right\} + \dots + o\left\{ \frac{\{\log(\lambda\omega)\}^r}{\{\log(\lambda\omega)\}^r} \right\} + o\left\{ \frac{(\log \omega)^r}{(\log \omega)^r} \right\} \\ &= r!\{B^r(\lambda\omega) - B^r(\omega)\} + o(1) = r! \frac{2l}{\pi} \frac{\log \lambda}{r+1} + o(1) \end{aligned}$$

by Lemma 3. This completes the proof.

REFERENCES

1. S. Kumari, *On the logarithmic summability of the successively derived series of a Fourier series and of its conjugate series*, Indian J. Math. **1** (1959), 87-101.
2. F. Lukács, *Über die Bestimmung des Sprunges einer Funktion aus ihrer Fourierreihe*, J. Reine Angew. Math. **150** (1920), 107-112.
3. M. L. Misra, *On the determination of the jump of a function by its Fourier coefficients*, Quart. J. Math. Oxford Ser. (2), **18** (1947), 147-156.
4. R. Mohanty and M. Nanda, *On the logarithmic mean of the derived conjugate series of a Fourier series*, Proc. Amer. Math. Soc. **7** (1956), 397-400.
5. N. Obrechhoff, *Sur la sommation des séries trigonometriques de Fourier par les moyennes arithmetiques*, Bull. Soc. Math. France **62** (1934), 84-109, (Suit et fin) 167-184.
6. M. Riesz, *Sur les séries de Dirichlet et les séries entières*, C. R. Acad. Sci. Paris **149** (1908), 309-312.
7. ———, *Sur un théorème de la moyenne et ses applications*, Acta Sci. Math. Szeged) **1** (1923), 114-126.
8. F. T. Wang, *On the summability of conjugate Fourier series by Riesz's logarithmic means*, Tôhoku Math. J. **40** (1935), 393-397.

UNIVERSITY OF GORAKHPUR, GORAKHPUR, INDIA