ON DECOMPOSITION INTO STOCHASTICALLY INDEPENDENT COMPONENTS

S. P. LLOYD

1. Introduction. Let $\mathcal{G}$ be an arbitrary class of measurable sets in the probability space $(\Omega, \mathcal{F}, P)$. We denote by $\mathcal{G}^*$ the class of sets which are stochastically independent of $\mathcal{G}$; that is,

$$\mathcal{G}^* = \{H \in \mathcal{F}: P(H \cap G) = P(H)P(G) \text{ for all } G \in \mathcal{G}\}.$$

If $\mathcal{G} \vee \mathcal{G}^*$ denotes the smallest σ -field containing $\mathcal{G}$ and $\mathcal{G}^*$, we say that decomposition (of $\mathcal{F}$ relative to $\mathcal{G}$) holds if $\mathcal{G} \vee \mathcal{G}^* = \mathcal{F}$. In other words, when decomposition holds, every statistical experiment in the class under consideration (every $\mathcal{F}$ set) can be expressed as a combination of (is in the σ -field generated by) experiments from the designated subclass (sets in $\mathcal{G}$) and experiments independent of the subclass (sets in $\mathcal{G}^*$). Examples will be given where decomposition holds and where it fails, and a sufficient condition for decomposition to hold will be established.

2. Some properties of independence. We digress briefly to exhibit some elementary properties of the $*$ operation. We denote by $\emptyset$ the σ -field consisting of sets of probability 0 and their complements. The following relations hold for arbitrary subclasses $\mathcal{G}, \mathcal{H}, \dots$, of $\mathcal{F}$.

- (i) $\mathcal{G}^*$ is closed under the formation of complements, proper differences, and (countable) disjoint unions (but not necessarily intersections),
- (ii) $\emptyset^* = \mathcal{F}, \mathcal{F}^* = \emptyset$,
- (iii) $\mathcal{G}^* \supset \emptyset$,
- (iv) If $\mathcal{G} \subset \mathcal{H}$ then $\mathcal{G}^* \supset \mathcal{H}^*$,
- (1) (v) $(\mathcal{G} \cup \mathcal{H})^* = \mathcal{G}^* \cap \mathcal{H}^*$,
- (vi) $(\mathcal{G} \cap \mathcal{H})^* \supset \mathcal{G}^* \cup \mathcal{H}^*$,
- (vii) $\mathcal{G} \subset \mathcal{G}^{**} (= (\mathcal{G}^*)^*)$,
- (viii) $\mathcal{G}^{***} = \mathcal{G}^*$,
- (ix) $\mathcal{G} \cap \mathcal{G}^* \subset \emptyset$,
- (x) $(\mathcal{G} \cup \mathcal{G}^*)^* = \emptyset, (\mathcal{G} \cup \mathcal{G}^*)^{**} = \mathcal{F}$,
- (xi) $(\mathcal{G} \vee \mathcal{G}^*)^* = \emptyset, (\mathcal{G} \vee \mathcal{G}^*)^{**} = \mathcal{F}$.

Proofs of the above properties are simple and straightforward, and are omitted. It is to be noted that the problem of when $\mathcal{G} \vee \mathcal{G}^* = \mathcal{F}$ holds is subsumed by the problem of when $\mathcal{H}^{**} = \mathcal{H}$ holds, from (xi).

Presented to the Society January 22, 1962 under the title *On decomposition into independent components*; received by the editors February 2, 1962.

3. Conditional cumulative distributions. In all that follows we assume that $\mathcal{G}$ is a sub- σ -field of $\mathcal{F}$.

THEOREM 1. $H \in \mathcal{G}^*$ if and only if $P(H|\mathcal{G}) = P(H)$ with probability 1.

PROOF. $P(H|\mathcal{G})$ is the unique element of $L_\infty(\Omega, \mathcal{G}, P)$ satisfying

$$(2) \quad P(H \cap G) = \int_{\mathcal{G}} P(H|\mathcal{G})_\omega dP(\omega) \quad \text{for all } G \in \mathcal{G}.$$

When $H \in \mathcal{G}^*$ the left-hand side is equal to $P(H)P(G)$, and a solution of (2) is clearly $P(H|\mathcal{G}) = \text{const} = P(H)$ with probability 1. Conversely, if $P(H|\mathcal{G})$ is constant with probability 1 then $P(H \cap G) = P(H)P(G)$ holds for every $G \in \mathcal{G}$, whence $H \in \mathcal{G}^*$.

If x is a real valued random variable we denote by $[x]$ the smallest σ -field with respect to which x is Borel measurable, that is, $[x]$ is the class of sets $\{\omega: x(\omega) \in B\}$ for all linear Borel sets B . The conditional cumulative distribution of x relative to $\mathcal{G}$ will be denoted by

$$F(\lambda, \omega) = P(x \leq \lambda | \mathcal{G})_\omega, \quad -\infty < \lambda < \infty, \omega \in \Omega.$$

We can and do assume that the conditional probability on the right-hand side is determined for each λ in such a way that $F(\cdot, \omega)$ is a bona fide cumulative distribution function for each $\omega \in \Omega$ [1, p. 29].

THEOREM 2. Suppose $F(\lambda, \omega)$, $-\infty < \lambda < \infty$, $\omega \in \Omega$, is continuous on $-\infty < \lambda < \infty$ for each ω in the complement of a set of probability 0. Then the random variable ξ defined by

$$\xi(\omega) = F(x(\omega), \omega), \quad \omega \in \Omega$$

has the properties

- (i) $[\xi] \subset [x] \vee \mathcal{G}$,
- (ii) $[x] \subset [\xi] \vee \mathcal{G}$,
- (iii) $[\xi] \subset \mathcal{G}^*$.

REMARK. P. Lévy is the first to use the independence property of random conditional cumulatives; Lévy gives no proof [2]. Since there seems to be no published proof, and since proof is not as simple as would seem off hand, we give it here.

We need first a measure theoretic lemma. Let $\mathcal{R}$ denote the σ -field of linear Borel sets, and let $(\Omega \times \Lambda, \mathcal{G} \times \mathcal{R})$ denote the product measurable space of $(\Omega, \mathcal{G})$ with $(\Lambda, \mathcal{R})$, where Λ is the real line. If $\Gamma \subset \Omega \times \Lambda$ then Γ_ω denotes the ω section $\Gamma_\omega = \{\lambda \in \Lambda: (\omega, \lambda) \in \Gamma\}$. For any Borel set B , define

$$F(B, \omega) = \int_B F(d\lambda, \omega).$$

LEMMA 1. If $\Gamma \in \mathcal{G} \times \mathcal{R}$ then

$$(3) \quad P(x(\omega) \in \Gamma_\omega) = \int F(\Gamma_\omega, \omega) dP(\omega), \quad \Gamma \in \mathcal{G} \times \mathcal{R}.$$

PROOF. The properties of $F(\cdot, \cdot)$ insure that $\Delta_1(\Gamma) = \int F(\Gamma_\omega, \omega) dP(\omega)$ is a measure on $\mathcal{G} \times \mathcal{R}$. With $\delta: \Omega \rightarrow \Omega \times \Lambda$ defined by $\delta(\omega) = (\omega, x(\omega))$, let Δ_2 be the measure $\Delta_2(\Gamma) = P(\delta^{-1}(\Gamma))$, $\delta^{-1}(\Gamma) \in \mathcal{F}$, noting that $\Delta_2(\Gamma) = P(x(\omega) \in \Gamma_\omega)$. When Γ is a $\mathcal{G} \times \mathcal{R}$ rectangle, $\Gamma = G \times B$, $G \in \mathcal{G}$, $B \in \mathcal{R}$, we have

$$\begin{aligned} \Delta_2(\Gamma) &= P(G \cap \{x \in B\}) \\ &= \int_G P(x \in B \mid \mathcal{G})_\omega dP(\omega) \\ &= \int F(\Gamma_\omega, \omega) dP(\omega) = \Delta_1(\Gamma). \end{aligned}$$

Since measures Δ_1 and Δ_2 coincide on $\mathcal{G} \times \mathcal{R}$ rectangles, they coincide on $\mathcal{G} \times \mathcal{R}$, proving the lemma.

PROOF OF THEOREM 2. Define the pseudoinverses of $F(\cdot, \omega)$ as

$$\begin{aligned} \overline{G}(u, \omega) &= \sup \lambda: F(\lambda, \omega) \leq u \\ \underline{G}(u, \omega) &= \min \lambda: F(\lambda, \omega) \geq u, \\ 0 &< u < 1, \omega \in \Omega. \end{aligned}$$

We have, for all values of the arguments,

$$(4) \quad \begin{aligned} F(\lambda, \omega) &< u \quad \text{if and only if} \quad \lambda < \underline{G}(u, \omega), \\ 0 &< u < 1, \quad -\infty < \lambda < \infty, \quad \omega \in \Omega. \end{aligned}$$

The assumed continuity of $F(\cdot, \omega)$ gives

$$F(\underline{G}(u, \omega), \omega) = F(\overline{G}(u, \omega), \omega) = u, \quad 0 < u < 1,$$

with probability 1.

We will show that the conditional distribution of ξ given $\mathcal{G}$ is the uniform-on-the-unit-interval distribution with probability 1. For each $G \in \mathcal{G}$ and each u , $0 < u < 1$, we have

$$\begin{aligned} P(G \cap \{\xi < u\}) &= P(G \cap \{F(x(\omega), \omega) < u\}) \\ &= P(G \cap \{x(\omega) < \underline{G}(u, \omega)\}) \\ &= P(x(\omega) \in \Gamma_\omega) \end{aligned}$$

where $\Gamma \subset \Omega \times \Lambda$ is the set

$$\Gamma = (G \times \Lambda) \cap \Delta,$$

$$\Delta = \{(\omega, \lambda): \lambda < \underline{G}(u, \omega)\}.$$

It follows from (4) that $\underline{G}(u, \cdot)$ is $\mathfrak{g}$ measurable for each fixed u , $0 < u < 1$; to see that Δ is $\mathfrak{g} \times \mathfrak{R}$ measurable, observe that $\Delta = \lim_n \Delta_n$, with

$$\Delta_n = \bigcup_{j=-\infty}^{\infty} D_{n,j} \times (-\infty, j/2^n)$$

$$D_{n,j} = \{\omega: j/2^n < \underline{G}(u, \omega) \leq (j+1)/2^n\},$$

and the approximating sequence is manifestly $\mathfrak{g} \times \mathfrak{R}$ measurable. Applying Lemma 1, we have, for each $G \in \mathfrak{g}$ and each u , $0 < u < 1$,

$$\begin{aligned} \int_G P(\xi < u | \mathfrak{g})_\omega dP(\omega) &= P(G \cap \{\xi < u\}) = P(x(\omega) \in \Gamma_\omega) \\ &= \int F(\Gamma_\omega, \omega) dP(\omega) \\ &= \int_G F(\underline{G}(u, \omega) - 0, \omega) dP(\omega) \\ &= uP(G), \quad G \in \mathfrak{g}, \quad 0 < u < 1, \end{aligned}$$

whence $P(\xi < u | \mathfrak{g}) = u$ with probability 1, and this for each u , $0 < u < 1$.

That ξ is $[x] \vee \mathfrak{g}$ measurable follows easily from (4). It remains to show that x is $[\xi] \vee \mathfrak{g}$ measurable. Define random variables $\underline{x}$ and $\bar{x}$ by

$$\begin{aligned} \underline{x}(\omega) &= \underline{G}(\xi(\omega), \omega) \\ \bar{x}(\omega) &= \overline{G}(\xi(\omega), \omega), \end{aligned} \quad (\text{where } \omega \in \Omega),$$

observing that $\underline{x}$ is $[\xi] \vee \mathfrak{g}$ measurable, from (4), and that $\underline{x}(\omega) \leq x(\omega) \leq \bar{x}(\omega)$ holds for each $\omega \in \Omega$. We will show that $\underline{x} = \bar{x}$ with probability 1. For each $\omega \in \Omega$ let $I(\omega)$ denote the union of the disjoint nontrivial closed intervals on which $F(\cdot, \omega)$ is constant. Since $\{\omega: \underline{x}(\omega) < \bar{x}(\omega)\} \subset \{\omega: x(\omega) \in I(\omega)\}$, it will suffice to prove that $P(x(\omega) \in I(\omega)) = 0$. If Lemma 1 is applicable we have $P(x(\omega) \in I(\omega)) = \int F(I(\omega), \omega) dP(\omega) = 0$, under the assumption that the $\Omega \times \Lambda$ set Γ whose ω sections are $\Gamma_\omega = I(\omega)$ is $\mathfrak{g} \times \mathfrak{R}$ measurable. To see that this is the case, observe that $\Gamma = \lim_n \Gamma_n$, with

$$\begin{aligned} \Gamma_n &= \bigcup_{-\infty < j < k < \infty} G_{n,j,k} \times [(j-1)/2^n, (k+1)/2^n], \\ G_{n,j,k} &= \{\omega: F((j-1)/2^n, \omega) < F(j/2^n, \omega) \\ &= F(k/2^n, \omega) < F((k+1)/2^n, \omega)\}. \end{aligned}$$

Our main result is the following, to the effect that if there exists one such random variable ξ then there exist many.

THEOREM 3. *Suppose that there exists a random variable independent of $\mathcal{G}$ whose distribution is continuous. Then decomposition holds; that is, $\mathcal{G} \vee \mathcal{G}^* = \mathcal{F}$.*

PROOF. From Theorem 2, we may assume given a random variable ξ with the property $P(\xi < u | \mathcal{G}) = u$ with probability 1, $0 < u < 1$. For arbitrary $A \in \mathcal{F}$ consider the random variable $\eta_A = \chi_A + \xi$, where χ_A denotes the characteristic function of set A . The conditional distribution of η_A has the property

$$P(\lambda_1 \leq \eta_A < \lambda_2 | \mathcal{G}) = P(A \cap \{\lambda_1 - 1 \leq \xi < \lambda_2 - 1\} | \mathcal{G}) \\ + P(A' \cap \{\lambda_1 \leq \xi < \lambda_2\} | \mathcal{G}) \leq 2(\lambda_2 - \lambda_1)$$

with probability 1, implying that some version of $P(\eta_A < \lambda | \mathcal{G})$ is continuous in λ with probability 1. Applying Theorem 2 to η_A , we obtain a random variable ξ_A such that η_A is $[\xi_A] \vee \mathcal{G}$ measurable and such that $[\xi_A] \subset \mathcal{G}^*$. Since $0 < \xi < 1$ with probability 1, set A differs from the set $\{\eta_A > 1\}$ by a set of probability 0, whence $A \in [\eta_A] \subset [\xi_A] \vee \mathcal{G} \subset \mathcal{G}^* \vee \mathcal{G}$ modulo set of probability 0. But A is an arbitrary element of $\mathcal{F}$, so that $\mathcal{F} \subset \mathcal{G} \vee \mathcal{G}^*$, or since the reverse inclusion is trivial, $\mathcal{F} = \mathcal{G} \vee \mathcal{G}^*$.

4. Examples.

Example 1. Decomposition holds. Let $(\Omega, \mathcal{F}, P)$ be atomless, and let $\mathcal{G}$ be purely atomic. That is, there is a finite or countable partition of Ω into disjoint sets G_i , $P(G_i) > 0$, $i = 1, 2, \dots$, and $\mathcal{G}$ is the σ -field generated by the G_i , $i = 1, 2, \dots$. From Theorem 1, $H \in \mathcal{G}^*$ if and only if $P(H \cap G_i) / P(G_i)$ is independent of i . To see that decomposition holds, let $A \in \mathcal{F}$ be an arbitrary measurable set, and define $H_{ii} = A \cap G_i$, $i = 1, 2, \dots$. Since $(\Omega, \mathcal{F}, P)$ is atomless, for each i there exist for all j sets $H_{ij} \subset G_j$ such that $P(H_{ij}) / P(G_j) = P(H_{ii}) / P(G_i)$. With $H_i = \bigcup_j H_{ij}$, we have $H_i \in \mathcal{G}^*$, $i = 1, 2, \dots$, and finally $A = \bigcup_i H_{ii} = \bigcup_i (G_i \cap H_i) \in \mathcal{G} \vee \mathcal{G}^*$. We will prove also that $\mathcal{G} = \mathcal{G}^{**}$ modulo sets of probability 0. Suppose $A \in \mathcal{F}$ does not differ from some $\mathcal{G}$ set by a set of probability 0. Then both $P(A \cap G_i) > 0$ and $P(A' \cap G_i) > 0$ hold for at least one i , say $i = j$. Choose $H_j \subset A \cap G_j$ and $K_j \subset A' \cap G_j$ so that $P(H_j) = P(K_j) > 0$. For all $k \neq j$ choose sets $H_k = K_k \subset G_k$ so that $P(H_k) / P(G_k) = P(H_j) / P(G_j)$. Defining $H = \bigcup_i H_i$, $K = \bigcup_i K_i$, we have $H, K \in \mathcal{G}^*$ and $P(H) = P(K)$. Since $P(A \cap H) - P(A \cap K) = 2P(H_j) > 0$, at least one of H, K is not independent of A , implying $A \notin \mathcal{G}^{**}$. Thus $\mathcal{G}^{**} \subset \mathcal{G}$ modulo sets of probability 0, whence $\mathcal{G}^{**} = \mathcal{G}$ modulo sets of probability 0, from (1.vii).

Example 2. Decomposition fails. Let $\Omega = \{(x, i): 0 \leq x \leq 1, i = 1, 2\}$, and let probability have density $p_i(x)$ at (x, i) relative to linear Borel measure on the two intervals which comprise Ω . Assume that $p_1(x) + p_2(x) = 1$, $0 \leq x \leq 1$. If B_1, B_2 are Borel subsets of the unit interval we denote by $[B_1, B_2]$ the ω set $\{(x, 1): x \in B_1\} \cup \{(x, 2): x \in B_2\}$. Let $\mathcal{G}$ be the sub- σ -field of sets of the form $[B, B]$ for all Borel B . It is readily verified that conditional probabilities relative to $\mathcal{G}$ have the form

$$(5) \quad P([B_1, B_2] \mid \mathcal{G})_{(x, i)} = \chi_{B_1}(x)p_1(x) + \chi_{B_2}(x)p_2(x), \\ 0 \leq x \leq 1, i = 1, 2.$$

For each ω the measure $P(\cdot \mid \mathcal{G})_\omega$ is purely atomic, the measures of the two atoms being $p_1(x), p_2(x)$. It is easy to choose p_1, p_2 in such a way that the only constant functions of the form (5) are 0 and 1, which is to say, $\mathcal{G}^* = \emptyset$. Thus decomposition fails. Note also that $\mathcal{G}^{**} = \mathcal{G}^* = \mathcal{F} \neq \mathcal{G}$.

Example 2 suggests the following conjecture: for decomposition to hold it is sufficient that the conditional probability $P(\cdot \mid \mathcal{G})$ be an atomless measure with probability 1. (This statement makes sense in the Stone representation space; see [3].)

5. Acknowledgment. Example 2 was suggested by Professor Murray Rosenblatt, and the author wishes to express his thanks to Professor Rosenblatt for a number of interesting conversations on this problem.

REFERENCES

1. J. L. Doob, *Stochastic processes*, John Wiley, New York, 1953.
2. P. Lévy, *Théorie de l'addition variables aléatoires*, Gauthier-Villars, Paris, 1937.
3. P. R. Halmos, *On a theorem of Dieudonné*, Proc. Nat. Acad. Sci. U.S.A. **35** (1949), 38-42.

BELL TELEPHONE LABORATORIES,
MURRAY HILL, NEW JERSEY