THE RAYLEIGH FUNCTION

NAND KISHORE

The present paper is a study of a set of symmetric functions of the zeros of $J_\nu(z)$, the Bessel function of the first kind. Let the zeros of $z^{-\nu}J_\nu(z)$ be denoted by $j_{\nu,m}$, $m=1, 2, \dots$, where $|R(j_{\nu,m})| \leq |R(j_{\nu,m+1})|$. And let

$$(1) \quad \sigma_{2n}(\nu) \equiv \sum_{m=1}^{\infty} (j_{\nu,m})^{-2n}, \quad n = 1, 2, \dots$$

We propose to call $\sigma_{2n}(\nu)$ the Rayleigh function of order $2n$. The Rayleigh functions of odd orders are identically zero.

These functions were first used by Euler to determine the three smallest zeros of $J_0(2\sqrt{z})$; and Rayleigh, independently, calculated the smallest positive zero of $J_\nu(z)$ with the aid of these functions. The Rayleigh functions have been the subject of a number of investigations by Cayley, Graeffe, Graf and Gubler, Watson, Kapteyn, Forsyth and others [6, p. 502].

In this paper we shall understand by the Bernoulli and Genocchi numbers, the entities B_n and G_n defined as follows:

$$(2) \quad B_n = \sum_{k=0}^n \binom{n}{k} B_k, \quad n \neq 1,$$

$$B_0 = 1, \quad B_1 = -\frac{1}{2}, \quad B_2 = \frac{1}{6}, \quad B_3 = 0, \quad B_4 = -\frac{1}{30}, \dots$$

$$(3) \quad G_n = 2(1 - 2^n)B_n;$$

(see [4; 3]).

From the well-known formula

$$z^{-1/2}J_{1/2}(z) = \sqrt{\frac{2}{\pi}} \sin z/z,$$

(see [6, p. 54]), the roots of $z^{-1/2}J_{1/2}(z)$ are seen to be $k\pi$, $k=1, 2, \dots$. Hence

$$(4) \quad \sigma_{2n}\left(\frac{1}{2}\right) = \sum_{k=1}^{\infty} (k\pi)^{-2n} = \pi^{-2n} \zeta(2n) = (-1)^{n-1} \frac{2^{2n-1}}{(2n)!} B_{2n}.$$

Similarly, since $z^{1/2}J_{-1/2}(z) = \sqrt{2/\pi} \cos z$, the roots of $z^{1/2}J_{-1/2}(z)$ are

Received by the editors December 14, 1961 and, in revised form, May 14, 1962.

$(k - \frac{1}{2})\pi$, $k = 1, 2, \dots$. Therefore,

$$\begin{aligned} \sigma_{2n} \left(-\frac{1}{2} \right) &= \pi^{-2n} \sum_{k=1}^{\infty} \left(k - \frac{1}{2} \right)^{-2n} = (2^{2n} - 1) \pi^{-2n} \zeta(2n) \\ (5) \qquad &= (-1)^n \frac{2^{2n-2}}{(2n)!} G_{2n}. \end{aligned}$$

A generating function for $\sigma_{2n}(\nu)$ can be obtained. Let $J_\nu(z)$ be expressed as a Weierstrassian product

$$J_\nu(z) = \frac{\left(\frac{1}{2} z \right)^\nu}{\Gamma(\nu + 1)} \prod_{n=1}^{\infty} \left\{ 1 - \frac{z^2}{j_{\nu,n}^2} \right\},$$

(see [6, p. 498]). Differentiating logarithmically with respect to z ,

$$\begin{aligned} J'_\nu(z)/J_\nu(z) &= \nu/z + \sum_{k=1}^{\infty} (-2z)/(j_{\nu,k}^2 - z^2) \\ &= 1/z \left\{ \nu - 2 \sum_{k=1}^{\infty} \sum_{n=1}^{\infty} z^{2n} / j_{\nu,k}^{2n} \right\}, \end{aligned}$$

$$zJ'_\nu(z)/J_\nu(z) = \nu - 2 \sum_{n=1}^{\infty} \sigma_{2n}(\nu) z^{2n},$$

$$(6) \qquad \frac{1}{2} z J_{\nu+1}(z)/J_\nu(z) = \sum_{n=1}^{\infty} \sigma_{2n}(\nu) z^{2n}.$$

Substituting $z = 1$ in (6) yields

$$(7) \qquad \sum_{n=1}^{\infty} \sigma_{2n}(\nu) = \frac{1}{2} J_{\nu+1}(1)/J_\nu(1);$$

$$(8) \qquad \sum_{n=1}^{\infty} \sigma_{2n}(\nu + k) = \frac{1}{2} J_{\nu+k+1}(1)/J_{\nu+k}(1), \qquad k = 0, 1, 2, \dots$$

Setting $\nu = +\frac{1}{2}$ and $-\frac{1}{2}$ in (6), the following are obtained in view of (4) and (5),

$$(9) \qquad \sum_{n=1}^{\infty} (-1)^{n-1} \frac{2^{2n}}{(2n)!} B_{2n} z^{2n} = 1 - z \cot z,$$

$$(10) \qquad \sum_{n=1}^{\infty} (-1)^n \frac{2^{2n-1}}{(2n)!} G_{2n} z^{2n} = z \tan z.$$

Taking the continued fraction representation of the ratio of two

Bessel functions,

$$J_{\nu+1}(z)/J_{\nu}(z) = \frac{z}{2(\nu+1)} - \frac{z^2}{2(\nu+2)} - \dots,$$

(see [6, p. 153]) then substituting $z=1$, and using (7),

$$(11) \quad \sum_{n=1}^{\infty} \sigma_{2n}(\nu) = \frac{1}{2} - \frac{1}{2(\nu+1)} - \frac{1}{2(\nu+2)} - \dots.$$

Set $\nu = \pm \frac{1}{2}$ in (11), then

$$(12) \quad \sum_{n=1}^{\infty} (-1)^{n-1} \frac{2^{2n}}{(2n)!} B_{2n} = \frac{1}{3} - \frac{1}{5} - \frac{1}{7} - \dots$$

$$= 1 - \cot(1) = .3579,$$

$$(13) \quad \sum_{n=1}^{\infty} (-1)^n \frac{2^{2n-1}}{(2n)!} G_{2n} = \frac{1}{1} - \frac{1}{3} - \frac{1}{5} - \dots$$

$$= \tan(1) = 1.5574.$$

A recurrence formula for the functions $\sigma_{2n}(\nu)$ may be derived from (6). Let (6), be written as

$$\frac{1}{2} z J_{\nu+1}(z) = J_{\nu}(z) \sum_{n=1}^{\infty} \sigma_{2n}(\nu) z^{2n}.$$

Substituting the series for $J_{\nu}(z)$ and $J_{\nu+1}(z)$, and identifying the coefficients of z^{2n} on both sides,

$$(14) \quad \sum_{k=1}^n (-1)^{k-1} 4^k (k!)^2 \binom{n}{k} \binom{\nu+n}{k} \sigma_{2k}(\nu) = n.$$

Substitute $\nu = \pm \frac{1}{2}$ in (14), then in view of (4) and (5),

$$(15) \quad \sum_{k=1}^n 2^{2k-1} \binom{2n+1}{k} B_{2k} = n,$$

(see [5, p. 174])

$$(16) \quad - \sum_{k=1}^n 2^{2k-2} \binom{2n}{2k} G_{2k} = n;$$

which are well-known Bernoulli and Genocchi recurrence formulas.

A determinant representation of $\sigma_{2n}(\nu)$ may be given. If in (14) the upper limit of the summation is taken as $1, 2, \dots, n$, then n linear equations involving n functions $\sigma_2(\nu), \sigma_4(\nu), \dots, \sigma_{2n}(\nu)$ are obtained. The determinant Δ of this system is triangular. Hence, the value of Δ is equal to the product of its elements on the main diagonal,

$$\Delta = (-1)^{n(n-1)/2} \cdot 2^{n(n+1)} \cdot n!! \prod_{k=1}^n (\nu + k)^{n-k+1}.$$

Then by Cramer's rule,

$$(17) \quad \sigma_{2n}(\nu) = (-1)^{n-1} \frac{(n-1)!!}{n!} 4^{-n} \prod_{k=1}^n (\nu + k)^{-n+k-1} \cdot D,$$

where,

$$D = \begin{vmatrix} \binom{1}{1} \binom{\nu+1}{1} & 0 & 0 & \dots & 0 & 1 \\ \binom{2}{1} \binom{\nu+2}{1} & \binom{2}{2} \binom{\nu+2}{2} & 0 & \dots & 0 & 2 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \binom{n}{1} \binom{\nu+n}{1} & \binom{n}{2} \binom{\nu+n}{2} & \dots & \binom{n}{n-1} \binom{\nu+n}{n-1} & n \end{vmatrix}.$$

Substitution of $\nu = \pm \frac{1}{2}$ in (17) yields, respectively,

$$(18) \quad B_{2n} = \frac{n!}{(2n+1)!} 2^{-n+1} D_1,$$

$$(19) \quad G_{2n} = -2^{-2n+2} D_2,$$

where

$$D_1 = \begin{vmatrix} \binom{3}{2} & 0 & 0 & \dots & 0 & 1 \\ \binom{5}{2} & \binom{5}{4} & 0 & \dots & 0 & 2 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \binom{2n+1}{2} & \dots & \binom{2n+1}{2n-2} & n \end{vmatrix},$$

$$D_2 = \begin{vmatrix} \binom{2}{2} & 0 & 0 & \cdots & 0 & 1 \\ \binom{4}{2} & \binom{4}{4} & 0 & \cdots & 0 & 2 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ \binom{2n}{2} & \cdots & \binom{2n}{2n-2} & \cdots & n \end{vmatrix}.$$

Other formulas for $\sigma_{2n}(\nu)$ may be derived. Let (6) be written as

$$\begin{aligned} & \{zJ_{r+2}(z)/J_{r+1}(z)\} \{zJ_{r+1}(z)/J_r(z)\} \\ &= 4 \left\{ \sum_{k=1}^{\infty} \sigma_{2k}(\nu+1)z^{2k} \right\} \left\{ \sum_{k=1}^{\infty} \sigma_{2k}(\nu)z^{2k} \right\}. \end{aligned}$$

Then in view of the well-known formula,

$$J_{r+2}(z) = \frac{2(\nu+1)}{z} J_{r+1}(z) - J_r(z),$$

(see [6, p. 45]), the above becomes

$$-\frac{1}{4}z^2 + \frac{1}{2}(\nu+1)zJ_{r+1}(z)/J_r(z) = \left\{ \sum_{k=1}^{\infty} \sigma_{2k}(\nu+1)z^{2k} \right\} \left\{ \sum_{k=1}^{\infty} \sigma_{2k}(\nu)z^{2k} \right\}.$$

Identifying the coefficients of z^{2n} on both sides

$$(20) \quad (\nu+1)\sigma_{2n}(\nu) = \sum_{k=1}^{n-1} \sigma_{2k}(\nu+1)\sigma_{2n-2k}(\nu).$$

Substitution of $\nu = -\frac{1}{2}$, in (20) yields

$$(21) \quad G_{2n} = - \sum_{k=1}^{n-1} \binom{2n}{2k} B_{2k} G_{2n-2k}.$$

Consider the well-known formulas

$$\frac{d}{dz} (z^{-\nu} J_{\nu}(z)) = -z^{-\nu} J_{\nu+1}(z),$$

(see [6, p. 45]),

$$\frac{d}{dz} (z^{\nu+1} J_{\nu+1}(z)) = z^{\nu+1} J_{\nu}(z).$$

It follows that

$$\frac{d}{dz} \left(\frac{z^{\nu+1} J_{\nu+1}(z)}{z^{-\nu} J_{\nu}(z)} \right) = z^{2\nu+1} \frac{(1 + J_{\nu+1}^2(z))}{J_{\nu}^2(z)}$$

That is,

$$\left(\frac{z}{2} \frac{J_{\nu+1}(z)}{J_{\nu}(z)} \right)^2 = -\frac{z^2}{4} + \frac{1}{2} z^{-2\nu+1} \frac{d}{dz} \left(\frac{z}{2} \frac{J_{\nu+1}(z)}{J_{\nu}(z)} z^{2\nu} \right).$$

Substituting (6) in this relation and identifying the coefficients of z^{2n} ,

$$(22) \quad (\nu + n) \sigma_{2n}(\nu) = \sum_{k=1}^{n-1} \sigma_{2k}(\nu) \sigma_{2n-2k}(\nu).$$

Substitute $\nu = \pm \frac{1}{2}$ in (22), then

$$(23) \quad -(1 + 2n) B_{2n} = \sum_{k=1}^{n-1} \binom{2n}{2k} B_{2k} B_{2n-2k},$$

(see [5, p. 66])

$$(24) \quad -2(1 - 2n) G_{2n} = \sum_{k=1}^{n-1} \binom{2n}{2k} G_{2k} G_{2n-2k}.$$

Let (6) be multiplied by itself,

$$J_{\nu+1}^2(z) = 4J_{\nu}^2(z) \left(\sum_{k=1}^{\infty} \sigma_{2k}(\nu) z^{2k-1} \right)^2,$$

then substituting the well-known series for $J_{\nu+1}^2(z)$ and $J_{\nu}^2(z)$, viz.,

$$J_{\nu+1}^2(z) = \sum_{k=0}^{\infty} (-1)^k \frac{\Gamma(2\nu + 3 + 2k)}{k! \Gamma(2\nu + 3 + k) \Gamma^2(\nu + 2 + k)} \left(\frac{z}{2} \right)^{2\nu+2+2k},$$

$$J_{\nu}^2(z) = \sum_{k=0}^{\infty} (-1)^k \frac{\Gamma(2\nu + 1 + 2k)}{k! \Gamma(2\nu + 1 + k) \Gamma^2(\nu + 1 + k)} \left(\frac{z}{2} \right)^{2\nu+2k},$$

(see [6, p. 147]), and identifying the coefficients of z^{2n} on both sides, the following is obtained

$$(25) \quad \sum_{k=1}^n (-1)^{k-1} \binom{2\nu + 2n - 2k}{n - k} (k!)^2 \binom{\nu + n}{k}^2 4^{k+1} \sum_{s=1}^k \sigma_{2s}(\nu) \sigma_{2(k-s+1)}(\nu) = \binom{2\nu + 2n}{n - 1}.$$

In view of (22) this reduces to

$$\begin{aligned}
 \sum_{k=1}^n (-1)^{k-1} 4^k \binom{2\nu+2n-2k}{n-k} \{(k-1)!\}^2 \binom{\nu+n-1}{k-1}^2 (\nu+R) \sigma_{2k}(\nu) \\
 (26) \qquad \qquad \qquad = \frac{2\nu+1}{2\nu+n} \binom{2\nu+2n-2}{n-1}.
 \end{aligned}$$

Substitute $\nu = \pm \frac{1}{2}$ in (26), then

$$(27) \qquad \sum_{k=1}^{n-1} (1+2k) \binom{2n}{2k} B_{2k} = 2n-1,$$

$$(28) \qquad \sum_{k=1}^{n-1} (1-2k) \binom{2n}{2k} G_{2k} = 2(2n-1)G_{2n}.$$

The Rayleigh functions of odd order are zero,

$$\sigma_1(\nu) = 0, \quad \sigma_3(\nu) = 0, \quad \sigma_5(\nu) = 0, \quad \dots$$

Identifying the coefficients of z^2 in (6), an explicit expression for $\sigma_2(\nu)$ is obtained. Then (20) may be used to get explicit expressions for the Rayleigh functions of higher orders. Thus,

$$\begin{aligned}
 \sigma_2(\nu) &= \frac{1}{2^2(\nu+1)}; \\
 \sigma_4(\nu) &= \frac{1}{2^4(\nu+1)^2(\nu+2)}; \\
 \sigma_6(\nu) &= \frac{2}{2^6(\nu+1)^3(\nu+2)(\nu+3)}; \\
 \sigma_8(\nu) &= \frac{5\nu+11}{2^8(\nu+1)^4(\nu+2)^2(\nu+3)(\nu+4)}.
 \end{aligned}$$

The first twelve Rayleigh functions are given in [1, pp. 405-407].

BIBLIOGRAPHY

1. D. H. Lehmer, *Zeros of the Bessel function $J_\nu(x)$* , Math. Comp. 1 (1943-1945), 405-407.
2. ———, *On the maxima and minima of Bernoulli polynomials*, Amer. Math. Monthly 47 (1940), 534-538.
3. ———, Ann. of Math. (2) 36 (1935), 637-649.
4. E. Lucas, *Theorie des nombres*.
5. N. Nielsen, *Traité élémentaire des nombres des Bernoulli*.
6. G. N. Watson, *A treatise on the theory of Bessel functions*, Macmillan, New York, 1944.

UNIVERSITY OF SAN FRANCISCO