A CLASS OF UNIVALENT FUNCTIONS^{1,2}

THOMAS H. MACGREGOR

1. Introduction. The condition $\operatorname{Re} f'(z) > 0$ is known to be sufficient for the univalence of an analytic function in any convex domain. In a recent paper [3] the author investigated the class of functions which satisfy $\operatorname{Re} f'(z) > 0$ for $|z| < 1$ and are normalized by $f(0) = 0$, $f'(0) = 1$. In this paper we study the subclass denoted by F and defined by the condition $|f'(z) - 1| < 1$ for $|z| < 1$. Some of our results have already been proven for the particular functions in F whose coefficients satisfy $\sum_{n=2}^{\infty} n|a_n| \leq 1$. Particular reference should be made of a paper by Schild [5].

2. Distortion theorems. Suppose that $f(z) \in F$. By applying Schwarz's lemma to the function $f'(z) - 1$ we obtain $|f'(z) - 1| \leq |z|$. This gives the estimates $1 - |z| \leq |f'(z)| \leq 1 + |z|$. Bounds for $|f(z)|$ can be obtained from $|f'(z) - 1| \leq |z|$ by integration, as follows. Integrating along the line segment from 0 to z we may write

$$\begin{aligned} f(z) - z &= \int_0^z (f'(s) - 1) ds = z \int_0^1 (f'(tz) - 1) dt. \\ |f(z) - z| &\leq |z| \int_0^1 |f'(tz) - 1| dt \\ &\leq |z| \int_0^1 t |z| dt = (1/2) |z|^2. \end{aligned}$$

From this estimate for $|f(z) - z|$ we immediately obtain

$$|z| - (1/2) |z|^2 \leq |f(z)| \leq |z| + (1/2) |z|^2.$$

Each estimate is precise only for the functions $f(z) = z + a_2 z^2$, where $|a_2| = 1/2$.

The bounds for $|f(z)|$ imply the following theorem.

THEOREM 1. *Each function in F assumes every complex number in the circle $|w| < 1/2$. No values outside of the circle $|w| < 3/2$ are assumed.*

Received by the editors January 22, 1963.

¹ Most of the results in this paper are contained in the author's dissertation, which was completed at the University of Pennsylvania in June, 1961.

² The preparation of this paper was partially supported by the National Science Foundation under the grant NSF-GP-161.

3. An area theorem. If $g(z) = \sum_{n=0}^{\infty} b_n z^n$ is regular for $|z| < 1$ and $|g(z)| \leq 1$ then $\sum_{n=0}^{\infty} |b_n|^2 \leq 1$ [1 p. 7]. Applying this estimate to $f'(z) - 1$ where $f(z) = z + \sum_{n=2}^{\infty} a_n z^n \in F$ we get $\sum_{n=2}^{\infty} n^2 |a_n|^2 \leq 1$. We will use this estimate in the next theorem. It also shows that the coefficients of functions in F satisfy $|a_n| \leq 1/n$ for $n=2, 3, \dots$ and equality for a given n holds only for functions of the form $f(z) = z + a_n z^n$.

THEOREM 2. *The area of the image of $|z| < 1$ under each function in F satisfies $A \leq (3/2)\pi$.*

PROOF. Suppose that $f(z) = z + \sum_{n=2}^{\infty} a_n z^n \in F$ and $0 < r < 1$. Let D_r and D be the images of $|z| < r$ and $|z| < 1$ under $f(z)$ and let A_r and A be the areas of D_r and D . The area of D exists since D is open and bounded. A_r is given by the following well-known formula.

$$A_r = \pi \left\{ r^2 + \sum_{n=2}^{\infty} n |a_n|^2 r^{2n} \right\}.$$

We will prove that

$$(1) \quad A = \pi \left\{ 1 + \sum_{n=2}^{\infty} n |a_n|^2 \right\}.$$

The convergence of this series follows from the fact that $\sum_{n=2}^{\infty} n^2 |a_n|^2$ converges. Let $\{r_n\}$ be any increasing sequence of positive numbers such that $r_n \rightarrow 1$. Then $\{D_{r_n}\}$ is an increasing sequence of sets whose union is D . From the theorem in measure theory $A_{r_n} \rightarrow A$. However the convergence of (1) implies that A_r is continuous for $0 \leq r \leq 1$. Consequently $A_{r_n} \rightarrow A_1$. This proves (1).

From (1) and $\sum_{n=2}^{\infty} n^2 |a_n|^2 \leq 1$, we obtain

$$\begin{aligned} A &= \pi \left\{ 1 + \sum_{n=2}^{\infty} n |a_n|^2 \right\} \\ &\leq \pi \left\{ 1 + (1/2) \sum_{n=2}^{\infty} n^2 |a_n|^2 \right\} \\ &\leq (3/2)\pi. \end{aligned}$$

This proves $A \leq (3/2)\pi$ and $A = (3/2)\pi$ only for the functions $f(z) = z + a_2 z^2$ where $|a_2| = 1/2$.

4. The boundary of the image domain.

THEOREM 3. *Each function in F maps $|z| < 1$ onto a domain whose boundary is a rectifiable Jordan curve.*

PROOF. If $f(z) \in F$ then $|f'(z)| < 2$. From $f(z_2) - f(z_1) = \int_{z_1}^{z_2} f'(z) dz$ we obtain $|f(z_2) - f(z_1)| \leq 2|z_2 - z_1|$. This implies that $f(z)$ is uniformly continuous in $|z| < 1$ and consequently can be extended continuously onto $|z| = 1$. Let C be defined by $w = f(e^{i\theta})$, $0 \leq \theta \leq 2\pi$.

Let us prove that C is rectifiable. This follows easily from the estimate $|f(z_2) - f(z_1)| \leq 2|z_2 - z_1|$ for $|z_1| = |z_2| = 1$. Namely, if $0 = \theta_0 < \theta_1 < \theta_2 < \cdots < \theta_n = 2\pi$ then

$$\sum_{k=1}^n |f(e^{i\theta_k}) - f(e^{i\theta_{k-1}})| \leq 2 \sum_{k=1}^n |e^{i\theta_k} - e^{i\theta_{k-1}}| < 4\pi.$$

Next we show that each function $f(z)$ in F is univalent in $|z| \leq 1$. This follows from the facts: $\operatorname{Re} f'(z) > 0$ for $|z| < 1$ and $f(z)$ is continuous in $|z| \leq 1$. It suffices to prove that $f(z)$ is univalent on $|z| = 1$. Suppose that $z_1 \neq z_2$, $|z_1| = |z_2| = 1$. Let l denote the line segment from z_1 to z_2 . We will consider several points, denoted by $z_1, s_1, z_3, z_4, z_5, s_2, z_2$. These will be distinct points on l and arranged in the order as written. Fix z_4 and then choose z_3 and z_5 such that $\operatorname{Re} f'(z) \geq (1/2) \operatorname{Re} f'(z_4)$ for all points on l between z_3 and z_5 . Then, integrating along l we obtain

$$\begin{aligned} |f(z_2) - f(z_1)| &= \left| \int_{s_1}^{s_2} f'(z) dz \right| \\ &\geq (1/2) |z_5 - z_3| \operatorname{Re} f'(z_4) = a. \end{aligned}$$

With z_3, z_4, z_5 fixed we can choose s_1 and s_2 so close to z_1 and z_2 that

$$|f(z_2) - f(s_2) - (f(z_1) - f(s_1))| < a.$$

Therefore, $f(z_2) - f(z_1) \neq 0$. This proves that $f(z)$ is univalent in $|z| \leq 1$. In particular this shows that C is simple.

THEOREM 4. Suppose $0 < r < 1$. The length of the image of $|z| = r$ under functions in F is maximal for $f_0(z) = z + (1/2)z^2$. Moreover, this length is less than 8.

PROOF. Suppose that $f(z) \in F$ and $L_r(f)$ is the length of the image of $|z| = r$ under $f(z)$. Since $|f'(z) - 1| < 1$ and $f'(z) - 1$ vanishes at $z = 0$, $f'(z)$ is subordinate to $f'_0(z) = 1 + z$ in $|z| < 1$. J. E. Littlewood [2, p. 484, Theorem 2] has shown that if $h(z)$ is subordinate to $H(z)$ in $|z| < 1$ then

$$\int_0^{2\pi} |h(re^{i\theta})|^k d\theta \leq \int_0^{2\pi} |H(re^{i\theta})|^k d\theta,$$

for any $k > 0$. Applying this result to $f'(z)$ for $k = 1$ we prove one part of the theorem, as follows.

$$L_r(f) = r \int_0^{2\pi} |f'(re^{i\theta})| d\theta \leq r \int_0^{2\pi} |f'_0(re^{i\theta})| d\theta = L_r(f_0).$$

$L_r(f) < 8$ will hold if we prove $L_r(f_0) < 8$. If $R \geq 1$ then $f'_0(z) = 1 + z$ is subordinate to $w = 1 + Rz$ in $|z| < 1$. With $R = 1/r$ we again apply Littlewood's inequality.

$$\begin{aligned} L_r(f_0) &< \int_0^{2\pi} |f'_0(re^{i\theta})| d\theta \\ &\leq \int_0^{2\pi} |1 + e^{i\theta}| d\theta \\ &= 2 \int_0^{2\pi} |\cos(\theta/2)| d\theta \\ &= 8. \end{aligned}$$

REMARKS. 1. It seems likely that the length of the boundary of the image of $|z| < 1$ under each function in F satisfies $L \leq 8$. This could not be improved since $L = 8$ for the functions $f(z) = z + a_n z^n$, $|a_n| = 1/n$, $n = 2, 3, \dots$.

2. Littlewood's inequality for $k = 2$ gives another proof of the estimate $A \leq (3/2)\pi$ in Theorem 2.

5. Radii of convexity and starlikeness.

THEOREM 5. *Each function in F maps $|z| < 1/2$ onto a convex domain.*

PROOF. Suppose that $f(z) \in F$. Then $f'(z) - 1 = zg(z)$, where $g(z)$ is regular for $|z| < 1$ and $|g(z)| \leq 1$. For such functions we have the estimate

$$|g'(z)| \leq \frac{1 - |g(z)|^2}{1 - |z|^2}$$

[4, p. 168]. Also,

$$\frac{f''(z)}{f'(z)} = \frac{g(z) + zg'(z)}{1 + zg(z)}.$$

Using the triangle inequalities and then the estimate for $|g'(z)|$ we find that

$$\left| \frac{f''(z)}{f'(z)} \right| \leq \frac{|g(z)| + |z|}{1 - |z|^2}.$$

Since $|g(z)| \leq 1$ we obtain

$$\left| \frac{f''(z)}{f'(z)} \right| \leq \frac{1}{1 - |z|}.$$

The condition $\operatorname{Re}\{zf''(z)/f'(z) + 1\} > 0$ for $|z| < r$ is necessary and sufficient for $f(z)$ to map $|z| < r$ onto a convex domain. This condition is satisfied if $|zf''(z)/f'(z)| < 1$. Thus, $f(z)$ is convex if $|z|/(1 - |z|) < 1$, i.e., if $|z| < 1/2$.

It is not difficult to see that $f(z) = z + a_2 z^2$, $|a_2| = 1/2$, are the only functions in F which are not convex in $|z| < r$ for some $r > 1/2$.

THEOREM 6. *Each function in F maps $|z| < (4/5)^{1/2}$ onto a domain which is starlike with respect to the origin.*

PROOF. The condition $\operatorname{Re}\{zf'(z)/f(z)\} > 0$ for $|z| < r$ is necessary and sufficient for $f(z)$ to be starlike in $|z| < r$. In §2 we proved that $|f'(z) - 1| \leq |z|$ and $|f(z) - z| \leq (1/2)|z|^2$ if $f(z) \in F$. Taking advantage of the geometric location of $f'(z)$ and $f(z)/z$ as given by these inequalities we obtain

$$\begin{aligned} |\arg f'(z)| &\leq \sin^{-1} |z|, \\ \left| \arg \frac{f(z)}{z} \right| &\leq \sin^{-1} (|z|/2). \end{aligned}$$

If $|z| < (4/5)^{1/2}$ then $\sin^{-1}|z| + \sin^{-1}(|z|/2) < \pi/2$. Therefore, for $|z| < (4/5)^{1/2}$ we have

$$\left| \arg \frac{zf'(z)}{f(z)} \right| \leq |\arg f'(z)| + \left| \arg \frac{f(z)}{z} \right| < \pi/2,$$

i.e., $\operatorname{Re}\{zf'(z)/f(z)\} > 0$ for $|z| < (4/5)^{1/2}$.

The estimates used in this proof are precise only for the functions $f(z) = z + a_2 z^2$, where $|a_2| = 1/2$. Since these functions map the whole circle $|z| < 1$ onto a starlike domain, $(4/5)^{1/2}$ is not the radius of starlikeness for the class F .

6. Functions with initial zero coefficients. Some of the results obtained for functions in F can be improved if $f(z)$ has the form $f(z) = z + a_n z^n + a_{n+1} z^{n+1} + \dots$. In this situation we can write $f'(z) - 1 = z^{n-1}g(z)$, where $g(z)$ is regular for $|z| < 1$ and $|g(z)| \leq 1$. With this as the starting point we can argue as in §§2 and 3 to prove the following theorem. The extremal functions for this theorem are $f(z) = z + a_n z^n$, where $|a_n| = 1/n$.

THEOREM 7. Suppose that $f(z) = z + a_n z^n + a_{n+1} z^{n+1} + \dots \in F$. Then $f(z)$ assumes all values in the circle $|w| < 1 - (1/n)$. No values outside of the circle $|w| < 1 + (1/n)$ are assumed. The area of the image domain satisfies $A \leq \pi(1 + 1/n)$.

THEOREM 8. If $f(z) = z + a_n z^n + a_{n+1} z^{n+1} + \dots \in F$ then $f(z)$ maps $|z| < (1/n)^{1/(n-1)}$ onto a convex domain.

PROOF. From $f'(z) - 1 = z^{n-1}g(z)$ it follows that

$$\frac{f''(z)}{f'(z)} = z^{n-2} \frac{(n-1)g(z) + zg'(z)}{1 + z^{n-1}g(z)}.$$

Using the triangle inequalities and then the estimate $|g'(z)| \leq (1 - |g(z)|^2)/(1 - |z|^2)$, we obtain

$$\left| \frac{f''(z)}{f'(z)} \right| \leq \frac{|z|^{n-2}}{1 - |z|^2} \frac{|z| + (n-1)(1 - |z|^2)|g(z)| - |z||g(z)|^2}{1 - |z|^{n-1}|g(z)|}.$$

To estimate the right side of this inequality let us consider the function

$$y = \frac{a + (n-1)(1-a^2)x - ax^2}{1 - a^{n-1}x},$$

$$a = |z|, \quad x = |g(z)|, \quad 0 < a < 1, \quad 0 \leq x \leq 1.$$

$$p = (1 - a^{n-1}x)^2 \frac{dy}{dx} = (n-1)(1-a^2) + a^n - 2ax + a^n x^2$$

$$\frac{dp}{dx} = 2a(a^{n-1}x - 1) < 0.$$

Thus p is decreasing.

$$\begin{aligned} p(x) &\geq p(1) = (n-1)(1-a^2) - 2a(1-a^{n-1}) \\ &= (1-a)[(n-1)(1+a) - 2a(1+a+a^2+\dots+a^{n-2})] \\ &> (1-a)[(n-1)(1+a) - 2a(n-1)] \\ &= (n-1)(1-a)^2 > 0. \end{aligned}$$

Thus y is an increasing function. Therefore

$$y(x) \leq y(1) = \frac{(n-1)(1-a^2)}{1-a^{n-1}},$$

$$\left| \frac{f''(z)}{f'(z)} \right| \leq \frac{(n-1)|z|^{n-2}}{1 - |z|^{n-1}}.$$

If $|zf''(z)/f'(z)| < 1$ then $f(z)$ is convex. Thus, $f(z)$ is convex if $(n-1)|z|^{n-1}/(1-|z|^{n-1}) < 1$, i.e., if $|z| < (1/n)^{1/(n-1)}$.

The functions $f(z) = z + a_n z^n$, $|a_n| = 1/n$, are extremal, i.e., they are the only functions in F of the form $f(z) = z + a_n z^n + a_{n+1} z^{n+1} + \dots$ which are not convex in some circle $|z| < r$, $r > (1/n)^{1/(n-1)}$.

REFERENCES

1. E. Landau, *Darstellung und Begründung einiger neuer Ergebnisse der Funktionentheorie*, Berlin, Springer, 1916.
2. J. E. Littlewood, *On inequalities in the theory of functions*, Proc. London Math. Soc. (2) **23** (1925), 481-519.
3. T. H. MacGregor, *Functions whose derivative has a positive real part*, Trans. Amer. Math. Soc. **104** (1962), 532-537.
4. Z. Nehari, *Conformal mapping*, McGraw-Hill, New York, 1952.
5. A. Schild, *On a class of functions schlicht in the unit circle*, Proc. Amer. Math. Soc. **5** (1954), 115-120.

LAFAYETTE COLLEGE