CONTRACTION OF WALSH FOURIER SERIES

CHINAMI WATARI

The purpose of this note is to prove an analogue for Walsh Fourier series (abbreviated WFS) of a theorem of R. P. Boas [1]. For basic properties of Walsh functions, the reader is referred to N. J. Fine [2].

We begin with some notations and definitions:

The Walsh functions are denoted by $\psi_n(x)$ ($n=0, 1, 2, \dots$) and considered in the "real version."

A complex-valued function $T(z)$ of a complex variable z is called a *contraction* if it satisfies the inequality

$$|T(z_1) - T(z_2)| \leq |z_1 - z_2|.$$

An integrable function $f(x)$ ¹ is *contractible* if it has absolutely convergent WFS and the same is true for $T(f(x))$ where T is any contraction.

Our theorem now reads as follows:

THEOREM. Let $\{\omega_n\}$ be a sequence of non-negative numbers with $\sum_{n=1}^{\infty} \omega_n < \infty$ and

$$(1) \quad \sum_{n=1}^{\infty} n^{-3/2} \left(\sum_{\nu=1}^n \nu^2 \omega_{\nu}^2 \right)^{1/2} + \sum_{n=1}^{\infty} n^{-1/2} \left(\sum_{\nu=n}^{\infty} \omega_{\nu}^2 \right)^{1/2} < \infty.$$

Then an integrable function $f(x)$ with

$$f(x) \sim \sum_{n=0}^{\infty} c_n \psi_n(x), \quad |c_n| \leq \omega_n$$

is contractible.

The proof of this theorem is essentially the same as Boas', the only difference being Lemma 3 below, in which an elementary property of Walsh function is used.

LEMMA 1. For a non-negative sequence $\{\omega_n\}$, (1) is equivalent to

$$\sum_{n=1}^{\infty} 2^{-n/2} \left(\sum_{\nu=1}^{2^n-1} 2^{2k} \omega_{\nu}^2 \right)^{1/2} + \sum_{n=1}^{\infty} 2^{n/2} \left(\sum_{\nu=2^n}^{\infty} \omega_{\nu}^2 \right)^{1/2} < \infty$$

where $k=k(\nu)$ is the integer satisfying $2^k \leq \nu < 2^{k+1}$.

Received by the editors December 27, 1962.

¹ We consider measurable functions with period 1 only; thus the integrability is meant over an interval of length 1, say $[0, 1)$.

PROOF. The equivalence between

$$\sum_{n=1}^{\infty} n^{-1/2} \left(\sum_{\nu=n}^{\infty} \omega_{\nu}^2 \right)^{1/2} < \infty \quad \text{and} \quad \sum_{n=1}^{\infty} 2^{n/2} \left(\sum_{\nu=2^n}^{\infty} \omega_{\nu}^2 \right)^{1/2} < \infty$$

is nothing but Cauchy's condensation theorem. On the other hand,

$$\begin{aligned} & \sum_{n=1}^{\infty} n^{-3/2} \left(\sum_{\nu=1}^n \omega_{\nu}^2 \right)^{1/2} \\ &= \sum_{j=0}^{\infty} \sum_{n=2^j}^{2^{j+1}-1} n^{-3/2} \left(\sum_{\nu=1}^n \omega_{\nu}^2 \right)^{1/2} \leq \sum_{j=0}^{\infty} 2^{-3j/2} \sum_{n=2^j}^{2^{j+1}-1} \left(\sum_{\nu=1}^n \omega_{\nu}^2 \right)^{1/2} \\ &\leq A \sum_{j=0}^{\infty} \sum_{n=2^j}^{2^{j+1}-1} 2^{-3j/2} \left(\sum_{\nu=1}^{2^{j+1}-1} 2^{2k} \omega_{\nu}^2 \right)^{1/2} = A \sum_{j=0}^{\infty} 2^{-j/2} \left(\sum_{\nu=1}^{2^{j+1}-1} 2^{2k} \omega_{\nu}^2 \right)^{1/2} \\ &\leq A \sum_{j=1}^{\infty} 2^{-j/2} \left(\sum_{\nu=1}^{2^j-1} 2^{2k} \omega_{\nu}^2 \right)^{1/2}, \end{aligned}$$

and

$$\begin{aligned} \sum_{j=0}^{\infty} \sum_{n=2^j}^{2^{j+1}-1} n^{-3/2} \left(\sum_{\nu=1}^n \omega_{\nu}^2 \right)^{1/2} &\geq A \sum_{j=0}^{\infty} \sum_{n=2^j}^{2^{j+1}-1} 2^{-3j/2} \left(\sum_{\nu=1}^n 2^{2k} \omega_{\nu}^2 \right)^{1/2} \\ &\geq A \sum_{j=0}^{\infty} 2^{-3j/2} \sum_{n=2^j}^{2^{j+1}-1} \left(\sum_{\nu=1}^{2^j-1} 2^{2k} \omega_{\nu}^2 \right)^{1/2} \\ &= A \sum_{j=0}^{\infty} 2^{-j/2} \left(\sum_{\nu=1}^{2^j-1} 2^{2k} \omega_{\nu}^2 \right)^{1/2}, \quad \text{q.e.d.} \end{aligned}$$

LEMMA 2. For any non-negative sequence $\{a_{\nu}\}$,

$$\sum_{\nu=1}^{\infty} a_{\nu} \leq \sum_{n=1}^{\infty} 2^{-n/2} \left(\sum_{\nu=1}^{2^n-1} 2^{2k} a_{\nu} \right)^{1/2}.$$

PROOF.

$$\begin{aligned} \sum_{\nu=1}^{\infty} a_{\nu} &= \sum_{j=0}^{\infty} \sum_{\nu=2^j}^{2^{j+1}-1} a_{\nu} = \sum_{j=0}^{\infty} 2^j \sum_{\nu=2^j}^{2^{j+1}-1} a_{\nu} \sum_{n=j+1}^{\infty} 2^{-n} \\ &= \sum_{n=1}^{\infty} 2^{-n} \sum_{j=0}^{n-1} 2^j \sum_{\nu=2^j}^{2^{j+1}-1} a_{\nu} = \sum_{n=1}^{\infty} 2^{-n} \sum_{j=0}^{n-1} \sum_{\nu=2^j}^{2^{j+1}-1} 2^k a_{\nu} \\ &= \sum_{n=1}^{\infty} 2^{-n} \sum_{\nu=1}^{2^n-1} 2^k a_{\nu} \leq \sum_{n=1}^{\infty} 2^{-n} \cdot 2^{n/2} \left(\sum_{\nu=1}^{2^n-1} 2^{2k} a_{\nu} \right)^{1/2} \\ &= \sum_{n=1}^{\infty} 2^{-n/2} \left(\sum_{\nu=1}^{2^n-1} 2^{2k} a_{\nu} \right)^{1/2}, \quad \text{q.e.d} \end{aligned}$$

LEMMA 3. Let $f(x) = \sum_{\nu=0}^{\infty} c_{\nu} \psi_{\nu}(x)$ with $\sum_{\nu=0}^{\infty} |c_{\nu}|^2 < \infty$ and let $g(x) \sim \sum_{\nu=0}^{\infty} a_{\nu} \psi_{\nu}(x)$ be a contraction of $f(x)$. Then

$$\sum_{\nu=1}^{2^n-1} 2^{2k} |a_{\nu}|^2 \leq A \sum_{\nu=1}^{2^n-1} 2^{2k} |c_{\nu}|^2 + A 2^n \sum_{\nu=2^n}^{\infty} |c_{\nu}|^2.$$

PROOF. For any h , $0 < h < 1$, choose the integer m so that

$$2^{-m-1} \leq h < 2^{-m}.$$

We have, by the definition of Walsh functions,

$$(2) \quad \psi_{\nu}(h) = -1 \quad (2^m \leq \nu < 2^{m+1}).$$

From Parseval's relation combined with (2), it follows that

$$\begin{aligned} 4 \sum_{\nu=2^m}^{2^{m+1}-1} |a_{\nu}|^2 &\leq \sum_{\nu=2^m}^{\infty} |a_{\nu}|^2 |1 - \psi_{\nu}(h)|^2 = \int_0^1 |g(x+h) - g(x)|^2 dx \\ &\leq \int_0^1 |f(x+h) - f(x)|^2 dx = \sum_{\nu=2^m}^{\infty} |c_{\nu}|^2 |1 - \psi_{\nu}(h)|^2 \\ &\leq 4 \sum_{\nu=2^m}^{\infty} |c_{\nu}|^2. \end{aligned}$$

Thus we have

$$(3) \quad \sum_{\nu=2^m}^{2^{m+1}-1} |a_{\nu}|^2 \leq \sum_{\nu=2^m}^{\infty} |c_{\nu}|^2 = \sum_{j=m}^{\infty} \sum_{\nu=2^j}^{2^{j+1}-1} |c_{\nu}|^2.$$

Since (3) is true for each m , multiplication by 2^{2m} and summation with respect to m give

$$\begin{aligned} \sum_{m=0}^{n-1} 2^{2m} \sum_{\nu=2^m}^{2^{m+1}-1} |a_{\nu}|^2 &= \sum_{m=0}^{n-1} \sum_{\nu=2^m}^{2^{m+1}-1} 2^{2k} |a_{\nu}|^2 \leq \sum_{m=0}^{n-1} 2^{2m} \sum_{j=m}^{\infty} \sum_{\nu=2^j}^{2^{j+1}-1} |c_{\nu}|^2 \\ &= \sum_{m=0}^{n-1} 2^{2m} \sum_{j=m}^{n-1} \sum_{\nu=2^j}^{2^{j+1}-1} |c_{\nu}|^2 + \sum_{m=0}^{n-1} 2^{2m} \sum_{j=n}^{\infty} \sum_{\nu=2^j}^{2^{j+1}-1} |c_{\nu}|^2 \\ &= \sum_{j=0}^{n-1} \sum_{\nu=2^j}^{2^{j+1}-1} |c_{\nu}|^2 \sum_{m=0}^j 2^{2m} + \sum_{j=n}^{\infty} \sum_{\nu=2^j}^{2^{j+1}-1} |c_{\nu}|^2 \sum_{m=0}^{n-1} 2^{2m} \\ &\leq A \sum_{j=0}^{n-1} 2^{2j} \sum_{\nu=2^j}^{2^{j+1}-1} |c_{\nu}|^2 + A \cdot 2^{2n} \sum_{j=n}^{\infty} \sum_{\nu=2^j}^{2^{j+1}-1} |c_{\nu}|^2 \\ &\leq A \sum_{\nu=1}^{2^n-1} 2^{2k} |c_{\nu}|^2 + A \cdot 2^{2n} \sum_{\nu=2^n}^{\infty} |c_{\nu}|^2, \quad \text{q.e.d.} \end{aligned}$$

PROOF OF THEOREM. By Lemmas 2 and 3, we have

$$\begin{aligned}
 \sum_{\nu=1}^{\infty} |a_{\nu}| &\leq A \sum_{n=1}^{\infty} 2^{-n/2} \left(\sum_{\nu=1}^{2^n-1} 2^{2k} |a_{\nu}|^2 \right)^{1/2} \\
 &\leq A \sum_{n=1}^{\infty} 2^{-n/2} \left(\sum_{\nu=1}^{2^n-1} 2^{2k} |c_{\nu}|^2 + 2^{2n} \sum_{\nu=2^n}^{\infty} |c_{\nu}|^2 \right)^{1/2} \\
 &\leq A \sum_{n=1}^{\infty} 2^{-n/2} \left(\sum_{\nu=1}^{2^n-1} 2^{2k} \omega_{\nu}^2 \right)^{1/2} + A \sum_{n=1}^{\infty} 2^{n/2} \left(\sum_{\nu=2^n}^{\infty} \omega_{\nu}^2 \right)^{1/2},
 \end{aligned}$$

which is convergent by Lemma 1 and the assumption of Theorem.

REFERENCES

1. R. P. Boas, *Beurling's test for absolute convergence of Fourier series*, Bull. Amer. Math. Soc. **66** (1960), 24-27.
2. N. J. Fine, *On Walsh functions*, Trans. Amer. Math. Soc. **65** (1949), 372-414.

TÔHOKU UNIVERSITY, SENDAI, JAPAN