REFERENCES

1. E. Bishop, *A generalization of the Stone-Weierstrass theorem*, Pacific J. Math. **11** (1961), 777-783.
2. R. C. Buck, *Bounded continuous functions on a locally compact space*, Michigan Math. J. **5** (1958), 95-104.
3. I. Glicksberg, *Bishop's generalized Stone-Weierstrass theorem for the strict topology*, Proc. Amer. Math. Soc. **14** (1963), 329-333.

UNIVERSITY OF WISCONSIN AND
UNIVERSITY OF CONNECTICUT

WEAK LIMITS OF POWERS OF A CONTRACTION IN HILBERT SPACE¹

S. R. FOGUEL

Let T be an operator, on the Hilbert space H , with $\|T\| \leq 1$. Let

$$H_0 = \{x \mid \text{weak } \lim T^n x = 0\}, \quad H_1 = H_0^\perp.$$

We shall use the facts, proved in [1], that

1. $x \in H_0$ if and only if $\lim (T^n x, x) = 0$ (Theorem 3.1).
2. On H_1 the operator T is unitary (Theorem 1.1).

Given $x \in H$ let $x = x_0 + x_1$, where $x_0 \in H_0$ and $x_1 \in H_1$. The purpose of this note is to find conditions, on the sequence $(T^n x, x)$, that will imply that x_1 is generated by eigenvectors of T . This is related to the notion of mixing for ergodic transformations.

THEOREM 1. *Let y be in the subspace generated by $T^n x$, $T^{*n} x$, $n = 1, 2, \dots$. If $\lim (T^n y, x) = 0$, then $\text{weak } \lim T^n y = 0$.*

PROOF. Since H_0 and H_1 are invariant under T , it is enough to prove the theorem for the case when $x \in H_1$ (and thus also $y \in H_1$). If n_i is any subsequence of the integers and $z = \text{weak } \lim T^{n_i} y$, then z is orthogonal to $T^k x$, $k = 0, \pm 1, \pm 2, \dots$ (since T is unitary on H_1). But z belongs to the subspace generated by $T^{\pm n} x$. Hence, $z = 0$ and, thus, $\text{weak } \lim T^n y = 0$.

COROLLARY. *Let $P(\lambda)$ be a polynomial whose roots of modulus one are $\lambda_1, \dots, \lambda_k$. If $\lim (T^n P(T)x, x) = 0$, then $x = x_0 + x_1$ where: $x_0 \in H_0$, $x_1 = \sum_{i=1}^k z_i$, $T z_i = \lambda_i z_i$.*

PROOF. Let $x = x_0 + x_1$, where $x_0 \in H_0$ and $x_1 \in H_1$. By Theorem 1,

Received by the editors March 25, 1964.

¹ This work was partially supported by NSF Grants GP87 and G14736 at Northwestern University.

$P(T)x \in H_0$ and, thus, $P(T)x_1 \in H_0$ or $P(T)x_1 = 0$. The theorem now follows from the spectral representation of T on H_1 .

Consider the example: $H = L_2([0, 2\pi], \mu)$ and $(Tf)(\theta) = e^{i\theta}f(\theta)$. Then

$$(T^n x, x) = \int_0^{2\pi} e^{in\theta} |f(\theta)|^2 \mu(d\theta)$$

are the Fourier coefficients of $|f|^2 \mu$. Thus if

$$a_0(T^n x, x) + a_1(T^{n+1} x, x) + \cdots + a_p(T^{n+p} x, x) \xrightarrow{n \rightarrow \infty} 0,$$

then

$$f = f_0 + f_1, \quad \int e^{in\theta} |f_0(\theta)|^2 \mu(d\theta) \rightarrow 0$$

and

$$f_1 = 0 \text{ except for the points } \theta_1, \cdots, \theta_k, \text{ which} \\ \text{are the roots of } a_0 + a_1 e^{i\theta} + \cdots + a_p e^{ip\theta} = 0.$$

Consider, for any contraction T , the polynomial $\lambda - 1$:

If $\lim((T^n x, x) - (T^{n-1} x, x)) = 0$, then $\lim(T^n x, x)$ exists. In fact, weak $\lim T^n x$ exists since $x = x_0 + x_1$, where $Tx_1 = x_1$.

In [2] Sucheston proved that if T is given by an ergodic transformation on a measure space and $\lim((T^n x, x) - (T^{n-1} x, x)) = 0$ for n outside of a lacunary sequence and for every $x \in H$, then T is strongly mixing. In our treatment we cannot exclude a lacunary sequence but T can be any contraction, and the condition is assumed for one x only.

Let $g(\theta) = f(e^{i\theta})$ be a continuous function of bounded variation. Let α_k be the Fourier coefficients of g . It is known that the Fourier sums converge uniformly to g .

LEMMA A. Let U be a unitary operator; then $(U^n f(U)x, x) = \sum_{k=-\infty}^{\infty} (U^{n+k} x, x) \alpha_k$.

PROOF. Let $E(\theta)$ be the spectral measure of U , $0 \leq \theta < 2\pi$. Then

$$(U^n f(U)x, x) = \int_0^{2\pi} g(\theta) e^{in\theta} (E(d\theta)x, x) = \sum_{k=-\infty}^{\infty} \alpha_k \int_0^{2\pi} e^{i(n+k)\theta} (E(d\theta)x, x).$$

LEMMA B. The sum $\sum_{k=-\infty}^{\infty} (T^{n+k} x, x) \alpha_k$ is convergent and if $x_0 \in H_0$ then

$$\sum_{k=-\infty}^{\infty} (T^{n+k} x_0, x_0) \alpha_k \xrightarrow{n \rightarrow \infty} 0.$$

PROOF. Let U be the unitary dilation of T , in the sense of [3]. Then

$$\sum_{k=-\infty}^{\infty} (T^{n+k}x, x)\alpha_k = \sum_{k=-\infty}^{\infty} (U^{n+k}x, x)\alpha_k = (U^n f(U)x, x).$$

Now if $\text{weak lim } T^n x_0 = 0$, then $(U^n x_0, x_0) = (T^n x_0, x_0) \rightarrow 0$, hence, $\text{weak lim } U^n x = \text{weak lim } U^{*n} x = 0$. Thus

$$(U^n f(U)x, x) = (f(U)x, U^{*n}x) \rightarrow 0.$$

THEOREM 2. *Let g vanish on at most a countable set. If*

$$\lim_{n \rightarrow \infty} \sum_{k=-\infty}^{\infty} (T^{n+k}x, x)\alpha_k = 0,$$

then $x = x_0 + x_1$, where $x_0 \in H_0$ and x_1 is generated by eigenvectors of T .

PROOF. Let $x = x_0 + x_1$, where $x_0 \in H_0$ and $x_1 \in H_1$. Now

$$\sum_{k=-\infty}^{\infty} (T^{n+k}x, x)\alpha_k = \sum_{k=-\infty}^{\infty} (T^{n+k}x_0, x_0)\alpha_k + \sum_{k=-\infty}^{\infty} (T^{n+k}x_1, x_1)\alpha_k.$$

By Lemma B the first sum tends to zero; thus $\sum_{k=-\infty}^{\infty} (T^{n+k}x_1, x_1)\alpha_k \rightarrow 0$. Hence, by Lemma A and Theorem 1, $f(T)x_1 = 0$. The theorem now follows from the spectral representation of T on H_1 .

BIBLIOGRAPHY

1. S. R. Foguel, *Powers of a contraction in Hilbert space*, Pacific J. Math. **13** (1963), 551-562.
2. L. Sucheston, *An ergodic application of almost convergence sequences*, Duke Math. J. **30** (1963), 417-422.
3. B. Sz-Nagy, *Sur les contractions de l'espace de Hilbert*, Acta Sci. Math. **15** (1953), 87-92.

NORTHWESTERN UNIVERSITY AND
HEBREW UNIVERSITY, JERUSALEM, ISRAEL