

ON DENSE SUBSPACES OF MOORE SPACES

BEN FITZPATRICK, JR.

By a development for a topological space is meant a sequence of collections of basis elements (called regions) satisfying conditions 1, 2, and 3 of Axiom 1 of [4]; by a complete development is meant a development which satisfies condition 4 of this axiom. A (complete) Moore space is a topological space which has a (complete) development. A Moore space is completable if and only if some complete Moore space contains it as a subspace. Clearly if S is completable, then S is dense in some complete Moore space S' , which will be called a completion of S . A development G for S satisfies Axiom C at the point p of S if and only if, for every region R containing p there is an integer n such that every element of G_n which intersects an element of G_n containing p is a subset of R . Younglove proved [7] that every complete development for a complete Moore space S satisfies Axiom C at each point of a dense subset M of S , so that M , regarded as space, satisfies Axiom C and is thus metrizable [5]. In this note, it is shown that every completable Moore space contains a dense metrizable subspace. It is not true, however, that every development of a completable Moore space (even a metrizable space) satisfies Axiom C at some point. It is proved that in order for some development for S to satisfy Axiom C at each point of a dense subspace of S , it is necessary and sufficient that S contain a dense subspace which is strongly screenable in S . Throughout this note certain terminology and theorems from [4] are used without explicit mention.

LEMMA 1. *If S is a topological space, M is a dense subset of S , and U and V are mutually exclusive domains with respect to M , then there exist mutually exclusive domains D_U and D_V in S , containing U and V , respectively.*

THEOREM 1. *If S is a completable Moore space, then every subspace of S contains a dense metrizable subspace.*

PROOF. It suffices to prove that every completable Moore space contains a dense metrizable subspace. Suppose S is a Moore space, and T is a completion of S . Then S is dense in T . Let G denote a complete development for T . Let G'_1 denote a maximal collection of mutually exclusive regions of G_1 . Now every region in T intersects S .

Presented to the Society, August 27, 1964; received by the editors October 24, 1964.

Denote by K_1 a subset of S which intersects each element of G'_1 at only one point; $K_1 \subset G'_1$. Denote by G'_2 a maximal collection of mutually exclusive regions of G_2 whose closures are subsets of elements of G'_1 , which is such that K_1 is a subset of G'_1 . Let K_2 denote a subset of $S \cdot G'_2$ which contains K_1 and intersects each element of G'_2 at only one point. Continue this process, obtaining sequences G' and K , such that K_n is a subset of $S \cdot G'_n$ containing only one point of each element of G'_n , $K_n \subset K_{n+1}$, and G'_n is dense in T . Let $M = K_1 + K_2 + \dots$. Then by a part of the argument for Theorem 164 of [4], M is dense in T . Clearly, then, M is dense in S . For each positive integer n , let H_n denote the collection to which the point set h belongs if and only if, for some integer $m \geq n$ and some element g of G'_m , h is $g \cdot M$. Then $H = (H_1, H_2, H_3, \dots)$ is an Axiom C development for M , so that M is metrizable [5].

THEOREM 2 (RUDIN). *There exists a Moore space which contains no dense metrizable subspace.*

PROOF. Mary Ellen Estill Rudin [6] showed the existence of a Moore space which is not separable but in which every collection of mutually exclusive domains is countable. Consider such a space S . Suppose it contains a dense metrizable subspace M . Let G be a collection of mutually exclusive domains in M . There exists, by Lemma 1, a collection G' of mutually exclusive domains in S covering G , each element of which contains only one element of G . By hypothesis G' is countable. Therefore G is also. Since M is metrizable, and every collection of mutually exclusive domains in M is countable, then M is separable. Denote by K a countable dense subset of M . Then K is a countable dense subset of S . This involves a contradiction.

REMARK. In [2], D. R. Traylor and the author constructed, starting with a Moore space S^0 , a Moore space S^w such that (1) if α is an infinite cardinal, S^0 is α -separable if and only if S^w is, (2) S^0 is normal if and only if S^w is, (3) every open set in S^w contains a copy of S^0 . Thus if S^0 is not metrizable (completable), S^w is not locally metrizable (completable) at any point.

THEOREM 3 (HEATH). *There exists a Moore space S with a dense, topologically complete metrizable subspace M such that no development for S satisfies Axiom C at each point of M .*

PROOF. R. W. Heath [3] constructed an example of a nonmetrizable Moore space S which is the sum of two topologically complete metrizable subspaces S_1 and S_2 each dense in S . Suppose there are developments G and H for S which satisfy Axiom C at each point of

S_1 and S_2 , respectively. There is a development I for S which is a common refinement of G and H . Then I satisfies Axiom C at each point of S_1 and S_2 .

THEOREM 4 (YOUNGLOVE). *There exists a metrizable space with a development not satisfying Axiom C anywhere.*

PROOF. Younglove proved [7] that if the Moore space S is not compact, but is complete, M is a dense inner limiting set in S , and some development for S satisfies Axiom C at each point of M , then some development for S satisfies Axiom C at each point of M and at no other point. Consider the line E^1 ; there exists a development which satisfies Axiom C everywhere; the irrationals, I , form a dense inner limiting set in E^1 ; so there is a development G for E^1 which satisfies Axiom C at each point of I and at no other point of E^1 . Let R denote the rationals, and for each n , let G'_n denote the collection to which g' belongs if and only if g' is $g \cdot R$ for some g in G_n . Then G' is clearly a development for the subspace R . Now if x is in R , there exists a domain D containing x such that, for each positive integer n , there exist regions g_n and h_n of G_n such that x is in g_n , h_n intersects g_n and contains a point not in \bar{D} . In the subspace R , let $D' = D \cdot R$, $g'_n = g_n \cdot R$, and $h'_n = h_n \cdot R$. Since R is dense in E^1 , every domain intersects R . Now g'_n and h'_n belong to G'_n , x is in g'_n , h'_n intersects g'_n , and h'_n contains a point not in D' . Thus G' satisfies Axiom C nowhere.

DEFINITION. The subset M of the topological space S is said to be strongly screenable in S if and only if, for each collection of domains G in S covering M , there exist discrete collections H_1, H_2, \dots of mutually exclusive domains in S such that for each i , H_i is a refinement of G and $\sum H_i^* \supset M$.

THEOREM 5. *In a Moore space S , the following are equivalent:*

- (i) *There exists a development for S which satisfies Axiom C at each point of a dense subset.*
- (ii) *There exists a dense subset of S which is strongly screenable in S .*

PROOF. Suppose (i) is true, and G is such a development, and M is such a dense subspace of S . There exists a maximal subcollection G_1'' of G_1 such that no region of G_1 intersects two regions of G_1'' . Note that if R is in G_1 some region of G_1 intersects both R and $G_1''^*$. Let K_1 denote a subset of M containing only one point of each element of the discrete collection G_1' . Let G_2' denote the set of all regions g of G_2 such that \bar{g} is a subset of $S - G_1'^*$ or of some element of G_1'' . There exists a maximal subcollection G_2'' of G_2' such that no region of G_2 intersects two regions of G_2'' , and such that K_1 is a subset of $G_2''^*$.

Continue this process, obtaining sequences $G_1'', G_2'', G_3'', \dots$, and K_1, K_2, K_3, \dots such that each G_n'' is a discrete subcollection of G_n such that no region of G_n intersects two regions of G_n'' but if R is in G_n some region of G_n intersects both R and $G_n''^*$, and such that K_n is a subset of K_{n+1} and of $G_n''^*$, and of M , and K_n contains only one point of each element of G_n'' . Let $K = K_1 + K_2 + \dots$. Suppose K is not dense in S . There is a region R which does not intersect \bar{K} but which does intersect M at some point x . There exists a positive integer n such that if g is in G_n and contains x , h is in G_n and intersects g , and k is in G_n and intersects h , then k is a subset of R . But some region of G_n contains x and intersects a region of G_n intersecting $G_n''^*$. Let k be a region of G_n'' which is a subset of R . Then $k \cdot K_n$ intersects R . Thus K is dense in S .

Suppose I is a collection of domains covering K . Let H_n denote the set of all regions of G_n'' that are subsets of elements of I . Then H_n is, for each n , a discrete refinement of I . Moreover, $H_1^* + H_2^* + \dots$ contains K , for suppose x is in K_n . Thus for each $m \geq n$, x is in some element of G_m'' . Some region R of G contains x . There is an integer $m \geq n$ such that every region of G_m that contains x is a subset of R . Then x is in H_n^* . Thus K is strongly screenable in S .

Now suppose (ii) is true and M is a dense subset of S that is strongly screenable in S . Let G denote a development for S . Let K_1 denote a maximal subset of M such that no region of G_1 contains two points of K_1 . Let H_1, H_2, \dots be a sequence of discrete refinements of G_1 covering M . Let $K_{1i} = H_i^* \cdot K_1$. Then K_{1i} is a closed and isolated point set (i.e., no point of it is a limit point of it) such that some discrete collection of regions covers it, and each element of that discrete collection contains only one element of it. Similarly, define K_2, K_3, \dots . Thus there is a dense subset K of S which is the sum of countably many point sets K_{ij} such that each is covered by a discrete collection of regions intersecting it at only one point. It suffices to prove that each such point set has Younglove's property Q [7] for if so there is a development G^{ij} which satisfies Axiom C at each point of K_{ij} , and one development refining all of these, so that there is a development satisfying Axiom C at each point of K . So now suppose that L is a closed and isolated point set, that there exists a discrete collection H_1 of regions covering L , and that G is a collection of domains covering S . Let H_2 denote a discrete refinement of H_1 and of G covering L . Let H_3 denote a discrete collection of regions covering L the closure of each of which is a subset of some element of H_2 . For each point x in $S - H_2^*$ there is a region g_x which contains x , is a subset of some element of G and does not intersect H_3^* . Let H_4 be the collection to

which g belongs if and only if, for some x in $S - H_2^*$, g is g_x . Then $H_2 + H_4$ covers S and is locally finite at each point of L . Thus L has Younglove's property Q .

The author does not know whether the existence of a dense metrizable subspace of a Moore space implies conditions (i) and (ii) of Theorem 5, although this is the case if S is normal. To see this, consider a development G for S , and let M denote a dense metrizable subspace. There exists a sequence K such that, for each n , $K_n \subset K_{n+1}$, no region of G_n contains two points of K_n , if x is in M some region of G_n contains x and intersects K_n . Let $L = K_1 + K_2 + \dots$; then L is metrizable and dense in S . Also, each K_i is a closed and isolated subset of L and thus has an open covering of mutually exclusive regions, each containing only one point of K_i . Then by Lemma 1 there is, in S , such an open covering of K_i . Using normality, one can obtain a discrete open covering of each K_i and, as in Theorem 5, K_i has Younglove's property Q ; so that some development satisfies Axiom C at each point of K_i . It follows that some development satisfies Axiom C at each point of L .

REFERENCES

1. R. H. Bing, *Metrisation of topological spaces*, Canad. J. Math. **3** (1951), 175-186.
2. B. Fitzpatrick and D. R. Traylor, *Two theorems on metrizability of Moore spaces*, Pacific J. Math. (to appear).
3. R. W. Heath, *Screenability, pointwise paracompactness and metrisation of Moore spaces*, Canad. J. Math. **16** (1964), 763-770.
4. R. L. Moore, *Foundations of point set theory*, Amer. Math. Soc. Colloq. Publ. Vol. 13, rev. ed., Amer. Math. Soc., Providence, R. I., 1963.
5. ———, *A set of axioms for plane analysis situs*, Fund. Math. **25** (1935), 13-28.
6. Mary Ellen Estill Rudin, *Concerning abstract spaces*, Duke Math. J. **17** (1950), 317-327.
7. J. N. Younglove, *Concerning dense metric subspaces of certain non-metric spaces*, Fund. Math. **48** (1959), 15-25.

AUBURN UNIVERSITY