ON THE DE LA VALLÉE POUSSIN DERIVATIVE¹

C. KASSIMATIS

Let f be a real valued function over D , a subset of the reals. For any integer $n > 0$ and any set $P \subset D$ consisting of $n+1$ distinct points $p_1, \dots, p_{n+1}$, the n th divided difference of f corresponding to P , $V_n[f; P]$, is given by

$$V_n[f; P] = V_n[f; p_1, \dots, p_{n+1}] \\ = \sum_{j=1}^{n+1} f(p_j) / [(u - p_1) \cdots (u - p_{n+1})]'_{u=p_j},$$

the "prime" denoting ordinary differentiation.

Now let D be the interval (a, b) and f continuous over D . Denote by E the set consisting of the $n+1$ points $x_1, \dots, x_{n+1}$. If

$$\lim_{h \rightarrow 0} n! V_n[f; x + x_1 h, \dots, x + x_{n+1} h]$$

exists and is finite, it is called; following Denjoy [1, p. 305], the n th *E-generalized derivative* of f at x , $f_{n,E}(x)$. If $f_{n,E}(x)$ is independent of the choice of $E \subset S$ for a subset S of the reals, we will call it the n th *S-generalized derivative* $f_{n,S}(x)$.

Let f and D be as above. If for $x \in (a, b)$ we have

$$(1) \quad f(x+h) = a_0 + a_1 h + \cdots + a_n h^n / n! + o(h^n),$$

where the numbers $a_i = a_i(x)$, $i=0, \dots, n$, are independent of h , then a_n will be called, following Marcinkiewicz and Zygmund [2, p. 1], the n th *de la Vallée Poussin derivative* of f at x , $f_{(n)}(x)$.

If $f_{(n)}(x)$ exists so does $f_{(i)}(x)$ for $i=1, \dots, n-1$. If f is differentiable n times at x , $f_{(i)}(x)$ exists for $i=1, \dots, n$ and is equal to the ordinary derivative of the corresponding order. The converse is not true for $n \geq 2$. Let $f(x) = e^{-x^{-2}} \sin e^{x^{-2}}$ for $x \neq 0$, and $f(0) = 0$. Then, $f_{(n)}(0) = 0$ for $n=1, 2, 3, \dots$. However, the ordinary derivative of f of order at least 2 does not exist at $x=0$.

Denjoy shows [1, p. 289] that if $f_{(n)}(x)$ exists then $f_{n,S}(x)$ exists for S arbitrary and $f_{n,S}(x) = f_{(n)}(x)$. Since the existence of $f_{(n)}(x)$ implies that of $f_{(i)}(x)$, $i=1, \dots, n-1$, it follows that $f_{i,S_i}(x)$ exists and $f_{i,S_i}(x) = f_{(i)}(x)$ for S_i arbitrary. Further, we can deduce easily from relation (1) that

Received by the editors November 4, 1964.

¹ Research supported by National Research Council, Canada, Grant MCA-31.

$$\lim_{h \rightarrow 0} \{ (y - x - x_i h) V_{n+1}[f: x + x_1 h, \dots, x + x_{n+1} h, y] \} = \theta(x, y - x),$$

$$i = 1, \dots, n + 1,$$

where $\theta(x, y - x) \rightarrow 0$ as $y \rightarrow x$, and $x_1, \dots, x_{n+1}$ is any $(n+1)$ -tuple of points of S .

In this paper we obtain a converse of the above Denjoy theorem.

Let f and E be as above. Let $y_i = x + x_i h$ for $x \in (a, b)$. Set $E_i = \{x_1, \dots, x_i\}$ and $Y_i = \{y_1, \dots, y_i\}$ for $i = 1, \dots, n + 1$.

THEOREM. *If*

(i) $f_{j, E_{j+1}}(x)$ *exists for all* $j = 1, \dots, n$,

(ii) $\lim_{h \rightarrow 0} \{ (y - y_{n+1}) V_{n+1}[f: y_1, \dots, y_{n+1}, y] \} = \theta(x, y - x)$ *where*
 $\theta(x, y - x) \rightarrow 0$ *as* $y \rightarrow x$, *then* $f_{(n)}(x)$ *exists and* $f_{(n)}(x) = f_{n, E}(x)$.

PROOF. By Newton's interpolation formula we have

$$\begin{aligned} f(y) &= f(y_1) + (y - y_1) V_1[f: Y_2] + \dots \\ &\quad + (y - y_1) \dots (y - y_n) V_n[f: Y_{n+1}] \\ &\quad + (y - y_1) \dots (y - y_n) \{ (y - y_{n+1}) V_{n+1}[f: y_1, \dots, y_{n+1}, y] \}. \end{aligned}$$

For $h \rightarrow 0$, we obtain

$$f(y) = f(x) + \dots + (y - x)^n f_{n, E}(x) / n! + (y - x)^n \theta(x, y - x).$$

Taking into account relation (1), the above result implies that $f_{(n)}(x)$ exists and $f_{(n)}(x) = f_{n, E}(x)$.

REMARK. Under the assumption that $f_{n, E}(x)$ exists, the 2nd condition of our theorem is equivalent to

$$\lim_{h \rightarrow 0} V_n[f: y_1, \dots, y_n, y] = (f_{n, E}(x) / n!) + \theta(x, y - x).$$

This follows directly from the relation

$$\begin{aligned} V_n[f: y_1, \dots, y_n, y] - (y - y_{n+1}) V_{n+1}[f: y_1, \dots, y_{n+1}, y] \\ = V_n[f: y_1, \dots, y_{n+1}]. \end{aligned}$$

REFERENCES

1. A. Denjoy, *Sur l'intégration des coefficients différentiels d'ordre supérieur*, Fund. Math. 25 (1935), 273-326.
2. J. Marcinkiewicz and A. Zygmund, *On the differentiability of functions and summability of trigonometrical series*, Fund. Math. 26 (1936), 1-43.

UNIVERSITY OF WINDSOR, ONTARIO, CANADA