

# HADAMARD MATRICES OF ORDER CUBE PLUS ONE

K. GOLDBERG

1. **Result.** Let  $A$  be an Hadamard design of type 1, and let  $(X, Y, Z)$  denote the direct product of matrices  $X, Y$  and  $Z$  (the direction of the product is unimportant here). Later we shall show that

$$B = (I, A, J) + (J, I, A) + (A, J, I) \\ + (A, A, A) + (A, A^T, A^T) + (A^T, A, A^T) + (A^T, A^T, A)$$

is also an Hadamard design of type 1. This construction will prove the theorem:

**THEOREM.** *If there is an Hadamard matrix of type 1 and order  $h$ , then there is an Hadamard matrix of type 1 and order  $(h-1)^3+1$ .*

Williamson [2] shows that there exist Hadamard matrices of type 1 for all orders

$$(1) \quad \left. \begin{array}{l} 2^a (p_1^{a_1} + 1) \cdots (p_r^{a_r} + 1) \\ a, r = 0, 1, 2, \dots, \\ a_1, \dots, a_r = 1, 3, 5, \dots, \end{array} \right\}$$

where each  $p_i$  is a prime congruent to 3 modulo 4.

For example, an Hadamard matrix of type 1 and order 16 exists. By our theorem, one also exists of order  $15^3+1=16 \cdot 211$ , which is not one of the numbers (1). However, another construction of Williamson [2] yields an Hadamard matrix (not of type 1) for this order. The first "new" order is  $39^3+1$ .

2. **Definitions and proof.** Throughout this paper  $I$  and  $J$  denote the identity matrix and the matrix with 1 in every position respectively, of the order required by the context. An  $a, b$  matrix is one in which each element is either  $a$  or  $b$ .

An Hadamard matrix is a 1,  $-1$  matrix  $H$  of order  $h$  such that  $HH^T = hI$ . (Necessarily either  $h=2$  or  $h$  is divisible by 4.) It is of type 1 if  $H+H^T = 2I$ .

An Hadamard design  $A$  is a 0, 1 matrix of order  $h-1$  such that  $AA^T = A^T A = (h/4)I + (h/4-1)J$ . (Necessarily  $AJ = JA = (h/2-1)J$ .) It is of type 1 if  $A+A^T = J-I$ .

If  $H$  is an Hadamard matrix it can be multiplied by generalized permutation matrices to bring it into the form

---

Presented to the Society, April 14, 1960; received by the editors February 12, 1964.

$$\begin{pmatrix} 1 & 1 & \dots & 1 \\ -1 & & & \\ \vdots & J - 2A & & \\ \vdots & & & \\ -1 & & & \end{pmatrix},$$

where  $A$  is an Hadamard design. Then  $H$  is of type 1 if and only if  $A$  is of type 1.

To prove that  $B$  is a 0, 1 matrix we write it in the form

$$B = (I, X) + (A, Y) + (A^T, Z)$$

where  $(P, Q)$  denotes the direct product of  $P$  and  $Q$ , and

$$X = (I, A) + (A, J),$$

$$Y = (I, I + A) + (A, I + A) + (A^T, I + A^T),$$

$$Z = (I, A) + (A, A^T) + (A^T, A).$$

Since  $I, A$  and  $A^T$  are mutually disjoint we need only show that  $X, Y$  and  $Z$  are 0, 1 matrices. And the same reasoning, applied to each, confirms this.

To prove that  $B + B^T = J - I$  we need only note that

$$X + X^T = J - I \quad \text{and} \quad Y + Z^T = Y^T + Z = J.$$

It remains to prove that  $BB^T$  is a linear combination of  $I$  and  $J$ . This is straightforward algebraic manipulation. First

$$BB^T = (I, U) + (A, V + (n - 1)W) + (A^T, V^T + (n - 1)W)$$

where

$$U = XX^T + (2n - 1)(YY^T + ZZ^T),$$

$$V = XZ^T + YX^T + ZY^T,$$

$$W = (Y + Z)(Y + Z)^T$$

and  $n = h/4$ . Evaluating  $U, V$  and  $W$  we obtain

$$U = mI + (m - 1)J,$$

$$V = -(n - 1)(4n - 1)I + 6n(2n - 1)J + (n - 1)(I, J),$$

$$W = (4n - 1)I + (4n - 1)^2J - (I, J),$$

where  $m = ((h - 1)^3 + 1)/4$ . It follows that

$$BB^T = mI + (m - 1)J.$$

This completes the proof of the theorem.

It is clear that three different Hadamard designs of type 1 of the same order can be used in constructing  $B$ . However, all attempts to apply this method using designs of different orders, have failed.

#### REFERENCES

1. A. T. Butson, *Generalized Hadamard matrices*, Proc. Amer. Math. Soc. **13** (1962), 894–898.
2. J. Williamson, *Hadamard's determinant theorem and the sum of four squares*, Duke Math. J. **11** (1944), 65–81.

NATIONAL BUREAU OF STANDARDS