A METRIZATION THEOREM

JANET S. ALLSBROOK

A characterization of metrizable topological spaces in terms of subtopologies is given. First, several terms are defined in order to describe the pertinent subtopologies. Then, the characterization is readily established as a result of a metrization theorem due to Bing [1] and a metrization theorem due to Ceder [2].

DEFINITION 1. Let $(X, \mathfrak{J})$ be a topological space having the property that the intersection of an arbitrary number of open sets is open. Then $(X, \mathfrak{J})$ is called a *fundamental topological space*.

DEFINITION 2. Let $(X, \mathfrak{J})$ be a topological space with a base $\beta = \{B_j: j \in J\}$ such that for each $x \in \bigcap \{B_i: i \in I, I \text{ an arbitrary subset of } J\}$ there exists $B_x \in \beta$ where $x \in B_x \subset \bigcap \{B_i: i \in I\}$. Then, β is called a *fundamental base* for $\mathfrak{J}$.

REMARK. If $(X, \mathfrak{J})$ has a fundamental base for $\mathfrak{J}$, it is a fundamental space.

DEFINITION 3. Let $(X, \mathfrak{J})$ be a topological space. Let β_i , a subcollection of $\mathfrak{J}$, be a base for some topology, $\mathfrak{J}_i$, on X . Consider the subcollection of $\mathfrak{J}$, $\bar{\beta}_i$, where $\bar{\beta}_i = \{X - \bar{T}_i: T_i \in \mathfrak{J}_i \text{ and the closure is with respect to } \mathfrak{J}\}$. Then, $\bar{\beta}_i$ is a base for a topology on X , $\mathfrak{J}/\mathfrak{J}_i$. The topology, $\mathfrak{J}/\mathfrak{J}_i$ is called the *dual topology of $\mathfrak{J}_i$ relative to $\mathfrak{J}$* . $\bar{\beta}_i$ is called the *dual base of β_i* .

We state two metrization theorems, without proof, to be used in the establishment of a new metrization theorem. For other proofs see [1], [2].

A METRIZATION THEOREM OF BING. *A topological space is metrizable if and only if it is T_1 and regular and the topology has a σ -discrete base.*

A METRIZATION THEOREM OF CEDER. *A topological space $(X, \mathfrak{J})$ is metrizable if and only if it is T_1 and regular and the topology has a σ -closure preserving base, $\beta = \bigcup_{i=1}^{\infty} \{B_i\}$, such that for each $x \in X$, and n , a positive integer, $\bigcap \{B: x \in B, B \in B_n\}$ is a neighborhood of x .*

We now state a metrization theorem which characterizes a topological space, $(X, \mathfrak{J})$, as metrizable in terms of a countable family of bases, β_i , $i = 1, 2, \dots$ for subtopologies, $\mathfrak{J}_i$ of $\mathfrak{J}$, and in terms of the dual bases, $\bar{\beta}_i$, $i = 1, 2, \dots$. More precisely:

THEOREM. *A topological space $(X, \mathfrak{J})$ is metrizable if and only if it is T_1 and regular and has a base $\beta = \bigcup_{i=1}^{\infty} \beta_i$ where for each i , β_i is a fun-*

Received by the editors July 29, 1964.

fundamental base for a topology, $\mathfrak{I}_i$, on X , and for each i , the dual base, $\bar{\beta}_i$, is a fundamental base for $\mathfrak{I}/\mathfrak{I}_i$, the dual topology of $\mathfrak{I}_i$ relative to $\mathfrak{I}$.

PROOF. Let $(X, \mathfrak{I})$ be a topological space satisfying the hypotheses above. Then, we shall show that $(X, \mathfrak{I})$ satisfied the hypotheses of Ceder's Theorem. Let $\beta = \bigcup_{i=1}^{\infty} \beta_i$ be the given base for $\mathfrak{I}$. Then, for each $x \in X$, n a positive integer, $\bigcap \{B: x \in B, B \in \beta_n\}$ is a neighborhood of x in $\mathfrak{I}_n \subset \mathfrak{I}$. Also, $X - \bigcup \{\bar{B}_i: B_i \in \beta'_n \subset \beta_n\} = \bigcap \{X - \bar{B}_i\} \in \mathfrak{I}/\mathfrak{I}_i \subset \mathfrak{I}$. Therefore, $\bigcup \{\bar{B}_i\}$ is closed. Hence, $\bigcup \{\bar{B}_i\} = \text{cl}[\bigcup B_i]$. Thus, β_n is a closure preserving family.

Now, let $(X, \mathfrak{I})$ satisfy the hypotheses of Bing's Theorem, and let $\beta = \bigcup_{i=1}^{\infty} \beta_i$ be the σ -discrete base. We define $\beta'_i = \beta_i \cup \{X\}$, $i = 1, 2, \dots$. It is apparent that β'_i is a fundamental base for some subtopology, $\mathfrak{I}'_i$, of $\mathfrak{I}$ for each i . It remains only to show that $\bar{\beta}'_i$ is a fundamental base, also. Now, $\bigcap_{\alpha} \{X - \bar{T}_{\alpha}: T_{\alpha} \in \mathfrak{I}'_i\} = X - \bigcup_{\alpha} \bar{T}_{\alpha} = X - \text{cl}[\bigcup_{\alpha} T_{\alpha}] \in \mathfrak{I}/\mathfrak{I}'_i$, since β'_i is a closure preserving subfamily of $\mathfrak{I}$ implies that $\mathfrak{I}'_i$ is also a closure preserving subfamily of $\mathfrak{I}$. Thus, $\bar{\beta}'_i$ is a fundamental base.

REFERENCES

1. R. H. Bing, *Metrization of topological spaces*, Canad. J. Math. **3** (1951), 175-186.
2. J. G. Ceder, *Some generalizations of metric spaces*, Pacific J. Math. **11**(1961), 105-125.

UNIVERSITY OF FLORIDA