ON CUBIC TRANSFORMATIONS OF ORTHOGONAL POLYNOMIALS

PIERRE BARRUCAND AND DAVID DICKINSON

After investigating generalizations of the quadratic transformation relating Hermite and Laguerre polynomials

$$H_{2n}(x) = (-1)^n 2^{2n} n! L_n^{(-1/2)}(x^2),$$

T. S. Chihara asked [1] for an example of a pair of orthogonal polynomial sets $\{R_n(x)\}$ and $\{S_n(x)\}$ such that $R_{3n}(x) = S_n(z)$ where z is a cubic in x . It is the purpose of this paper to exhibit such examples.

Let $\{S_n(z)\}$ be a set of monic polynomials with $S_n(-z) = (-1)^n S_n(z)$. For this set to be a set of orthogonal polynomials, it is both necessary and sufficient (Shohat [3, pp. 454 and 456]) that there exist a sequence of positive numbers $\{B_n\}$ such that

$$(1) \quad zS_n(z) = S_{n+1}(z) + B_n S_{n-1}(z), \quad n = 1, 2, \dots,$$

or equivalently,

$$(2) \quad (x^3 + \beta x)S_n(x^3 + \beta x) = S_{n+1}(x^3 + \beta x) + B_n S_{n-1}(x^3 + \beta x),$$

$$n = 1, 2, \dots$$

We look for a monic set of orthogonal polynomials $\{R_n(x)\}$ with $R_n(-x) = (-1)^n R_n(x)$ and with a subset $\{R_{3n}(x)\}$ that satisfies a recurrence relation with the same coefficients,

$$(3) \quad (x^3 + \beta x)R_{3n}(x) = R_{3n+3}(x) + B_n R_{3n-3}(x), \quad n = 1, 2, \dots$$

Such a set of orthogonal polynomials exists if (3) can be formed by iterating

$$(4) \quad xR_n(x) = R_{n+1}(x) + b_n R_{n-1}(x), \quad n = 1, 2, \dots,$$

the condition (with $b_n > 0$, $n = 1, 2, \dots$) that is equivalent to the condition that $\{R_n(x)\}$ is an orthogonal set.

Iterating (4), we may write successively

$$xR_{3n}(x) = R_{3n+1}(x) + b_{3n} R_{3n-1}(x),$$

$$x^2 R_{3n}(x) = (R_{3n+2}(x) + b_{3n+1} R_{3n}(x)) + b_{3n} (R_{3n}(x) + b_{3n-1} R_{3n-2}(x)),$$

$$\begin{aligned} x^3 R_{3n}(x) = & (R_{3n+3}(x) + b_{3n+2} R_{3n+1}(x)) \\ & + (b_{3n+1} + b_{3n})(R_{3n+1}(x) + b_{3n} R_{3n-1}(x)) \\ & + b_{3n} b_{3n-1}(R_{3n-1}(x) + b_{3n-2} R_{3n-3}(x)) \end{aligned}$$

and thus

$$\begin{aligned} (x^3 + \beta x) R_{3n}(x) = & R_{3n+3}(x) \\ & + (b_{3n+2} + b_{3n+1} + b_{3n} + \beta) R_{3n+1}(x) \\ & + (b_{3n+1} + b_{3n} + b_{3n-1} + \beta) b_{3n} R_{3n-1}(x) \\ & + b_{3n} b_{3n-1} b_{3n-2} R_{3n-3}(x), \quad n = 1, 2, \dots \end{aligned}$$

is the same recurrence relation as (3) provided β and the b_n are such that

$$\begin{aligned} (5) \quad & b_{3n+2} + b_{3n+1} + b_{3n} + \beta = 0, \quad n = 1, 2, \dots, \\ & b_{3n+1} + b_{3n} + b_{3n-1} + \beta = 0, \quad n = 1, 2, \dots, \\ & b_{3n} b_{3n-1} b_{3n-2} = B_n, \quad n = 1, 2, \dots \end{aligned}$$

Thus, assuming these three conditions (5) on the recurrence relations (1) and (4), the polynomial sets $\{S_n(x^3 + \beta x)\}$ and $\{R_{3n}(x)\}$ satisfy the same recurrence relations ((2) or (3)) and if we further place the initial restrictions

$$(6) \quad S_0(x^3 + \beta x) = R_0(x)$$

and

$$(7) \quad S_1(x^3 + \beta x) = R_3(x),$$

then we will have

$$S_n(x^3 + \beta x) = R_{3n}(x), \quad n = 0, 1, 2, \dots$$

Condition (6) is satisfied because our assumption that polynomial sets are monic implies $S_0(z) = R_0(x) = 1$. Since $S_1(z) = z$, condition (7) becomes $x^3 + \beta x = R_3(x)$ or, because $R_3(x) = xR_2(x) - b_2R_1(x) = x^3 - (b_1 + b_2)x$, we may replace (7) by the condition

$$(8) \quad b_1 + b_2 + \beta = 0.$$

Thus, after combining (8) and the three conditions (5), we have established the

THEOREM. *If $\{S_n(z)\}$ is a set of orthogonal polynomials with $S_0(z) = 1$, $S_1(z) = z$ and*

$$(1) \quad zS_n(z) = S_{n+1}(z) + B_n S_{n-1}(z), \quad n = 1, 2, \dots$$

and if there exist numbers β and b and a sequence of positive numbers $\{b_n\}$ such that the four conditions

$$\begin{aligned} b_1 + b + \beta &= 0, \\ b_{3n-1} &= b, & n &= 1, 2, \dots, \\ b_{3n+1} + b_{3n} + b + \beta &= 0, & n &= 1, 2, \dots, \\ b_{3n} b_{3n-2} &= B_n, & n &= 1, 2, \dots, \end{aligned}$$

are met, then the polynomials $\{R_n(x)\}$ defined by $R_0(x) = 1$, $R_1(x) = x$, and

$$(4) \quad xR_n(x) = R_{n+1}(x) + b_n R_{n-1}(x), \quad n = 1, 2, \dots$$

form an orthogonal set with the property

$$(9) \quad S_n(x^3 + \beta x) = R_{3n}(x), \quad n = 0, 1, \dots$$

After this theorem, one naturally inquires about expressing the polynomials of the sets $\{R_{3n+1}(x)\}$ and $\{R_{3n+2}(x)\}$ in terms of the polynomials of the set $\{S_n(x^3 + \beta x)\}$. We can express $R_{3n+1}(x)$ in terms of $R_{3n}(x)$ and $R_{3n+3}(x)$ by successively combining recurrence relations:

$$\begin{aligned} xR_{3n+1}(x) &= R_{3n+2}(x) + b_{3n+1}R_{3n}(x), \\ x^2R_{3n+1}(x) &= (R_{3n+3}(x) + b_{3n+2}R_{3n+1}(x)) + xb_{3n+1}R_{3n}(x), \\ (x^2 - b_{3n+2})R_{3n+1}(x) &= R_{3n+3}(x) + xb_{3n+1}R_{3n}(x). \end{aligned}$$

Thus we are led to the

COROLLARY. *For polynomial sets meeting the hypotheses of the theorem,*

$$\begin{aligned} R_{3n+1}(x) &= (x^2 - b)^{-1}(S_{n+1}(x^3 + \beta x) + xb_{3n+1}S_n(x^3 + \beta x)), \\ & \quad n = 0, 1, \dots, \\ R_{3n+2}(x) &= (x^2 - b)^{-1}(xS_{n+1}(x^3 + \beta x) + b_{3n+2}b_{3n+1}S_n(x^3 + \beta x)), \\ & \quad n = 0, 1, \dots \end{aligned}$$

A particularly simple set of constants meeting the conditions of the theorem is $b_n = B_n = 1$ for $n > 1$, $\beta = -3$, and $b_1 = B_1 = 2$. For this choice of constants, the two sets $\{R_n(x)\}$ and $\{S_n(x)\}$ are identical and $R_{3n}(x) = R_n(x^3 - 3x)$, $n = 0, 1, \dots$. The recurrence relation

$$\begin{aligned} xR_n(x) &= R_{n+1}(x) + R_{n-1}(x), & n &= 1, 2, \dots, \\ xR_1(x) &= R_1(x) + 2R_0(x) \end{aligned}$$

and the polynomials $\{R_n(x)\}$ when transformed by $R_0(2x) = T_0(x)$ and $R_n(2x) = 2T_n(x)$, $n = 1, 2, \dots$, lead us to the nonmonic set $\{T_n(x)\}$ defined by

$$\begin{aligned} T_0(x) &= 1, & T_1(x) &= x, \\ 2xT_n(x) &= T_{n+1}(x) + T_{n-1}(x), & n &= 1, 2, \dots \end{aligned}$$

But this is a definition of the Tchebichef polynomials of the first kind [2, (10.11)]. Thus we have shown that for the Tchebichef polynomials of the first kind $\{T_n(x)\}$,

$$\begin{aligned} T_{3n}(x) &= T_n(4x^3 - 3x), & n &= 0, 1, \dots, \\ T_{3n+1}(x) &= (4x^2 - 1)^{-1}(T_{n+1}(4x^3 - 3x) + 2xT_n(4x^3 + 3x)), \\ (10) \quad & n &= 0, 1, \dots, \\ T_{3n+2}(x) &= (4x^2 - 1)^{-1}(2xT_{n+1}(4x^3 - 3x) + T_n(4x^3 - 3x)), \\ & n &= 0, 1, \dots \end{aligned}$$

The cubic transformation (10), which may be written as $T_{3n}(x) = T_n(T_3(x))$, is a special case of the transformation of arbitrary degree

$$T_{mn}(x) = T_m(T_n(x)), \quad m, n = 0, 1, \dots,$$

that is an easy consequence of the definition $T_n(\cos \theta) = \cos n\theta$.

For a less trivial example of the theorem, let the $S_n(z)$ be the monic Legendre polynomials defined in terms of the ordinary Legendre polynomials $P_n(z)$ by $2^n(1/2)_n(n!)^{-1}S_n(z) = P_n(z)$, $n = 0, 1, \dots$, or equivalently, defined directly by

$$\begin{aligned} S_0(z) &= 1, \quad S_1(z) = z, \\ xS_n(z) &= S_{n+1}(z) + \frac{n^2}{(2n+1)(2n-1)}S_{n-1}(z), & n &= 1, 2, \dots \end{aligned}$$

Here we have

$$B_n = \frac{n^2}{(2n+1)(2n-1)}, \quad n = 1, 2, \dots,$$

and the remaining numbers of the theorem may be chosen as

$$\begin{aligned} b_{3n} &= \frac{n}{2n+1}, & n &= 1, 2, \dots, \\ b_{3n+1} &= \frac{n+1}{2n+1}, & n &= 0, 1, \dots, \end{aligned}$$

$$b_{3n+2} = b = 1, \quad \beta = -2, \quad n = 0, 1, \dots$$

That these polynomials $\{R_n(x)\}$ where

$$\begin{aligned} 2^n(1/2)_n(n!)^{-1}R_{3n}(x) &= P_n(x^3 - 2x), \\ 2^{n+1}(1/2)_{n+1}((n+1)!)^{-1}R_{3n+1}(x) &= (x^2 - 1)^{-1}(P_{n+1}(x^3 - 2x) \\ &\quad + xP_n(x^3 - 2x)), \\ 2^{n+1}(1/2)_{n+1}((n+1)!)^{-1}R_{3n+2}(x) &= (x^2 - 1)^{-1}(xP_{n+1}(x^3 - 2x) \\ &\quad + P_n(x^3 - 2x)) \end{aligned}$$

are orthogonal over a bounded domain follows from the boundedness of the coefficients in the recurrence relation (4) by a theorem due to Shohat ([3, p. 458]). Since the zeroes of the polynomials of the set $\{P_n(z)\}$ are dense in the domain $|z| \leq 1$, the zeroes of the polynomials of the set $\{P_n(x^3 - 2x)\}$ (which are the same as the zeroes of the set $\{R_{3n}(x)\}$, $(n=0, 1, \dots)$) are dense in the domain D consisting of the points x such that $|x^3 - 2x| \leq 1$. This D is the union of the three disjoint intervals $((-1 - (5)^{1/2})/2, -1)$, $((1 - (5)^{1/2})/2, (-1 + (5)^{1/2})/2)$, and $(1, (1 + (5)^{1/2})/2)$. It follows from Szegő ([4, Theorem 6.1.1]) that this domain D is precisely the "true" domain of orthogonality of this set $\{R_n(x)\}$.

BIBLIOGRAPHY

1. T. S. Chihara, *On kernel polynomials and related systems*, Boll. Un. Mat. Ital. (3) **19** (1964), 451-459.
2. A. Erdélyi et al., *Higher transcendental functions*, McGraw-Hill, New York, 1953.
3. J. Shohat, *The relation of the classical orthogonal polynomials to the polynomials of Appell*, Amer. J. Math. **58** (1936), 453-464.
4. G. Szegő, *Orthogonal polynomials*, Amer. Math. Soc. Colloq. Pub. Vol. 23, Amer. Math. Soc. Providence, R. I., 1959.

INSTITUT BLAISE PASCAL AND
UNIVERSITY OF MASSACHUSETTS