

ON OKAMURA'S UNIQUENESS THEOREM

JOHN H. GEORGE¹

Moyer [1] gives a uniqueness theorem for ordinary differential equations which includes many of the known criteria for uniqueness. It is demonstrated here that Moyer's results can be obtained as a special case of Okamura's theorem [2]. (See Yoshizawa, [3].)

If $f(t, y)$ is defined and continuous on $D: t_0 \leq t \leq t_0 + a, |y - y_0| \leq b$, the problem of uniqueness for a solution $z(t)$ to the initial value problem,

(1) $\dot{y} = f(t, y), y(t_0) = y_0$ is equivalent to the uniqueness of the solution $x(t) \equiv 0$ of

(2) $\dot{x} = F(t, x) = f(t, z(t)) - f(t, z(t) - x)$, where $F(t, 0) = 0$ on D .

THEOREM 1 (OKAMURA [2], SEE ALSO [3]). *The $x(t) \equiv 0$ solution of (2) is unique to the right iff there exists a function $V(t, x)$ defined on D such that (i) $V(t, 0) = 0$, (ii) $V(t, x) > 0, x \neq 0$, (iii) $V(t, x)$ satisfies a local Lipschitz condition with respect to x and*

$$V'(t, x) = \liminf_{h \rightarrow 0} [V(t + h, x + hF(t, x)) - V(t, x)] \leq 0.$$

Or, equivalently, the solution $z(t)$ of (1) is unique if there exists a function $V(t, x)$ such that $V(t, z(t) - y)$ has the properties of Theorem 1. Moyer's results are obtained by defining

$$V(t, z(t) - y) = \exp[2W(t, z(t) - y)].$$

Moyer's Theorem 2.1 can be stated in the following manner.

THEOREM 2. *If there exists a $V(t, x)$ as in Theorem 1 and $z(t), y(t)$ are two distinct solutions of (1) then $z(t) \equiv y(t)$ to the right of t_0 if and only if*

$$\liminf_{t \rightarrow t_0} V(t, z(t) - y(t)) = 0.$$

REMARK. Okamura's theorem shows it may be possible to have uniqueness of solutions of (1) but not assert the existence of the $W(t, r)$ function of Moyer [1].

Received by the editors January 9, 1967.

¹ The author would like to acknowledge the comments of Dr. T. F. Bridgland and Mr. R. Gunderson.

REFERENCES

1. R. D. Moyer, *A general uniqueness theorem*, Proc. Amer. Math. Soc. **17** (1966), 602-607.
2. H. Okamura, *Condition nécessaire et suffisante remplie par les équations différentielles ordinaires sans points de Peano*, Mem. Coll. Sci. Kyoto Univ. A **24** (1942), 21-28.
3. T. Yoshizawa, *Stability theory by Liapunov's second method*, The Mathematical Society of Japan, Maruzen, Tokyo, 1966.

NASA, MARSHALL SPACE FLIGHT CENTER, HUNTSVILLE, ALABAMA