2. ———, *Differential systems with general boundary conditions*, Seminar Reports in Mathematics, University of California Publications in Mathematics 2 (1944), 45–61.

3. R. H. Cole, *General boundary conditions for an ordinary linear differential system*, Trans. Amer. Math. Soc. 111 (1964), 521–550.

4. W. M. Whyburn, *On related difference and differential systems*, Amer. J. Math. 51 (1929), 265–286.

THE VIRGINIA MILITARY INSTITUTE

THE ASYMPTOTIC BEHAVIOR OF A CLASS OF NONLINEAR DIFFERENTIAL EQUATIONS¹

DONALD S. COHEN

We shall study the asymptotic behavior for $t \rightarrow \infty$ of solutions of the following nonlinear differential equation:

$$(1) \quad u'' + f(t, u) = 0.$$

We suppose that $f(t, u)$ satisfies the following conditions:

H-1: $f(t, u)$ is continuous in $D: t \geq 0, -\infty < u < \infty$.

H-2: The derivative f_u exists on D and satisfies $f_u(t, u) > 0$ on D .

H-3: $|f(t, u(t))| \leq f_u(t, 0)|u(t)|$ on D .

An important class of functions $f(t, u)$ which satisfy conditions H-1, 2, 3 is the class of twice continuously differentiable functions $f(t, u)$ which are odd and strictly monotone in u with $f_{uu} \geq 0$ for $u < 0$ and $f_{uu} \leq 0$ for $u > 0$. Nonlinear eigenvalue problems involving this class of functions have been studied extensively by G. H. Pimbley [1].

For the case $f(t, u) = \pm t^\nu u^n$, R. Bellman [2] has given an exhaustive treatment of the asymptotic behavior of proper solutions (i.e., solutions which exist and have continuous derivatives for $t \geq t_0$). For the case $f(t, u) = a(t)u^{2n+1}$ several results on asymptotic behavior exist depending on properties of $a(t)$. References can be found in the papers of P. Waltman [3] and R. A. Moore and Z. Nehari [4].

Our basic result is that there exist solutions of (1) which approach those of $u'' = 0$. More precisely, we prove the

Received by the editors October 5, 1966.

¹ This work was supported by the National Science Foundation under Grant No. GP-4597 at the California Institute of Technology.

THEOREM. Let $f(t, u)$ satisfy H-1, 2, 3, and in addition, suppose that

$$(2) \quad \int_0^\infty t f_\mu(t, 0) dt < \infty.$$

Then, equation (1) has solutions which are asymptotic to $a + bt$ as $t \rightarrow \infty$, where $b \neq 0$.

PROOF. Our proof is essentially a modification of one given by Bellman [2, pp. 114–115] for the linear case $f(t, u) = a(t)u$. Integrate (1) twice between 1 and t to obtain

$$(3) \quad u(t) = c_1 + c_2 t - \int_1^t (t-s)f(s, u(s))ds.$$

From this we obtain, for $t \geq 1$,

$$|u(t)| \leq (|c_1| + |c_2|)t + t \int_1^t |f(s, u(s))| ds.$$

Using properties H-2, 3, we then obtain

$$(4) \quad \frac{|u(t)|}{t} \leq (|c_1| + |c_2|) + \int_1^t s f_u(s, 0) \frac{|u(s)|}{s} ds.$$

We now invoke the fundamental Gronwall inequality which states that if $u, v \geq 0$, $c > 0$ and if $u(x) \leq c + \int_0^x u(\xi)v(\xi)d\xi$, then

$$u(x) \leq c \exp\left(\int_0^x v(\xi)d\xi\right).$$

Applying this to (4), we obtain

$$(5) \quad \frac{|u(t)|}{t} \leq (|c_1| + |c_2|) \exp\left(\int_1^t s f_u(s, 0) ds\right).$$

Finally, (2) and (5) imply

$$(6) \quad |u(t)|/t \leq c_3.$$

Now, by differentiating (3), we obtain

$$(7) \quad u'(t) = c_2 - \int_1^t f(s, u(s))ds.$$

Then, H-3 and (6) imply that

$$\int_1^t |f(s, u(s))| ds \leq \int_1^t f_u(s, 0) |u(s)| ds \leq c_3 \int_1^t s f_u(s, 0) ds.$$

Hence, as $t \rightarrow \infty$, the integral in (7) converges, and therefore u' has a limit as $t \rightarrow \infty$. To ensure that this limit is not zero, we choose $c_2 = 1$ and use as a lower limit, instead of 1, a point t_0 , where t_0 is chosen so that $c_3 \int_{t_0}^{\infty} s f_u(s, 0) ds < 1$. Q.E.D.

REFERENCES

1. G. H. Pimbley, Jr., *A sublinear Sturm-Liouville problem*, J. Math. Mech. **11** (1962), 121-138.
2. R. Bellman, *Stability theory of differential equations*, McGraw-Hill, New York, 1953.
3. P. Waltman, *On the asymptotic behavior of solutions of a nonlinear equation*, Proc. Amer. Math. Soc. **15** (1964), 918-923.
4. R. A. Moore and Z. Nehari, *Nonoscillation theorems for a class of nonlinear differential equations*, Trans. Amer. Math. Soc. **93** (1959), 30-52.

CALIFORNIA INSTITUTE OF TECHNOLOGY